

SEDAR 75 Gulf of Mexico Gray Snapper Operational Assessment Report

Gulf Fisheries Branch
Sustainable Fisheries Division
NOAA Fisheries - Southeast Fisheries Science Center

December 9, 2022

Table of Contents

| | |
|--|----|
| 1. Assessment Process Proceedings | 4 |
| 1.1. Introduction..... | 4 |
| 1.1.1. Workshop Time and Place | 4 |
| 1.1.2. Terms of Reference | 4 |
| 1.1.3 List of Participants | 5 |
| 1.1.4 List of Working Papers and Reference Documents | 6 |
| 2. Data Review and Update | 7 |
| 2.1. Stock Structure and Management Unit | 9 |
| 2.2. Life History Parameters | 9 |
| 2.2.1. Morphometric and Conversion Factors | 9 |
| 2.2.2. Age and Growth | 9 |
| 2.2.3. Natural Mortality | 10 |
| 2.2.4. Maturity..... | 10 |
| 2.2.5. Fecundity..... | 10 |
| 2.3. Fishery-Dependent Data | 10 |
| 2.3.1. Commercial Landings | 10 |
| 2.3.2. Recreational Landings | 11 |
| 2.3.3. Commercial Discards..... | 12 |
| 2.3.4. Recreational Discards | 13 |
| 2.3.5. Commercial Size Composition | 13 |
| 2.3.6. Recreational Size Composition..... | 13 |
| 2.3.7. Commercial Age Composition | 14 |

| | |
|---|----|
| 2.3.8. Recreational Age Composition | 14 |
| 2.3.9. Recreational Catch Per Unit of Effort Indices of Abundance..... | 14 |
| 2.4. Fishery-Independent Surveys..... | 14 |
| 2.4.1. Age-0 and Age-1 Surveys | 14 |
| 2.4.2. SEAMAP Trawl Survey | 15 |
| 2.4.3. Combined Video Survey..... | 15 |
| 2.4.4. Reef Fish Visual Survey | 15 |
| 2.5. Environmental Considerations & Contributions from Stakeholders | 16 |
| 3. Stock Assessment Model Configuration and Methods..... | 16 |
| 3.1. Stock Synthesis Model Configuration | 16 |
| 3.1.1. Initial Conditions | 17 |
| 3.1.2. Temporal Structure | 17 |
| 3.1.3. Spatial Structure..... | 17 |
| 3.1.4. Life History..... | 17 |
| 3.1.5. Recruitment Dynamics..... | 18 |
| 3.1.6. Fleet Structure and Surveys | 18 |
| 3.1.7. Selectivity | 19 |
| 3.1.8. Retention..... | 21 |
| 3.1.9. Landings and Age Compositions | 22 |
| 3.1.10. Discards..... | 22 |
| 3.1.11. Indices | 23 |
| 3.2. Goodness of Fit and Assumed Error Structure | 23 |
| 3.3. Estimated Parameters..... | 23 |
| 3.4. Model Diagnostics | 24 |
| 3.4.1. Residual Analysis..... | 24 |
| 3.4.2. Correlation Analysis | 24 |
| 3.4.3. Profile Likelihoods..... | 24 |
| 3.4.4. Jitter Analysis..... | 24 |
| 3.4.5. Retrospective Analysis..... | 25 |
| 3.4.6. Additional Diagnostics..... | 25 |
| 3.4.7. Sensitivity Runs | 25 |
| 4. Stock Assessment Model - Results..... | 26 |
| 4.1. Estimated Parameters and Derived Quantities..... | 26 |
| 4.2. Fishing Mortality | 26 |

| | |
|---|-----|
| 4.3. Selectivity | 26 |
| 4.4. Retention | 27 |
| 4.5. Recruitment..... | 27 |
| 4.6. Biomass and Abundance Trajectories..... | 28 |
| 4.7. Model Fit and Residual Analysis | 28 |
| 4.7.1. Landings..... | 28 |
| 4.7.2. Discards..... | 28 |
| 4.7.3. Indices | 29 |
| 4.7.4. Length Compositions | 29 |
| 4.7.5. Age Compositions..... | 29 |
| 4.8. Model Diagnostics | 29 |
| 4.8.1. Correlation Analysis | 29 |
| 4.8.2. Profile Likelihoods..... | 30 |
| 4.8.3. Jitter Analysis..... | 30 |
| 4.8.4. Retrospective Analysis..... | 30 |
| 4.8.5. Additional Diagnostics..... | 30 |
| 4.8.6. Bridging analysis | 30 |
| 4.8.6. Sensitivity Model Runs..... | 31 |
| 5. Discussion | 31 |
| 6. Projections..... | 33 |
| 6.1. Introduction..... | 33 |
| 6.2. Projection methods..... | 33 |
| 6.3. Projection results..... | 34 |
| 6.3.1. Biological Reference Points..... | 34 |
| 6.3.2. Stock Status..... | 34 |
| 6.3.3. Overfishing Limits and ABC projections | 34 |
| 7. Acknowledgements..... | 34 |
| 8. Research Recommendations | 35 |
| 9. References | 36 |
| Tables | 38 |
| Figures..... | 137 |
| 10. Appendix..... | 254 |

1. Assessment Process Proceedings

1.1. Introduction

SEDAR75 addressed the stock assessment for Gulf of Mexico Gray Snapper using data inputs through 2020 as implemented in the Stock Synthesis 3 modeling framework (Methot and Wetzel 2013).

1.1.1. Workshop Time and Place

SEDAR75 is an Operational Assessment and the assessment process consisted of two Topical Working Groups, one on Life History and the other on the Shore Mode recreational fleet. These were held from May through July of 2022.

1.1.2. Terms of Reference

The terms of reference approved by the Gulf of Mexico Fishery Management Council (GMFMC) are listed below.

1. Update the approved SEDAR 51 Gulf of Mexico Gray Snapper base model with data through 2019, or for the most recent year for which finalized data are available.
2. Document any changes or corrections made to model and input datasets and provide updated input data tables.
 - a. Evaluate the potential effects of red tide on Gray Snapper, with consideration of past red tide events through 2019.
 - b. Document changes in MRIP data, both pre- and post-recalibration, in terms of the magnitude of changes to catch and effort.
 - c. Re-evaluate gear selectivity, retention, and discards for the recreational shore mode.
 - d. Consider the SEFSC's improved approach for estimating commercial discards.
 - e. Evaluate how to correct for predicted commercial discards above the size limit, given no commercial trip limit or other regulatory limitation.
 - f. Consider SEDAR 51 recommendations for natural mortality (M):
 - i. Set the max age = 28 years
 - ii. Apply a Lorenzen age-specific M vector
 - iii. Consider bounding M between 0.13 and 0.17
 - g. Consider SEDAR 51 recommendations for growth:
 - i. Use all age data regardless of sex
 - ii. Determine whether to predict growth within the model, using the recommended growth parameters as priors, or to use fixed growth parameters
 - h. Consider combining available relevant video indices for the Gulf to allow for the greatest sample size across the longest potential time period. Consider other weighting alternatives for these surveys.
 - i. Consider SEDAR 51 recommendations, and any new information, for reproduction.

- j. Incorporate social and economic information into the stock assessment considerations as practicable.
- 3. Update model parameter estimates and their variances, model uncertainties, estimates of stock status and management benchmarks, and provide the probability of overfishing occurring at specified future harvest and exploitation levels. Provide commercial and recreational landings and discards in pounds and numbers.
 - a. Investigate bounding steepness between 0.81 and 0.99, based on the range considered in the SEDAR 15 Update assessment of mutton snapper.
 - b. Use the following status determination criteria (SDC) proposed in Amendment 51:
 - i. MSY proxy = yield at FMSY or FRebuild (if overfished)
 - ii. MSST = 0.5*BMSY
 - iii. MFMT = FMSY or FRebuild (if overfished)
 - iv. If different SDC are recommended, provide outputs for both the current and recommended SDC.
 - c. Unless otherwise recommended, use the geometric mean of the previous three years' fishing mortality to determine F_{Current} . If an alternative approach is recommended, provide justification and outputs for the current and alternative approach.
 - d. Provide yield streams for the overfishing limit and acceptable biological catch in pounds:
 - i. Annually for five years
 - ii. Under a "constant catch" scenario for both three and five years
 - iii. For the equilibrium yield at F_{MSY} , when estimable
- 4. Develop a stock assessment report to address these TORS and fully document the input data and results of the stock assessment and the comparison model.

1.1.3 List of Participants

Life History Topical Working Group Members

| | |
|------------------------------------|---------------------|
| Francesca Forrestal (Lead analyst) | NMFS Miami |
| Robert Allman | NMFS Panama City |
| Beverly Barnett | NMFS Panama City |
| Chris Bradshaw | FWC |
| Steve Garner | NMFS Panama City |
| Doug Gregory | SSC |
| Dominique Lazarre | FWC, St. Petersburg |
| Heather Moncrief-Cox | NMFS Miami |
| Jim Nance | SSC |
| Steven Scyphers | SSC |
| Katie Siegfried | NMFS Beaufort |
| Molly Stevens | NMFS Miami |
| Ted Switzer | FWC, St. Petersburg |
| Laura Thornton | NMFS Panama City |
| Jim Tolan | TPWD/ SSC |

Shore Mode Topical Working Group Members

Francesca Forrestal (Lead analyst) NMFS Miami
 Chris Bradshaw FWC
 Doug Gregory SSC
 Dominique Lazarre FWC, St. Petersburg
 Vivian Matter NMFS
 Jim Nance SSC
 Matt Nuttall NMFS
 Beverly Sauls FWC
 Eric Schmidt Industry Representative
 Steven Scyphers SSC
 Katie Siegfried NMFS Beaufort
 Molly Stevens NMFS Miami
 Ted Switzer FWC, St. Petersburg
 Jim Tolan TPWD/ SSC

Attendees

Skyler Sagarese NMFS
 Julie Vecchio FWC
 Yuying Zhang FIU

Staff

Julie Neer SEDAR
 Alisha Gray NOAA SERO
 Ryan Rindone GMFMC Staff

1.1.4 List of Working Papers and Reference Documents

| Document # | Title | Authors | Date Submitted |
|--|---|---|----------------|
| Documents Prepared for the Operational Assessment | | | |
| SEDAR75-WP-01 | Gray Snapper Abundance Indices from Inshore Surveys of Northeastern Gulf of Mexico estuaries (1996-2020) | Kerry E. Flaherty-Walia, Amanda J. Tyler-Jedlund, and Theodore S. Switzer | 29 April 2022 |
| SEDAR75-WP-02 | General Recreational Survey Data for Gray Snapper in the Gulf of Mexico | Matthew A. Nuttall | 15 June 2022 |
| SEDAR75-WP-03 | Gray Snapper Abundance Indices from SEAMAP Groundfish Surveys in the Northern Gulf of Mexico | Adam G. Pollack and David S. Hanisko | 7 June 2022 |
| SEDAR75-WP-04 | Electronic Monitoring Documentation of Gray Snapper (<i>Lutjanus griseus</i>) Catches in the Eastern Gulf of Mexico | Max Lee, Carole Neidig, and Katie Harrington | 16 June 2022 |

| | | | |
|---------------------------------------|---|--|--------------|
| | Commercial Reef Fish Bottom Longline Fishery | | |
| SEDAR75-WP-05 | Life History Data for SEDAR 75 Gulf of Mexico Gray Snapper | Steven Garner, Laura Thornton, Heather Moncrief-Cox and Robert Allman | 8 July 2022 |
| SEDAR75-WP-06 | Standardized Reef Visual Census index for Gray Snapper, <i>Lutjanus griseus</i> , for the Florida reef track from the Florida Keys and Dry Tortugas for 1997-2018 | Robert G. Muller | 11 July 2022 |
| SEDAR75-WP-07 | Gulf of Mexico Gray Snapper (<i>Lutjanus griseus</i>) Commercial and Recreational Landings Length and Age Compositions | Molly H. Stevens | 21 July 2022 |
| | | | |
| Final Stock Assessment Reports | | | |
| SEDAR75-SAR1 | Gulf of Mexico Gray Snapper | | |
| | | | |
| Reference Documents | | | |
| SEDAR75-RD01 | Characterizing gray snapper (<i>Lutjanus griseus</i>) life history in the northcentral Gulf of Mexico: age and growth, mortality, and reproduction | Edward S. M. Kim | |
| SEDAR75-RD02 | Social equity in shore-based fisheries: identifying and understanding barriers to access | Kelsi L. Furman, Sharon L. Harlan, Luiz Barbieri, and Steven B. Scyphers | |
| | | | |

2. Data Review and Update

A variety of data sources were used in the SEDAR75 Operational Assessment. Where practicable, the SEDAR75 base model used the same data sets as the SEDAR51 Benchmark model with an updated time series. However, there were a few new or revised datasets provided for the SEDAR75 data call. These included the National Marine Fisheries Service's (NMFS) Marine Recreational Information Program (MRIP) Fishing Effort Survey (FES) catch and discard time series, improved commercial fishery discard estimates, improved Southeast Region Headboat Survey discard proxy estimates, updated growth curve given newly available age-length pairs, updated information on maturity based on recent findings, and a new fishery-independent combined video survey with associated length compositions. Length composition data for two fishing independent surveys were provided, the SEAMAP Trawl Survey and the Reef Fish Visual Survey. Age compositions were provided and were able to be used in SEDAR75. These new data series were considered because they had not previously been available for the SEDAR 51 Benchmark assessment or represented improved data inputs for use in the assessment. The Commercial Vertical Line Index used in SEDAR51 was not used in this assessment as the data would cover the post-IFQ period and the index could not be extended.

The index in SEDAR 51 was mirrored to the non-Monroe Commercial Handline fleet in the SEDAR 51 assessment model and this fleet was not used in this assessment as the fleet structure was modified (**see Section 2.3.1**). The data utilized in the SEDAR75 base model are summarized below and illustrated in **Figure 1** along with their corresponding temporal scale. Comprehensive descriptions of individual data components are provided within each subsection below.

1. Life history
 - a. Meristics
 - b. Age and growth
 - c. Natural mortality
 - d. Maturity
 - e. Discard mortality
2. Landings
 - a. Commercial Vertical Line: 1945-2020 (metric tons whole weight)
 - b. Commercial Longline: 1945-2020 (metric tons whole weight)
 - c. Commercial Nets & Traps: 1945-2020 (metric tons whole weight)
 - d. Recreational Private: 1945-2020 (thousands of fish)
 - e. Recreational Shore: 1945-2020 (thousands of fish)
 - f. Recreational Charter & Headboat: 1945-2020 (thousands of fish)
3. Discards (thousands of fish)
 - a. Commercial Vertical Line: 1993-2020 (metric tons whole weight)
 - b. Commercial Longline: 1993-2020 (metric tons whole weight)
 - c. Recreational Private: 1981-2020 (thousands of fish)
 - d. Recreational Shore: 1981-2020 (thousands of fish)
 - e. Recreational Charter & Headboat: 1981-2020 (thousands of fish)
4. Age composition of landings (1-year age bins, plus group ages 21 and older)
 - a. Commercial Vertical Line: 1991-2020
 - b. Commercial Longline: 1982-2020
 - c. Recreational Private: 1992-2020
 - d. Recreational Charter & Headboat: 1981-2020
5. Length composition of landings (10:80, 2cm Fork Length bins)
 - a. Commercial Vertical Line: 1985-1992
 - b. Commercial Longline: 1990-2000
 - c. Commercial Nets & Traps: 1986-2003
 - d. Recreational Private: 1981-2001
 - e. Recreational Shore: 1981-2020
 - f. Recreational Charter & Headboat: 1983-1990
6. Abundance indices
 - a. Fishery-independent:
 - i. FWRI 0: 1998-2020
 - ii. FWRI 1: 1996-2020
 - iii. SEAMAP Trawl: 2010-2019

- iv. Combined Video Survey: 1993-2020
- v. Reef Fish Visual Survey: 1997-2018
- b. Fishery-dependent:
 - i. Private CPUE: 1981-2020
 - ii. Shore CPUE: 1981-2020
- 7. Length composition of surveys (10:80, 2cm Fork Length bins)
 - a. SEAMAP Trawl: 2010-2019
 - b. Combined Video Survey: 2006-2020
 - c. RF Visual Survey: 1997-2018

2.1. Stock Structure and Management Unit

Two regions (Atlantic and GOM) are currently used by the South Atlantic Fishery Management Council (SAFMC) and GMFMC for Gray Snapper management. The geographic boundary of these management units typically extends from approximately the Dry Tortugas through the Florida Keys (U.S. Highway 1) to mainland Florida. However, the management unit for Gulf of Mexico Gray Snapper extends from the United States–Mexico border in the west through the northern Gulf waters, the Florida Keys and all of Monroe County. Currently, the Council manages Gulf of Mexico Gray Snapper as one unit. There was no Data Workshop (DW) for the SEDAR75, therefore the stock definition was left unchanged from SEDAR51.

2.2. Life History Parameters

Life history data used in the assessment included length-length and length-weight relationships, age and growth, natural mortality, and maturity. Some of the life history data were input to the population model (Stock Synthesis) as fixed values, while other life history parameters were estimated.

2.2.1. Morphometric and Conversion Factors

The length-weight relationship ($W = aFL^b$) for sexes combined was developed at the SEDAR51 Data Workshop, and used as a fixed model input (**Table 1, Figure 2**).

2.2.2. Age and Growth

Additional pairs of length and age were made available during the SEDAR75 Life History Topical Working Group. To account for minimum size regulations beginning in 1990, two scenarios were proposed for size-adjusted growth models: Scenario 1 used all fishery dependent samples and assigned a 12" TL size limit after 1990. Scenario 2 split out the recreational samples caught in Florida state jurisdictional waters after 1990 and assigned those a 10" TL size limit. All other fishery dependent samples were assigned a 12" TL size limit (see SEDAR75-WP-05). The Life History Topical Working Group determined Scenario 2 was the best option for the updated growth model for SEDAR75.

Growth was estimated externally to Stock Synthesis using a single von Bertalanffy growth curve for both sexes combined (**Table 2, Figure 2**). During the modeling process, growth was able to be estimated internally by Stock Synthesis, however this model had poor retrospective patterns.

2.2.3. Natural Mortality

The age-specific vector of natural mortality (M) was not updated during the SEDAR75 (**Table 3, Figure 2**). During the modeling process, natural mortality was attempted to be estimated internally. As the internally estimated growth was not used, it was not necessary to internally estimate natural mortality.

2.2.4. Maturity

The maturity curve for Gray Snapper was updated during the Life History Topical Working Group. A logistic relationship with a logit link function based on fish exhibiting functional maturity as opposed to physiological maturity (see SEDAR75-WP-05). The age at 50% maturity predicted was around 2.5 years and the length at 50% maturity was estimated at 270 mm. (**Figure 3**). This was a slight increase from SEDAR 51, which had an age and length at 50% maturity of 2.3 years and 253 mm respectively. During SEDAR 51, there was a discussion around whether individuals below 300 mm had a significant contribution to the spawning biomass, however there were very few samples from individuals below 300 mm. The updated maturity model includes 126 females sampled below 300 mm as opposed to the 59 females sampled for SEDAR 51. Two models were proposed, one using physiological maturity and the other based on a functional maturity. The functional maturity model chosen had a larger L50 and A50 than the physiological maturity model, but both models had a L50 below the 300 mm discussed during SEDAR 51.

2.2.5. Fecundity

Weight was used as a proxy for fecundity in SEDAR75 as was the case for SEDAR51. Batch fecundity sample sizes for SEDAR75 increased to 12 from the 6 available during SEDAR51, which were insufficient to provide estimates.

2.3. Fishery-Dependent Data

2.3.1. Commercial Landings

The fleet structure used in the previous assessment of Gray Snapper (SEDAR51 Base Model) split the commercial landings into three fleets, handline from Monroe County, handline not from Monroe County and longline. When the landings were re-apportioned from the Data Workshop recommended fleets into the assessment model fleets, some landings were accidentally excluded (**Figure 4**). During the model building phase for SEDAR75, a continuity model was run with the corrected commercial landings (See **Section 4.8.5**.)

Commercial landings data (1963-2019) used in this assessment are presented in **Table 4** and **Figure 5**. The commercial landings are partitioned into three fleets: Commercial Vertical Line gear plus diving, Commercial Longline gear (plus all other gears not covered by Commercial Vertical Line or Commercial Nets & Traps) and Commercial Nets & Traps (all nets and traps gear code). They represent the main commercial harvesting gears capturing Gulf of Mexico Gray

Snapper. Commercial landings were reported in pounds whole weight and converted to metric tons for input to the assessment model.

The commercial fleet structure for SEDAR75 differs from the final fleet structure used in the SEDAR51 assessment. The fleet structure was changed from the recommended fleets from the SEDAR51 Data Workshop due to concerns about weighting length compositions from the Florida Keys. However, new methods for stratifying and weighting length composition data does not necessitate using the same fleet structure as SEDAR51 assessment (see SEDAR75-WP-07).

The majority of commercial landings over time have been from the Commercial Vertical Line fleet (**Figure 5**). Uncertainty estimates were not provided for commercial landings from the Gulf of Mexico. A CV of 0.05 was assigned to these landings.

As in SEDAR51, minor removals of Gray Snapper were assumed to have occurred in the Gulf of Mexico prior to 1945; however, for this evaluation the stock was assumed to be at near unfished condition at the start of the model.

2.3.2. Recreational Landings

Recreational landings data (1945-2020) used in the assessment are presented in **Table 5** and **Figure 5**. For the data period (1981-2019), final recreational landings were computed using fully calibrated estimates from the MRIP using FES, the Southeast Region Headboat Survey (SRHS), Louisiana Creel, and the Texas Parks and Wildlife Department (TPWD) data (see SEDAR75-WP-02). Recreational landings are reported by mode and include Charter, Headboat, Private, and Shore modes. For the assessment, recreational landings from the charterboat and headboat modes were aggregated, as was done in SEDAR51. Private landings represented the dominant mode in the total recreational landings by numbers since 1981. Recreational landings were reported in numbers of fish and input into the assessment model as 1000s of fish.

The fully calibrated estimates differed from the time series of recreational landings used in SEDAR51, particularly for the private mode where annual differences ranged from 4% to 224% (average 87%). Differences in the Shore mode were less, ranging from -23% to 547% (average 84%). Differences in charterboat and headboat for the MRIP period (1981-1985) ranged from -45% to 389% (average 12%).

The fully calibrated time series originally submitted for the Private mode exhibited a very strong peak in 1984 (9,689,657 fish). This peak was discussed during the SEDAR75 Shore Mode Topical Working Group. Given that: 1. it was beyond the range of the rest of the data series, 2. it was driven by a single stratum of western Florida, wave 6 and ocean less than 10 miles. This stratum consisted of 14 angler trips, with three of those trips seen by the observer (SEDAR75-WP-02), and 3. it had a major influence on the historical time series (which uses the average CPUE from 1981-1985 as a scalar, see SEDAR72-WP-05), the decision was made to replace the 1984 peak landings for private mode with the geometric mean of 1981, 1982, 1983, and 1985 private mode landings for that stratum. This was not possible as none of those years had any landings for the stratum. The complete lack of landings in the stratum for the surrounding years was further justification to smooth data. As an alternative, the landings from the 1986 stratum were used for the 1984 stratum. This resulted in a 68% decrease in the point estimate of landings for that year and mode (down to 3,929,241 fish).

Historical estimates (1945-1980) for recreational landings were estimated using the National Survey of Fishing, Hunting, & Wildlife-Associated Recreation (FHWAR) method (For a recent document detailing the methodology, see SEDAR72-WP-05). The FHWAR method utilizes a combination of information including U.S. angler population estimates and angling effort estimates from 1945 – 1985 to estimate effort (saltwater days) for the GOM for every five years when the survey is conducted. For the years in between, a linear interpolation of the estimates is applied. Estimates of effort for 1945-1980 are then multiplied by the mean CPUE for Gulf of Mexico Gray Snapper for 1981 to 1985 (MRIP, SRHS and TPWD combined) to estimate annual landings for the historical time period (1945-1980). For SEDAR75, total historical recreational catches were apportioned by mode using the ratios 64% Private:29% Shore:7% Headboat & Charterboat. These ratios were based on the average proportion of landings by fleet over the period 1981-1985.

Uncertainty estimates were provided for the Recreational Private and Recreational Shore landings for 1981-2019 and for the Recreational Charter & Headboat charterboat mode (SEDAR75-WP-02). However, CVs were not used in SEDAR51 and CVs were fixed at 0.1.

Starting the assessment model in 1945, when the stock is already in a fished state, requires the estimation of initial conditions via initial equilibrium catches, which are used to calculate initial fishing mortality rates. Initial equilibrium catches were calculated for the Recreational Shore fleet as the average landings over the first five years of the assessment time series. Initial runs attempted estimating initial fishing mortality rates for the Commercial Nets & Traps and Recreational Private fleets but the estimates bounded at 0 and were highly correlated with one another. As such the decision was made to only calculate initial fishing mortality rates for the Recreational Shore fleet and fix the initial F of the Commercial Nets & Traps and Recreational Private fleets at 0, as was done in SEDAR 51.

2.3.3. Commercial Discards

Commercial discards (1993-2020) used in SEDAR75 are presented in **Table 6** and **Figures 6-7**. The commercial discards for Gulf of Mexico Gray Snapper were estimated using methods revised since SEDAR 51. The improved methodology made use of CPUE from the coastal reef fish observer program and total fishing effort from the commercial reef logbook program to estimate total catch. As there was no Data Workshop for SEDAR75, a working paper was not submitted for the dataset used, however a full description of the improved methodology and CPUE-expansion estimation procedures can be found in SEDAR72-WP-16. The same methodology has been recently applied to other SEDAR assessments including for GOM Red Grouper, Gray Triggerfish, Vermilion Snapper, Scamp and Greater Amberjack.

The discard estimates reported in numbers were input into the assessment as 1,000s of fish with corresponding log-scale standard errors fixed at 0.3 (SE, **Table 6**). This differs from how discards were input in SEDAR 51. Due to the fleet structure change after the Data Workshop, discards needed to be input as proportion removed rather than total removals. Due to this difference, it is not possible to directly compare the inputs between SEDAR51 and SEDAR75. A discard mortality rate of 6.9%, as recommended by the SEDAR 51 DW, was applied to the commercial discards.

2.3.4. Recreational Discards

Recreational discards from the Recreational Private, Recreational Shore and Recreational Charter & Headboat fleets (1981-2020) used in the assessment are presented in **Table 7** and **Figure 8-10**. Final recreational discards were computed using fully calibrated estimates from MRIP using FES (SEDAR75-WP-02) for Recreational Private (1981-2020), Recreational Shore (1981-2020) and Recreational Charter & Headboat (1981-2020).

Recreational discards were reported as numbers of fish and input into the assessment as 1000s of fish with log-scale standard errors fixed at 0.3 (SE, **Table 7**). A discard mortality rate of 14%, as recommended by the SEDAR51 DW, was applied to all recreational fleets.

2.3.5. Commercial Size Composition

Commercial Vertical Line length compositions of landed (retained) (1985-1992) fish are presented in **Figure 11**, Commercial Longline length compositions of landed (retained) (1990-2000) fish are presented in **Figure 12** and Nets and Traps length compositions in **Figure 13**.

The annual length compositions were combined into 2-cm fork length interval bins (10:80). Length compositions of landings were constructed using the same data sources approved in SEDAR51 (the commercial trip intercept program (TIP) and GulfFIN) but were processed using revised practices for calculating final compositions. Length samples were weighted by the commercial landings at the finest spatial and temporal scale available. A description of the revised methods used to develop the length composition data was provided in SEDAR75-WP-07. The input sample size associated with each year/fleet was calculated by multiplying the number of trips sampled with the percentage of landings represented in the length composition for that fleet/year. Year/fleet combinations with less than 10 trips sampled were removed from the assessment model. Years with available age compositions were removed from the length compositions to prevent double counting of the same sample.

Data from the Reef Fish Observer Program (RFOP) were available for length compositions from commercial discards. However, there were insufficient sample sizes to be used in SEDAR75.

2.3.6. Recreational Size Composition

Recreational Private length compositions of landed (1981-2001) fish are presented in **Figure 14**. Recreational Shore length compositions of landed fish (1981-2001) fish are presented in **Figure 15**. Recreational Charter & Headboat length compositions of landed (1983-1990) fish are presented in **Figure 16**.

The annual length compositions were combined into 2-cm fork length interval bins (10:80). Length compositions of landings were constructed using the same data sources approved in SEDAR 51 (MFRSS/MRIP, SRHS, TPWD, the GulfFIN database, and the TIP database) but were processed using revised practices for calculating final compositions. Length samples were weighted by the recreational landings at the finest spatial and temporal scale available. A description of the revised methods used to develop the length composition data was provided in SEDAR75-WP-07. The input sample size associated with each year/fleet was calculated by multiplying the number of trips sampled with the percentage of landings that represented in the length composition for that fleet/year. Year/fleet combinations with less than 10 trips sampled

were removed from the assessment model. Years with available age compositions were removed from the length compositions to prevent double counting of the same sample.

Data from the Florida Fish and Wildlife Commission (FWC) Fish and Wildlife Research Institute (FWRI) At-Sea Observer Program (2006-2020) were used to characterize the length compositions from recreational discards. However, there were insufficient sample sizes to be used in SEDAR75.

2.3.7. Commercial Age Composition

Commercial age compositions of landed fish used in the assessment are presented in **Figure 17** and **Figure 18**. The Commercial Vertical Line age compositions were input as weighted ages Commercial Longline as nominal ages with sample sizes specified as number of trips (SEDAR75-WP-07).

Cohorts are visible in the Commercial Vertical Line age composition data (**Figure 21**) and in the Commercial Longline data (**Figure 22**). The main age classes captured were 4-12-year olds and 5-16-year olds for the Vertical Line and Longline fleets, respectively.

2.3.8. Recreational Age Composition

Recreational age compositions of landed fish used in the assessment are presented in **Figures 19-20**. The recreational age compositions were input as weighted ages with sample sizes specified as number of trips (SEDAR75-WP-07). Nominal age compositions for Recreational Shore were available, however during the modeling process, these were removed as the model had difficulty fitting the ages at the expense of the available length compositions for Recreational Shore.

The apparent cohorts in the Recreational Private data include 2005, 2012 and 2014 (**Figures 23**). The apparent cohorts in the Recreational Charter & Headboat data include the 1991 and 2005 cohorts. The main age classes captured by the Recreational Private fleet were 2-6-year olds. The main age classes captured by the Recreational Charter & Headboat were 3-7-year olds.

2.3.9. Recreational Catch Per Unit of Effort Indices of Abundance

The standardized CPUE indices for the Recreational fleets used in the assessment are summarized in **Table 8**. Two recreational indices were used in the SEDAR75 assessment model: the MRIP Private Mode index (**Figure 25**) and the MRIP Shore mode index (**Figure 26**). The MRIP Private mode and Shore mode indices tracks total catches of Gray Snapper (landed plus discards). As in SEDAR51, the guild approach was used to select trips for the Shore mode with higher probability of encountering Gray Snapper and the Stephens and MacCall (2004) approach was used for subsetting the trips for the Private mode. Annual CVs were scaled to a common mean of 0.2 for the two fishery-dependent indices and they were converted to log-scale SEs for input into SS (**Table 9**).

2.4. Fishery-Independent Surveys

2.4.1. Age-0 and Age-1 Surveys

Age-0 and age-1 Gray Snapper indices were developed for the Gulf of Mexico using the State of Florida FWC Estuarine fishery-independent monitoring (FIM) Survey (**Table 8, Figure 27**,

Figure 28). See SEDAR75-WP-01 for a full description of the methods used to develop this index.

Overall, the index remained relatively unchanged with the inclusion of additional years of data compared with SEDAR51. Annual CVs were converted to log-scale SEs for input in Stock Synthesis (**Table 9**) and an additional SE was estimated as part of the data weighting process (see **Section 3.2**).

2.4.2. SEAMAP Trawl Survey

The primary objective of SEAMAP trawl survey is to collect data on the abundance and distribution of demersal organisms in the northern Gulf of Mexico. Following the recommendation of the SEDAR 51 Indices Workgroup, a single abundance index was produced for Gray Snapper. The index utilized data solely from summer surveys in the eastern GOM due to the scarcity of Gray Snapper in the western GOM and the gaps in spatial coverage during the fall surveys (Pollack et al. 2017). See SEDAR75-WP-03 for a full description of the methods used to develop this index.

This index was updated through 2019 (**Table 8, Figure 29**). Annual CVs were converted to log-scale SEs for input into SS (**Table 9**).

Length composition for the survey comprised a total of 1,233 individuals measured 2010-2019. Length compositions were input as nominal lengths with sample sizes specified as the number of individuals measured (**Figure 30**).

2.4.3. Combined Video Survey

The combined video survey used SEDAR 51 was updated through 2020. This survey uses three different stationary video surveys for reef fish in the northern Gulf of Mexico (GOM). The NMFS SEAMAP reef fish video survey (SFRV), carried out by NMFS Mississippi Laboratory, has the longest running time series (1993-1997, 2002, and 2004+), followed by the NMFS Panama City lab survey (PC; 2005+), with the most recent survey being the Florida Fish and Wildlife Research Institute video survey (FWRI, starting year 2010). For more information on the survey methodology from the recent red snapper Data Workshop (SEDAR 74), see SEDAR74-WP-23. (**Table 8, Figure 31**). Annual CVs were converted to log-scale SEs for input into SS (**Table 9**).

Length composition for the survey comprised a total of 3,228 individuals measured in 1996-1997, 2002 and 2004-2020. Length compositions were input as nominal lengths with sample sizes specified as the number of survey stations from which successful measurements were obtained (**Figure 32**). Sample sizes below 10 trips annually were omitted.

2.4.4. Reef Fish Visual Survey

The Reef Fish Visual Census was updated through 2018 for SEDAR75. This survey was derived as the mean number observed per 20-minute observation period (See SEDAR75-WP-06). (**Table 8, Figure 33**). Annual CVs were converted to log-scale SEs for input into SS (**Table 9**).

Length composition for the survey comprised a total of 3,744 individuals measured in 1997-2012, 2014, 2016 and 2018. Length compositions were input as nominal lengths with sample sizes specified as the number individuals measured (**Figure 34**).

2.5. Environmental Considerations & Contributions from Stakeholders

No environmental or red tide data were submitted for SEDAR75.

3. Stock Assessment Model Configuration and Methods

3.1. Stock Synthesis Model Configuration

The assessment model used was Stock Synthesis (SS), version 3.30.19. Descriptions of SS algorithms and options are available in the SS User's Manual (Methot et al. 2020), the NOAA Fisheries Toolbox website (<http://nft.nefsc.noaa.gov/>), and Methot and Wetzel (2013). Stock Synthesis (SS) is a widely used integrated statistical catch-at-age model (SCAA) that has been tested for stock assessments in the United States (US), particularly on the West Coast and Southeast, and also throughout the world (see Dichmont et al. 2016 for review). SCAA models consist of three closely linked modules: the population dynamics module, an observation module, and a likelihood function. Input biological parameters (e.g., **Section 2.2**) are used to propagate abundance and biomass forward from initial conditions (population dynamics model) and SS develops predicted data sets based on estimates of fishing mortality, selectivity, and catchability (the observation model). The observed and predicted data are compared (the likelihood module) to determine best-fit parameter estimates using a statistical maximum likelihood framework (detailed in Methot and Wetzel (2013)). Because many inputs are correlated, the concept behind SS is that processes should be modeled together, which helps to ensure that uncertainties in the input data are properly accounted for in the assessment.

The Gulf of Mexico Gray Snapper SS model assumed a similar configuration structure as developed for the previous SEDAR 51 Gulf of Mexico Gray Snapper Benchmark, with the exception of the commercial fleet structure (**Section 2.3.1**). The fully configured SS model included observations of catch for six fishery fleets (Commercial Vertical Line, Commercial Longline, Commercial Nets & Traps, Recreational Private, Recreational Shore and Recreational Charter & Headboat) and discards for five fishery fleets (Commercial Vertical Line, Commercial Longline, Recreational Private, Recreational Shore and Recreational Charter & Headboat). The model included two fishery-dependent CPUE indices of abundance (Private CPUE and Shore CPUE), and three fishery independent time series (FWRI Age 0, FWRI Age 1, SEAMAP Trawl, Combined Video Survey, and RF Visual Survey). Model estimated parameters include growth parameters, fishing mortality by fleet for each year, selectivity and retention parameters for each directed fleet, parameters describing the stock-recruit function, stock-recruit deviation parameters, index catchabilities, and Dirichlet multinomial parameters.

The SS modeling framework provides estimates for key derived quantities including: time series of recruitment (units: 1,000s of age-0 recruits), abundance (units: 1,000s of fish), biomass (units: metric tons), SSB (units: metric tons), and harvest rate (units for Gray Snapper: total biomass killed age 2+ / total biomass age 2+). The r4ss software (Taylor et al. 2021) was utilized

extensively to develop various graphics for model outputs and was also used to summarize various output files and perform diagnostic runs.

Projections are implemented within SS starting from the year succeeding the terminal year of the assessment model utilizing the same population dynamics equations and modeling assumptions.

3.1.1. Initial Conditions

The Gulf of Mexico Gray Snapper assessment begins in 1945 and has a terminal year of 2020. Since removals of Gray Snapper were assumed to be negligible in the Gulf of Mexico prior to 1945 for both commercial and recreational fisheries, the stock was assumed to be at equilibrium.

3.1.2. Temporal Structure

The Gray Snapper population was modeled from age-0 through age-28 (the maximum age), with data bins spanning age-0 through age-21+, with the last age representing a plus group. Data collection and fishing activities were assumed relatively continuous throughout the year; therefore, inclusion of a seasonal component to the removals was not deemed necessary. The fishing season was assumed to be continuous and homogeneously distributed throughout the year.

3.1.3. Spatial Structure

A single area model was implemented where recruits are assumed to homogeneously settle across the entire Gulf of Mexico region.

3.1.4. Life History

A fixed length-weight relationship was used to convert body length (cm Fork Length, FL) to body weight (kg whole weight; **Table 1, Figure 2**). Stock Synthesis moves fish among age classes and length bins on January 1st of each modeled year starting from birth at age-0. Because the ‘true’ birth date often does not occur on January 1st, with peak spawning occurring around July 15 for Gray Snapper in the Gulf of Mexico, some slight alterations in growth (t_0 , or the age at length 0) and natural mortality parameters are required to account for the difference between true age and modeled age when parameters are input instead of estimated.

Growth was modeled with a three parameter von Bertalanffy equation: (1) L_{Amin} (cm FL), the mean size at age-1 Gray Snapper; (2) L_{Amax} (cm FL), the mean size at maximum aged Gray Snapper; and (3) K (year^{-1}), the growth coefficient. In Stock Synthesis, when fish recruit at the real age of 0.0 they have a body size equal to the lower limit of the first population bin (fixed at 10 cm FL). Fish then grow linearly until they reach a real age equal to the input value of A_{min} (growth age for L_{Amin}) and have a size equal to L_{Amin} . As they age further, they grow according to the von Bertalanffy growth equation (**Figure 2**). L_{Amax} was specified as equivalent to L_{inf} . Two additional parameters are used to describe the variability in size-at-age and represent the CV in length-at-age at A_{min} (age 1) and A_{max} (age 28). For intermediate ages, a linear interpolation of the CV on mean size-at-age is used.

The von Bertalanffy growth model parameters L_{Amax} and K were estimated externally to SS using updated length and age compositions while L_{Amin} was estimated by SS. Variance parameters

CV_{Amin} (0.1514) and CV_{Amax} (0.1922) were fixed at the values recommended at the SEDAR51 DW (Table 2).

The age-specific vector of M (Section 2.2.3) was fixed within the SS model (Table 3, Figure 2).

3.1.5. Recruitment Dynamics

A Beverton-Holt stock-recruit function was used to parameterize the relationship between spawning output and resulting recruitment of age-0 fish. The stock-recruit function (representing the arithmetic mean spawner-recruit levels) requires three parameters: (1) steepness (h) characterizes the initial slope of the ascending limb (i.e., the fraction of virgin recruits produced at 20% of the equilibrium spawning biomass); (2) the virgin recruitment (R_0 , estimated in log space) represents the asymptote or virgin recruitment levels; and (3) the variance or recruitment variability term (σ_R) which is the SD of the log of recruitment (it both penalizes deviations from the spawner-recruit curve and defines the offset between the arithmetic mean spawner-recruit curve and the expected geometric mean from which the deviations are calculated). Similar to SEDAR51, h was fixed at 0.99, in the SEDAR75 Base Model, while σ_R and virgin recruitment ($\ln R_0$) were freely estimated.

Annual deviations from the stock-recruit function were estimated in SS as a vector of deviations forced to sum to zero and assuming a lognormal error structure. A lognormal bias adjustment factor was applied to recruitment estimates as recommended by Methot et al. (2020), but only to the data-rich years in the assessment. This was done so that SS will apply the full bias-correction only to those recruitment deviations that have enough data to inform the model about the full range of recruitment variability (Methot et al. 2020). For the SEDAR75 Base Model, main period (i.e. data rich) recruitment deviations spanned 1981-2020, while early period (i.e. data poor) recruitment deviations spanned 1950 -1980. Full bias adjustment was used from 1991 to 2017 when length or age composition data are available. Bias adjustment was phased in linearly, from no bias adjustment prior to 1960 to full bias adjustment in 1991. Bias adjustment was phased out in 2017, decreasing from full bias adjustment to no bias adjustment in that year, because the age composition data contains less information on recruitment in more recent years. The years selected for full bias adjustment were estimated following the methods of Methot and Taylor (2011).

3.1.6. Fleet Structure and Surveys

Four fishing fleets were modeled and had both associated length and age compositions. The SS fleet codes for these were: Commercial Vertical Line (Com_VL_1), Commercial Longline (Com_LL_2), Recreational Private (Rec_PR_4), Recreational Shore (Rec_Shore_5) and Recreational Charter & Headboat (Rec_HB_CBT_6). Two fishing fleets were modeled with associated length compositions. The SS fleet codes for these were: Commercial Nets & Traps (Com_NT_3) and Recreational Shore (Rec_Shore_5). This structure differs from SEDAR51 for the commercial fleets and the inclusion of age compositions for four fleets. Fishing was assumed to be continuous and homogeneous across the entire year. Discards were modeled as total removals whereas SEDAR51 used the annual proportion discarded (discards/ (total landings + discards)).

Two fishery-dependent CPUE indices were included in the SEDAR75 Base Model: Private CPUE and Shore CPUE (CPUE units: number kept or discarded per angler hour). CPUE was

treated as an index of biomass or abundance where the observed standardized CPUE time series was assumed to reflect annual variation in population trajectories. Both the Private CPUE and Shore CPUE were input as surveys into SS (see **Section 2.3.10**) and the selectivities were mirrored to age selectivity of the Recreational Private and the length selectivity in Recreational Shore fleets.

Five fishery-independent surveys were included in the SEDAR75 Base Model: the FWRI 0, the FWRI 1, SEAMAP Trawl, Combined Video Survey and the RF Visual Survey. The FWRI 0 was set up as a special survey of Age-0 recruits (i.e. age based selectivity restricted to, and fully selecting, age 0) while FWRI 0 fully selected for all ages. The SEAMAP Trawl, Combined Video Survey and RF Visual Survey had length observations available which were fit directly based on estimated length-based selectivity functions.

3.1.7. Selectivity

Selectivity represents the probability of capture by age or length for a given fleet and represents the net result of multiple interrelated factors (e.g., gear type, targeting, and availability of fish due to spatial and temporal constraints). SS allows users to specify length-based selectivity, age-based selectivity, or both. The final selectivity curve governing each fleet/survey reflects the additive effect of both age- and length- based processes.

Selectivity patterns were assumed to be constant over time for each fleet and survey. The Gulf of Mexico Gray Snapper fishery has experienced changes in management regulations in 1990 when a minimum size limit was implemented, which was assumed to influence the discard patterns more so than selectivity. As such, these changes were accounted for in the assessment model using time-varying retention patterns (see **Section 3.1.8.**) and modeling discards explicitly (see **Section 3.1.10.**).

3.1.7.1. Length-based Selectivity

Length-based selectivity patterns were specified for each fleet and survey and were characterized as one of two functional forms: (1) a two-parameter logistic function (SS pattern 1) and (2) a six-parameter double normal function (SS pattern 24). A logistic curve implies that fish below a certain size range are not vulnerable, but then gradually increase in vulnerability with increasing size until all fish are fully vulnerable (asymptotic selectivity curve). Two parameters describe logistic selectivity: (1) the length at 50% selectivity, and (2) the difference between the length at 95% selectivity and the length at 50% selectivity, which were both estimated in this assessment. The double normal has the feature that it allows for domed or logistic selectivity and is a combination of two normal distributions; the first describes the ascending limb, while the second describes the descending limb. A line segment joins the maximum selectivity of the two functions. However, the double normal functional form can be more unstable than other selectivity functions due to the increased number of parameters. When robust length or age compositions are available with sufficient numbers of larger or older fish, it may be appropriate to freely estimate all parameters (especially the descending limb). If that is not the case, certain parameters can be fixed to improve model stability as long as fixing the parameter does not largely influence the point estimates of the remaining selectivity parameters. Unless strong evidence exists for domed selectivity, it is generally advisable to use the logistic function.

In the SEDAR75 Base Model, separate selectivity patterns were defined for each fleet/survey: 1) Commercial Vertical Line (double normal), 2) Commercial Longline (double normal), 3) Commercial Nets & Traps (double normal), 4) Recreational Private (double normal), 5) Recreational Shore (double normal), 6) Recreational Shore (double normal), 7) FWRI Age-0 (only age 0 fully selected), 8) FWRI Age-1 (full selectivity after age 0 +), 9) SEAMAP Trawl (double normal), 10) Combined Video Survey (logistic) and 11) RF Visual Survey (double normal). The fishery selectivity patterns remained the same as those used in SEDAR51; the survey selectivity patterns changed from SEDAR51 as length composition data was available and the use of fleet mirroring was not necessary.

A logistic selectivity pattern was attempted for the commercial longline fleet (with all parameters freely estimated) because there was little evidence in the age data suggesting availability issues that might make older fish less vulnerable. This was evident in catch curves developed for the longline fleet, where the lognormally distributed catch-at-age was regressed against age using the equation from Quinn and Deriso (1999):

$$\ln(C_a) = [\ln(\mu N_f) + fZ] - Z_a$$

where μ is the probability of catching a fish, N_f is the abundance at the start of age a , and Z is the total mortality at age- a . The estimate of Z is the negative of the slope estimated from the linear regression, and its SE is equal to the SE of the slope. The corresponding estimate of survival-at-age (S_a) is $\exp(Z)$. A catch curve typically shows an increasing section of the curve for younger ages, due to increasing availability of fish or selectivity of the gear, followed by a decreasing trend for older ages due to increased mortality stemming from full selectivity by the fishing or survey gear. Steep slopes (e.g., > 1) are generally evidence for dome-shaped selectivity, as was the case for the vertical line fleet (**Figure 35**). The catch curve for the longline commercial fleet showed increases in selection of younger fish, full selection by 11 years, and a gradual decline with age characterized by a relatively shallow slope (**Figure 35**). For the Commercial Longline fleet, the estimation ignored the last size bins and allowed SS to decay the large fish selectivity according to parameters the descending width to reduce the number of parameters being estimated and improve model stability. For Commercial Nets & Traps, the beginning size of the plateau, the first parameter, was fixed to improve model stability.

Double normal selectivity was implemented for all three recreational fleets because dome-shaped selectivity was considered highly likely due to areas fished (e.g., closer to shore, shallower) and targeting behavior. For the recreational fleets, the estimation ignored the first and last size bins and allowed SS to decay the small and large fish selectivity according to parameters of ascending width and descending width, respectively, to reduce the number of parameters being estimated and improve model stability. For Recreational Private and Recreational Shore, the beginning size of the plateau, the first parameter, was fixed to improve model stability.

Logistic selectivity was assumed for the Combined Video Survey since the survey targeted high relief areas that the largest individuals are known to occupy. Both parameters were freely estimated. For SEAMAP Trawl and RF Visual Survey, dome-shaped selectivities provided the best fits to the length compositions; the estimation ignored the first and last size bins and allowed SS to decay the small and large fish selectivity according to parameters of ascending width and descending width, respectively, to reduce the number of parameters being estimated and improve model stability.

The selectivity of the FWRI Age-0 Survey did not need to be specified as the survey was set up as a recruitment index (i.e. pre-specified to select age-0 fish only).

3.1.7.2. Age-based Selectivity

Age-based selectivity was specified for Commercial Vertical Line, Commercial Longline, Recreational Private, and Recreational Charter & Headboat. Initially, full age selectivity was restricted to ages 1+, however this provided very poor fits to the age composition and had poor model stability. All four fleets used a fully estimated dome-shaped selectivity, with one exception. The Vertical Line fleet had the age of inflection fixed due to bounding issues.

3.1.7.3. Mirroring

The age and length-based selectivity patterns of the Private CPUE and Shore CPUE indices were assumed to mirror the selectivity pattern of their respective fleets. The age-based selectivity patterns of the Private CPUE index was made to mirror the selectivity pattern of the Recreational Private fleet and the length-based selectivity patterns of the Shore CPUE index was made to mirror the selectivity pattern of the Recreational Shore.

3.1.8. Retention

Time-varying retention functions are commonly used in Gulf stock assessments to allow for varying discards at size due to the impacts of management regulations. For Gray Snapper, time blocks were based on changes in the Federal and Florida state waters minimum size limits. The time varying retention blocks were defined as:

For all fishing fleets:

1. 1945 - 1989: no minimum size limit regulation in place
2. 1990-2020: 12 inches FL Federal waters minimum size limit
3. 1990-2020: 10 inches FL Florida waters minimum size limit

For each fleet, the retention function was specified as a logistic function consisting of four parameters: (1) the inflection point, (2) the slope, (3) the asymptote, and (4) the male offset inflection (not applicable to this model and assumed to be zero). The blocks related to the minimum size limits were linked to the inflection point parameters.

For the period of 1945-1989, the inflection of the retention curve and the slope of the retention function was allowed to be freely estimated. From 1990 on, the inflection of the retention function was estimated for Commercial Vertical Line and Recreational Charter & Headboat with the asymptotes fixed at 1 (knife-edged). The inflection point was allowed to be estimated for these two fleets to improve model stability but had estimated values close to or at the initial parameters of the size limits. From 1990 on, the inflection of the retention function was fixed at the federal size limits for Commercial Longline with the asymptote fixed at 1. From 1990 on, the inflection of the retention function was fixed at the Florida state size limits for Recreational Private and Recreational Shore and with the asymptote fixed at 1. This was done to account for the majority of the recreational landings from the private and shore modes coming from Florida state waters.

3.1.9. Landings and Age Compositions

Landings by fleet and associated length and age compositions were estimated using fleet-specific continuous fishing mortality rates and length-specific selectivity curves following Baranov's catch equation.

The commercial landings were assumed to be the most representative and reliable data source in the model, especially over the most recent time period. This information was collected in the form of a census as opposed to being collected as part of a survey and a CV of 0.05 was assumed, as was the case in SEDAR51. The recreational landings were assumed to be less precise than the commercial landings and a CV of 0.1 was assumed for all three recreational fleets, as was the case in SEDAR51. All CVs were converted to a log-scale SE (see **Section 3.2.**).

A new feature available for fitting composition data in SS is the Dirichlet Multinomial (DM) which differs from the standard multinomial in that it included an estimable parameter (θ) which scales the input sample size (Thorson et al. 2017; Methot et al. 2020). The DM is self-weighting, which avoids the potential for subjectivity as when the Francis re-weighting procedure is applied (Francis 2011). The DM approach also allows for observed zeros in the data, and the effective sample sizes calculated are directly interpretable. The DM uses the input sample sizes directly, adjusted by an estimated variance inflation factor. The more positive the inflation factor, the more weight the data carry in the likelihood. The DM is considered an improved practice and recommended for use by the SS model developers, and was first used in a Gulf stock assessment during SEDAR70 in 2020 for Gulf of Mexico Greater Amberjack.

Because SS models the growth internally and tracks individual fish from birth, it actually grows fish by length bins before eventually converting lengths to ages (based on the growth curve). As such, it is possible to fit both age and length composition simultaneously. For SEDAR75, the age and length composition data for each fleet/survey were assumed to follow a Dirichlet multinomial error structure where sample size represented the number of trips (of adjusted number of trips, see **Sections 2.3.5.** and **2.3.6.**), adjusted by an estimated variance inflation factor. Input sample sizes were related to the number of trips/sets rather than the number of measurements taken because using the number of lengths can overestimate sample sizes in fisheries data, as samples are rarely truly random or independent (Hulson et al. 2012). In addition, using higher effective sample sizes can lead to the composition data dominating the likelihood and reduce fit to other data sources. See **Sections 2.3.5-2.3.8** and **Sections 2.4.2-2.4.3** for more detail on input sample sizes for each fleet/survey. The final effective sample sizes for each year are provided on the figures illustrating the age composition and length composition (given by N_{adj} in each panel; **Figures 17-20**).

3.1.10. Discards

Discard data for each fleet were directly fit in the SS model using size-based retention functions, and a log-normal error structure was assumed. The model estimates total discards based on the selectivity and retention functions, then calculates dead discards based on the discard mortality rates of 14% and 6.9% for the commercial and recreational fleets, respectively (**Sections 2.3.3-2.3.4**).

3.1.11. Indices

The indices are assumed to have a lognormal error structure. The CVs provided by the index standardization were converted to a log-scale SE required for input to SS for lognormal error structures (Section 3.2.).

3.2. Goodness of Fit and Assumed Error Structure

A maximum likelihood approach was used to assess goodness of model fit to each of the data sources (e.g., catch, indices, compositions, etc.). For each separate data set, an assumed error distribution and an associated likelihood component was specified, the value of which was determined by the difference in observed and predicted values along with the assumed variance of the error distribution. The total likelihood was the sum of each individual component. A nonlinear iterative search algorithm was used to minimize the total negative log-likelihood across the multidimensional parameter space to determine the parameter values that provide the best fit to the data. With this type of integrated modeling approach, data weighting (i.e., the variance associated with each data set) can impact model results, particularly if the various data sets indicate differing population trends.

Where lognormal error structures were used, annual CVs associated with each of the data sources were converted to log-scale SEs using the approximation: $\log_e(SE) = \sqrt{(\log_e(1 + CV^2))}$ provided in Methot et al. (2020).

Weak penalty functions were implemented to keep parameter estimates from hitting their bounds, which includes a symmetric-beta penalty on age selectivity parameters (Methot et al. 2020). Parameter bounds were set to be relatively wide and were unlikely to truncate the search algorithm.

Uncertainty in parameter estimates was quantified by computing asymptotic SEs for each parameter. Asymptotic SEs are calculated by inverting the Hessian matrix (i.e., the matrix of second derivatives) after the model fitting process (Methot and Wetzel, 2013). Asymptotic SEs provide a minimum estimate of uncertainty in parameter values.

3.3. Estimated Parameters

In all, 652 parameters were included in the analysis for the SEDAR75 Base Model, of which 576 were active parameters (Table 10). These parameters include: year specific (1945-2020) fishing mortality for each fleet, the stock-recruit deviations for the data-poor time period (1951-1980) the stock-recruit deviations for the data-rich time period (1981-2020), one von Bertalanffy growth parameter (L_{Amin}), two stock-recruit relationship parameters ($\ln(R_0)$ and σR), size selectivity parameters for each fleet or survey, logistic retention parameters for each fleet, catchability parameters for each index, 13 parameters informing the Dirichlet multinomial length and age composition weightings.

3.4. Model Diagnostics

3.4.1. Residual Analysis

The main approach used to address model fit and performance was residual analysis of model fit to each of the data sets (e.g., catch, indices, length/age compositions, discards). Any temporal trends in model residuals (or trends with age or length for compositions data) can be indicative of model mis-specification and poor performance. It is not expected that any model will perfectly fit any of the observed data sets, but ideally, residuals will be randomly distributed and conform to the assumed error structure for that data source. Any extreme patterns of positive or negative residuals are indicative of poor model performance and potential unaccounted for process or observation error.

3.4.2. Correlation Analysis

High correlation among parameters can lead to flat likelihood response surfaces and poor model stability. By performing a correlation analysis, modeling assumptions that lead to inadequate model parameterizations can be highlighted. Because of the highly parameterized nature of stock assessment models, it is expected that some parameters will always be correlated (e.g., stock recruit parameters). However, a large number of extremely correlated parameters warrant reconsideration of modeling assumptions and parametrization. A correlation analysis was carried out and correlations with an absolute value greater than 0.7 were reported.

3.4.3. Profile Likelihoods

Profile likelihoods are used to examine the change in log-likelihood for each data source in order to address the stability of a given parameter estimate, and to see how each individual data source influences the estimate. The analysis is performed by holding the given parameter at a constant value and rerunning the model. This is repeated for a range of reasonable parameter values. Ideally, the graph of negative log likelihood values against parameter values will give a well-defined minimum, indicating that data sources agree. When a given parameter is not well estimated, the profile plot may show conflicting signals across the data sources. The resulting total likelihood surface will often be flat, indicating that multiple parameter values are equally likely given the data. In such instances, the model assumptions need to be reconsidered.

For this assessment, a profile on the log of virgin recruitment $\ln(R_0)$ was conducted.

3.4.4. Jitter Analysis

Jitter analysis is a relatively simple method that can be used to assess model stability and to determine whether a global as opposed to local minima has been found by the search algorithm. The premise is that all of the starting values are randomly altered (or ‘jittered’) by an input constant value and the model is rerun from the new starting values. If the resulting population trajectories across a number of runs converge to the same final solution, it can be reasonably assumed that a global minimum has been obtained. This process is not fault-proof and no guarantee can ever be made that the ‘true’ solution has been found or that the model does not contain misspecification. However, if the jitter analysis results are consistent, it provides additional support that the model is performing well and has come to a stable solution. For this

assessment, a jitter value of 0.1 (10%) was applied to the starting values and 100 runs were completed.

3.4.5. Retrospective Analysis

A retrospective analysis is a useful approach for addressing the consistency of terminal year model estimates. The analysis sequentially removes a year of data at a time and reruns the model. If the resulting estimates of derived quantities such as SSB or recruitment differ significantly, particularly if there is serial over- or underestimation of any important quantities, it can indicate that the model has some unidentified process error, and requires reassessing model assumptions. It is expected that removing data will lead to slight differences between the new terminal year estimates and the updated estimates for that year in the model with the full data. Oftentimes additional data, especially compositional data, will improve estimates in years prior to the new terminal year, because the information on cohort strength becomes more reliable. Therefore, slight differences are expected between model runs as more years of data are peeled away. Ideally, the difference in estimates will be slight and more or less randomly distributed above and below the estimates from the model with the complete data sets. A five-year retrospective analysis was carried out for the SEDAR75 Base Model.

3.4.6. Additional Diagnostics

Additional diagnostics using the R package ‘SS3Diags’ are presented following the recommendations of Carvahlo et al. (2021). Joint residual plots were used to assess goodness of model fit by identifying conflicting time series and auto-correlation of residual patterns via a Loess smoother (Winker et al. 2018; Carvahlo et al. 2021). Undesirably high root mean squared error (RMSE) were values which exceeded 30%. Model misspecification was evaluated by exploring patterns in residuals of indices and compositions using a runs test, which indicates the presence of nonrandom variation (Carvahlo et al. 2021). In addition, outlier data points were identified via the 3-sigma limit, where any points beyond this limit would be unlikely given random process error in the observed residual distribution (Carvahlo et al. 2021).

Prediction skill of the model was tested using the hindcasting cross-validation approach of Kell et al. (2021). The mean absolute scaled error (MASE; Hyndman and Koehler 2006) was calculated for a 5-year period for each data input where available. The MASE scales the mean absolute error (MAE) of forecasts (i.e., prediction residuals) to the MAE of a naïve in-sample prediction (Carvahlo et al. 2021). A skilled model would improve the model forecast compared to the baseline (i.e., random walk), with a MASE value of 0.5 indicative of a forecast being twice as accurate as the baseline and values >1 indicative of average model forecasts worse than the baseline (Carvahlo et al. 2021; Kell et al. 2021).

3.4.7. Sensitivity Runs

Sensitivity runs were conducted with the SEDAR75 Base Model to investigate critical uncertainty in data and reactivity to modeling assumptions. An exhaustive evaluation of model uncertainty was not carried out, but the aspects of model uncertainty judged to be the most important for model performance and accuracy were investigated. Only the most important sensitivity runs are presented below, but many additional exploratory runs were also implemented. The order in which they are presented is not intended to reflect their importance;

each run included here provided important information for developing or evaluating the base case model and alternate states of nature. The focus of the sensitivity runs was on population trajectories, improvements in fit and important parameter estimates (e.g., recruitment).

Start year - A start year of 1986 was explored and all data inputs were modified to begin in 1986.

Natural Mortality (M) - Model sensitivity to the specification of the natural mortality rate was evaluated. The target M of the base model was 0.15, a sensitivity run using the SEDAR51 low (target = 0.13) and high (target = 17) M vectors was done (**Table 3**).

4. Stock Assessment Model - Results

4.1. Estimated Parameters and Derived Quantities

A summary of model parameters for the SEDAR75 Base Model can be found in **Table 10**. Results included are estimated parameter values and their associated CVs from SS, initial parameter values, minimum and maximum bounds on parameters, and the prior densities assigned to each parameter (if a prior was used). Most parameter estimates and variances were reasonably well estimated (i.e., $CV < 1$). Of the 576 active parameters, 23 exhibited CVs above 1, including 15 recruitment deviations, retention curves for Commercial Vertical Line and Commercial Longline, the parameter defining the downslope of the age selectivity for Commercial Vertical Line, Commercial Longline and Recreational Private, and the Dirichlet Multinomial parameter on the Commercial Vertical Line length compositions.

4.2. Fishing Mortality

The exploitation rate (total biomass killed age 2+ / total biomass age 2+) for the entire stock are provided in **Table 11** and **Figure 36**. Since 1945, the exploitation rate for the stock has averaged around 0.055, and ranged between 0.002 in 1945 to 0.133 in 1991. The exploitation rate has gradually increased from low levels (less than ~0.1) to near 0.1 in the 1980s and 1990s. It then remained near those values until the mid-2000s after which rates started to slightly increase with larger inter-annual variations. The terminal year (2020) exploitation rate for the entire stock was 0.088, which is slightly above the time series mean.

Table 12 and **Figure 37** provide estimates of exploitation rate by fleet and year. The results show that the exploitation rate for the stock was driven largely by the Recreational Shore fleet throughout the entire time series. The next largest exploitation rates were that of Recreational Shore.

4.3. Selectivity

A comparison of the SS estimated length-based selectivity functions for each directed fleet for Gulf of Mexico Gray Snapper from the SEDAR75 and SEDAR51 models is shown in **Figure 38**. **Figures 39-44** provide fleet specific terminal year (2020) selectivity, retention, discard mortality and fraction of fish kept, dead and discarded for the 6 directed fisheries for both the SEDAR75 and SEDAR51 assessments. **Figure 45** presents SS derived age-based selectivity for each fleet in 2020. Commercial Longline attained maximum selection at age 19. The Commercial Longline fleet reached 50% selectivity at age 15. The Commercial Nets & Traps fleet attained maximum

selection at age 6. Both the Recreational Private and Recreational Shore fleets attained maximum selection at the youngest age of 1. Recreational Charter & Headboat had the oldest age of maximum selection of 20.

The estimated length-based selectivity functions for the FWRI Age-0 Survey through RF Visual Survey for the SEDAR75 vs. SEDAR51 are shown in **Figure 46**. The derived age-based selectivity functions are shown in **Figure 47** only for SEDAR75 because no ages were included for SEDAR51.

All selectivity parameter estimates and associated uncertainty are listed in **Table 10** with the Label prefix “Size_”.

4.4. Retention

Time-varying retention functions, by time block, are provided for each directed fleet and are shown in **Figures 48-52**. All retention parameter estimates and associated uncertainty are listed in **Table 10** with the Label prefix “Retain_”.

Most retention parameters appeared well estimated except for the 1945 to 1989 time block on Commercial Vertical Line and Commercial Longline (**Table 10**).

4.5. Recruitment

As noted in the description of the SS model configuration, one of three of the S/R parameters were fixed at values agreed upon during SEDAR51: steepness (0.99). The corresponding Beverton-Holt stock recruit relationship is shown in **Figure 53**. Estimated annual recruitment of age-0 fish (1000s) from 1951-2021 including recruitment deviations and variance are shown in **Table 13** and **Figures 54-56**. Virgin recruitment in log-space ($\ln(R_0)$) was estimated at 10.052 (**Table 10**), which equates to 23.19 million age-0 Gray Snapper. The estimated (and applied) recruitment bias adjustment ramp is shown in **Figure 57**.

During the main recruitment period (1981-2020, see **Section 3.1.5.**), estimated recruitment averaged 23.08 million Gray Snapper and was lowest in 1989 at 9.88 million Gray Snapper and highest in 2018 at 46.23 million Gray Snapper (**Figure 54**). Recruitment deviations were characterized by a period of lower than average recruitment in the late 1980's and mid 1990's followed by a period of large interannual variations until a period of above average recruitment in the mid-2010s. There was a noticeable drop in recruitment in 2013 (an 56% drop from the previous year), which coincides with a strong signal of recruitment failure in the age-0 survey index (**Figure 27**) and the age composition of the Commercial Vertical Line (**Figures 21**).

CVs for recruitment deviations during the main recruitment period averaged 0.101 between 2018 and 2017, and ranged from 0.06 in 2006 to 0.208 in 1982 (**Figure 56**). For the last two years of the assessment (2019, 2020), recruitment deviations were largely informed by the age-0 index, as age-0 and 1 fish had not yet fully recruited to the fisheries. Estimated recruitment for those terminal years were above average, their estimated values and associated CVs were 44.247 million Gray Snapper (CV=0.127) and 31.341 million Gray Snapper (CV=0.153), respectively.

4.6. Biomass and Abundance Trajectories

The estimated annual total biomass (metric tons), exploitable biomass (ages 2+, metric tons), SSB (metric tons), SSB ratio (SSB/virgin SSB) and exploitable abundance (1,000s of fish) from 1945 to 2020 are provided in **Table 13**. Total biomass averaged 32,782 metric tons, and ranged from 16,115 metric tons in 1997 to 56,914 metric tons in 1945 (**Figure 58**). Exploitable biomass and numbers, which were comprised of Gray Snapper age-2 or older, averaged 31,771 metric tons and 37,409,629 Gray Snapper, respectively. Exploitable biomass was lowest in 1997 at 15,138 metric tons and peaked in 1945 at 55,844 metric tons, whereas exploitable numbers ranged from 22,122,100 Gray Snapper in 1997 to 54,481,600 Gray Snapper in 1945 (**Table 13**). SSB averaged 12,688 metric tons, and ranged from 6,324 metric tons in 1997 to 21,719 metric tons in 1945 (**Figure 59**). Both total biomass and SSB show a steady decline from 1945 to the early 1980s, followed by a plateauing off in the 1980s to early 2000s. Starting in the late 2000s, biomass trends show a gradual increase to the terminal year.

The SSB ratio averaged 0.58, and ranged from 0.29 in 1997 to 1 in 1945 (**Table 13**). Estimated SSB ratio has been above 30% since 1997, with estimated spawning stock biomass in the most recent year (2020) predicted to be at 48% of the corresponding unfished spawning stock biomass (**Table 13**).

4.7. Model Fit and Residual Analysis

4.7.1. Landings

Landings for the Commercial Vertical Line, Commercial Longline and Commercial Nets & Traps fleets were fit almost exactly given their relatively small SEs (**Table 14-16, Figure 60**). Given the larger SEs assigned to the recreational fleet landings, there were considerable differences between input and predicted landings in numbers (**Table 17-19, Figure 60**). **Section 4.7.2, Figure 66**). In general, there was a similar fit to the landings data in SEDAR51 compared with SEDAR75, although direct comparisons between the commercial fleets are not possible due to the change in fleet structure (**Figure 60**).

4.7.2. Discards

The time series of commercial discards begins in 1993, three years after the implementation of the first minimum size limit. Observed and expected values are summarized in **Tables 20-21 and Figure 61**. Generally, the discards were relatively low for both the Commercial Vertical Line and Commercial Longline fleets, though the Commercial Longline had fewer discards than the Commercial Vertical Line. Discards were estimated with a large assumed uncertainty, and therefore were characterized by large confidence intervals for the two commercial fleets with discards (**Figures 61-63**). For the Commercial Vertical Line fleet, the model expected fewer discards than observed prior to 2010. For the Commercial Longline fleet, the model had a difficult time predicting discards given the lack of discard information.

The time series of discards for the recreational fleets begins in 1981 (**Tables 22-24, Figures 63-65**). The model was able to fit discard observations relatively well throughout the time series for recreational fleets. Though, for Recreational Shore (**Figure 64**), the model overestimated the expected discards in 1991.

There was an improvement in fits to the discard data in SEDAR51 as compared with SEDAR75, however direct comparisons between the discard inputs are not possible due to how discards were modeled in the two assessments (**Figure 66**), as well as changes in the discard estimation methodology.

4.7.3. Indices

Observed and predicted CPUE are provided in **Tables 25 and 26** and **Figure 67**.

The fits to the two fishery dependent indices, Private CPUE and Shore CPUE, were similar (root mean squared error [RMSE] = 0.304 and 0.444, respectively). Both indices had similar trends (**Figure 67**), however Shore CPUE exhibited a more pronounced increasing trend.

Of the fishery independent indices, the model fit better to the FWRI age 1 (RMSE= 0.81) than to the FWRI age 0 (RMSE= 0.516) (**Figure 67**). The increased time series for Combined Video Survey improved the fit as compared to SEDAR51 0.425.

4.7.4. Length Compositions

Model fits to the retained length composition data are provided in **Figures 69-76**.

The aggregate fit to the retained length composition data were fairly similar between SEDAR51 and SEDAR75 (**Figure 75**), however, the available years for the length composition data for some fleets in SEDAR75 were decreased as compared to SEDAR51 (**Figure 76**) due to removing some years of length compositions where age compositions were available. This is result of including age compositions in SEDAR75.

Model fits to the survey length composition data are provided in **Figures 77-79**.

4.7.5. Age Compositions

Model fits to the age composition data are provided in **Figures 80-85**. Generally, the model fit the age composition well however there were some residual patterns observed in Commercial Vertical Line and Recreational Private.

Across all fleets, there was a tradeoff between fitting to the weighted retained length compositions and fitting to the nominal age compositions. Overall, the model fit more closely to the length compositions due to the larger sample sizes and larger contribution to the total likelihood.

4.8. Model Diagnostics

4.8.1. Correlation Analysis

A summary of correlations for the base model parameters considered as outliers is contained in **Table 27**. Given the highly parametrized nature of this model, some parameters were mildly correlated (correlation coefficient > 70%) and two combinations (the Recreational Charter & Headboat age selectivity parameters and the Combined Video Survey length selectivity parameters) displayed a strong correlation (> 95%; **Table 27**). Correlation among many of these parameters is not surprising, especially for the selectivity parameters, because the parameters of

selectivity functions are inherently correlated (i.e., as the value of one parameter changes the other value will compensate). Moderate correlations occurred between the parameters defining the peak and the width of the ascending and/or descending limb of the double normal selectivity functions for some fleets.

4.8.2. Profile Likelihoods

The total likelihood component from the $\ln R_0$ likelihood profile indicates that the global solution for this parameter is approximately 10.1 (**Figure 86**). The SEDAR75 Base Model estimating $\ln R_0$ at 10.052 (CV = 0.005; **Table 10**). Almost all data sources supported this estimate, with the exception of the length data which supported a slightly lower $\ln R_0$ near 10.0.

4.8.3. Jitter Analysis

A jitter analysis was conducted using a jitter value of 0.1. With this procedure, the starting model parameter values are randomly adjusted by 10% from the SEDAR75 Base Model best fit over 100 runs (**Figure 87**). No better solution was found, though the analysis shows there is some sensitivity to starting values.

4.8.4. Retrospective Analysis

Results from the retrospective analysis do not indicate any directional retrospective patterns. As the last few years of data are peeled off, the model estimates of SSB, recruitment and F in each successive terminal year do not change by a large margin (and remain within the confidence intervals; **Figure 88**).

4.8.5. Additional Diagnostics

The SEDAR75 displayed acceptable RMSE (<30%) for the joint residuals for the mean age and mean length data sources (**Table 29**). Residuals revealed some conflict in indices of abundance and mean age (evident by colored vertical lines in opposite directions) and trends in the residuals (evident by Loess smoothed line; **Figure 89-Figure 91**). The lowest RMSE was exhibited for the length composition, which exhibited the smallest residuals but did reveal some conflicts (**Table 29; Figure 89**). Runs test results revealed evidence of non-randomly distributed residuals for the Recreational Private and Combined Video Survey lengths compositions, FWRI 1 index of abundance, and Recreational Charter & Headboat age compositions (**Table 30; Figure 95**). A few outliers (evident by red points) were identified in residuals for mean age for all fleets, in the index residuals for the Recreational Private, Recreational Shore and the Combined Video Survey, and in residuals for length compositions for the Recreational Private fleet (**Figure 95**). Superior prediction skill (<1) was evident over the naive baseline forecast for the Recreational Private index (**Figure 92**), mean age for all fleets (**Figure 93**), and mean length for the commercial fleets (**Figure 94; Table 31**).

4.8.6. Bridging analysis

The general flow of model building runs that led to the final SEDAR75 base model is shown in **Table 32**. Changes in estimated quantities are shown in **Table 32** and **Figures 98-101**.

The SEDAR51 model that used the SS3.24_S version was successfully converted to the new SS3.30 version without any issues (Step 1). Key derived quantities and important parameters (e.g., S/R parameters, growth) were estimated similarly in SS3.30. The run with the corrected commercial landings (Run 2) had slight differences in biomass and annual exploitation rates from the S51 Base Model (**Figures 102-103**). When the new revised MRIP-FES landings and discard data were substituted for the recreational fleet inputs in Step 3, estimates of virgin SSB (SSB_0) and virgin recruitment (R_0) increased by 10%. Substituting the new MRIP-FES estimates generally increased SSB across the time series without drastically affecting the trajectory of the stock over the data period. The next step in the bridging analysis, Step 4, involved updating all new data streams (changing the terminal year to 2020, altering the commercial fleet structure to the SEDAR 51 Data Workshop approved fleets and including all new catches, discards, indices, length comps and associated CVs). The following step, Step 5, added in the fishery independent length compositions. Step 6 added all available age composition data and subsequently removed some length composition years with available age composition. Step 7 had the maximum age of 21 used in SEDAR 51 while step 8 adjusted the maximum age from 21 to the Terms of Reference requested maximum age of 28. Many additional runs were conducted during the development of SEDAR75 but the above mentioned 8 steps show the most important stepping stones that govern the changes observed between the SEDAR51 and SEDAR75. Most changes affecting the trajectory of the stock occurred prior to step 7.

4.8.6. Sensitivity Model Runs

Results for the sensitivity runs summarized in **Section 3.4.6** are discussed below.

Start Year of 1986

Truncating the data inputs to begin in 1986 greatly increased the biomass of Gray Snapper however this run is highly unstable and did not converge, as such the results are not included.

Natural Mortality

Using the SEDAR51 low and high M vectors equated to expected patterns in the spawning stock biomass and exploitation rates, however the low natural mortality run's Hessian matrix was not positive definite. This indicates that the available data does not support the lower M . (**Figure 104**).

5. Discussion

The SEDAR75 Gray Snapper assessment included several important changes to data inputs and model parameterization that affected the assessment results including the following:

1. incorporating the MRIP-FES in estimation of recreational landings and discards;
2. including fishery independent length compositions;
3. including age compositions for four fishing fleets;
4. revising historical landings for the recreational fleets;
5. applying the SEFSC's improved approach for estimating commercial discards;

6. updating estimates of maturity;
7. updating estimates of maximum age;
8. updating estimates of the growth curve;
9. applying an internal re-weighting approach to both age and length compositions.

The most significant of the SEDAR51 to SEDAR75 model changes (data or model configuration) was the addition of age compositions for the Commercial Vertical Line, Commercial Longline, Recreational Private and Recreational Charter & Headboat fleets as well the inclusion of fishery independent length compositions. (**Figure 98**), which ultimately scaled the population size upward due to higher estimated biomass. The change from the maximum age of 21 to 28 did not have a large impact on the overall assessment results and estimates of parameters (growth rate, R_0 , etc.) or key derived quantities. Converting the previous SEDAR 33 SS 3.24s model to the upgraded SS 3.30 version had virtually no impact on model results but was seen as an overall improvement in the assessment as the updated SS version (3.30_19) allows even greater flexibility in handling a number of processes including data weighting and projections.

The SEDAR75 model fit most of the data sources well with no major residual patterns and the fits to the discards were much improved from SEDAR51. The dominant data inputs were the length and age compositions as these produced the greatest impact on the model fit (as measured in the contribution to the total likelihood). There were only a few parameters with high correlations and they did not appear to be the source of any major model stability issues. No substantial retrospective patterns are present in the model fits, indicating internal consistency within the model. Likelihood profiles on R_0 showed general agreement between data sources.

It is important to note that uncertainties remain in some components of the Gray Snapper data series used in the assessment. The landings data are dominated by the recreational fishery, and recreational landings are more uncertain than commercial data. Additionally, before 1981, recreational data are estimated using a hindcasting procedure that is very sensitive to the assumed catches in 1981-1985 (as was shown by the impact that a single year's peak in Recreational Private landings had on the historical time series). The largest removals for Gray Snapper come from the Recreational Private and Recreational Shore fisheries. However, the length and age composition for the Recreational Shore fleet are very sparse given the magnitude of reported landings. Age compositions from the shore mode were not sufficient to be included in the model. Additionally, there was no discard compositions for any fleet.

Though scales differ (largely due to the introduction of MRIP-FES estimates), the SEDAR75 Gray Snapper assessment and SEDAR51 assessments both predict similar stock trends. This trend is characterized by a steady decline in stock biomass associated with a steady increase in exploitation rates from the beginning of the time series (1945) to 1980, followed by a period of relatively steady stock size until the mid-2000s when the stock begins to increase and exploitation rates from the recreational fleets ramp up (**Figure 59 and 37**). The SEDAR51 assessment found that the Gray Snapper GOM stock had been undergoing overfishing since 1976; with the inclusion of age compositions, fishery independent length compositions and updated life history estimates, the SEDAR75 assessment time series indicates that Gray Snapper

has not undergone overfishing, although the spawning stock biomass did decline and exhibited a similar overall trend as SEDAR51.

A number of research questions were raised during the SEDAR75 assessment process. While attempts were made to address these questions through sensitivity runs and preliminary data exploration, the Operational nature of this assessment did not leave enough time to thoroughly evaluate each and every one of these questions. The SEFSC strongly recommends that these topics (listed in **Section 8**) be more thoroughly examined during a future assessment with targeted topical working works.

Overall, the SEDAR75 base model is improved since the SEDAR51 Benchmark assessment, and it incorporates the best available data and addressed modeling issues evident in the prior assessment.

6. Projections

6.1. Introduction

The SEDAR75 projections were run to obtain the overfishing limit (OFL) for the F_{SPR30} fishing mortality scenario and the acceptable biological catch (ABC, 75% F_{SPR30}).

6.2. Projection methods

The simulated dynamics used for projections assumed nearly identical parameter values and population dynamics as the SS base model. **Table 33** provides a summary of projection settings. Projections were run assuming that selectivity, discarding and retention were the same as the terminal year. Forecast recruitment values were derived from the model-estimated Beverton-Holt stock-recruitment relationship.

The terminal year of the SEDAR75 assessment was 2020 and the first year of management advice was 2024. Retained catch for the interim years (2021-2023) used the average of the last 3 years of retained catches, see **Table 33**.

F_{SPR30} was determined using a long-term 100-year projection assuming that equilibrium was obtained over the last 10 years (2110-2120). For the OFL projection, the F_{SPR30} was applied to the stock starting in 2024 (**Table 33**).

The minimum stock size threshold (MSST) was determined by multiplying the reference spawning stock biomass, SSB_{SPR30} , by 0.5 (per the SEDAR75 TORS) and was used to determine stock status (**Table 34**). The maximum fishing mortality threshold (MFMT) was equivalent to the harvest rate (F_{SPR30} ; total biomass killed age 2+ / total biomass age 2+) that achieved SSB_{FSPR30} , and was used to assess whether overfishing was occurring in a given year (**Table 34**).

Once the proxy values were calculated, 2020 stock status was used to determine whether a rebuilding plan was required (i.e., if $SSB < MSST$ then Gulf of Mexico Gray Snapper would be considered overfished and a rebuilding plan would be required). As Gray Snapper was not determined to be overfished, no rebuilding plan is needed.

6.3. Projection results

Following the Terms of Reference, benchmarks and reference points were calculated by assuming an SSB defined in terms of males and females combined.

6.3.1. Biological Reference Points

The following status determination criteria (SDCs) were adopted for Gulf of Mexico Gray Snapper:

- MSY proxy = yield at F_{SPR30} ,
- $MSST = 0.5 * SSB_{SPR30}$ (Amendment 51),
- $MFMT = F_{SPR30}$ or $F_{rebuild}$ if overfished.

The harvest rate that results in SSB_{SPR30} over the long-term (100 years) was 0.134 for SSB and (Table 34). The resulting $SSB_{F_{SPR30}}$ was 6,477.13 metric tons, and the minimum stock size threshold (MSST) was 3,238.57 metric tons (Figure 105).

6.3.2. Stock Status

Benchmarks and reference points are shown in Table 34. Detailed time series are presented in Table 35. The Gulf of Mexico Gray Snapper stock is not undergoing overfishing ($F_{current} < MFMT$) and is not overfished ($SSB_{2020} > MSST$) based on the definition of MSST ($0.5 * SSB_{SPR30}$), $F_{current}$ (geometric mean of the harvest rate over 2018-2020 and MFMT ($F_{SPR30\%}$) for the final SEDAR75 base model (Table 34). The terminal year SSB is also above the recovery target, SSB_{SPR30} (Figure 105). In 2020, SSB was 160% of the biomass level needed to support MSY. From 2018 to 2020 the estimated stock harvest rate, using the geometric mean, was 0.091, which was equivalent to 68% (Table 34, Figure 105).

The Kobe plot (Figure 106) indicates that over the time horizon of the assessment (i.e., 1945 - 2020), the stock has not ever experienced overfishing.

6.3.3. Overfishing Limits and ABC projections

OFL projection results are provided in Table 36 and ABC projections results are provided in Table 37. Both the OFL and ABC projections are provided in Figure 107. Forecasts begin in 2024 because management based on this stock assessment is not expected to begin until that time. Since the stock is not overfished (Table 34), a rebuilding projection is not needed.

Since the stock is currently below the F_{SPR30} target, forecasts indicate that a reduction in yield is not required in the near-term.

7. Acknowledgements

The SEDAR75 Operational for Gulf of Mexico Gray Snapper would not have been possible without the efforts of the numerous NMFS, SEFSC, SERO, and GMFMC staff along with the many academic and research partners involved throughout the Gulf of Mexico. The following agencies contributed to the assessment and deserve notable attention and thanks for efforts extended to developing data inputs: NOAA SEFSC Fisheries Statistics Division (FSD), NOAA

SEFSC Gulf Fisheries Branch of the Sustainable Fisheries Division, NOAA SEFSC Panama City Laboratory, University of Florida, NOAA SEFSC Mississippi Laboratories, NOAA Southeast Regional Office (SERO), Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute, Cooperative Institute for Marine and Atmospheric Studies, Rosenstiel School for Marine and Atmospheric Science, NOAA Atlantic Oceanographic and Meteorological Laboratory, and the Gulf States Marine Fisheries Commission. Special thanks are also extended to the Topical Working Group members for their assistance and guidance throughout the process. Special thanks are extended to Dr. Richard Methot and team for continued discussions and modifications to the Stock Synthesis model. Thanks are extended to Skyler Sagarese and Lisa Ailloud for developing R code to implement the Markdown versions of the majority of the tables and figures.

8. Research Recommendations

Recommendations for considerations of future research are provided below and do not indicate any particular order of priority.

Recreational Landings and Discards data

- Further develop best practices for correcting for prominent peaks and troughs in the earlier part of the time series where uncertainty is high and catch/discard estimates are driven by few but influential intercept records.
- Develop estimates of uncertainty around Headboat discard estimates.

Age and length composition

- Quantify and evaluate appropriate modeling and weighting procedures of length and age compositions to ensure age and length composition inputs are representative of the segment of the population being modeled.
- Improve sampling programs for shore based recreational fishing modes to obtain length and age data for that mode.

Selectivity and catchability of the commercial fleets

- Further investigate and quantify changes in selectivity/catchability through time to improve fit to the discards and length compositions in recent years.
- Continue data collection from observer programs or electronic monitoring programs.

Selectivity and retention of the recreational fleets

- Further investigate and quantify changes in selectivity/catchability through time to improve fit to the length compositions across the time series.

Landings and Discards

- Explore approaches for assigning uncertainty estimates to commercial landings and revisit estimation of historic landings.
- Further investigate best practices for converting historical recreational landings from numbers to weight.

Recreational CPUE indices

- Additional research is needed to investigate if assumptions are appropriate across full time series (e.g., targeting, trip length, effects of various regulations on gray snapper as well on other species i.e. red snapper).

Natural mortality

- Explore ways to better reflect uncertainty about the mortality at age vector.

9. References

- Carvalho F, H Winker, D Courtney, M Kapur, L Kell, M Cardinale, M Schirripa, T Kitakado, D Yemane, KR Piner and MN Maunder. 2021. A cookbook for using model diagnostics in integrated stock assessments. *Fisheries Research* 240: 105959. doi: 10.1016/j.fishres.2021.105959
- Dichmont, CM, RA Deng, AE Punt, J Brodziak, YJ Chang, JM Cope, JN Ianelli, CM Legault, RD Methot, CE Porch and MH Prager. 2016. A review of stock assessment packages in the United States. *Fisheries Research* 183:447-460.
- Francis RICC. 2011. Data weighting in statistical fisheries stock assessment models. *Canadian Journal of Fisheries and Aquatic Sciences*. 68:1124-1138.
- Francis RICC, RJ Hurst and JA Renwick. 2003 Quantifying annual variation in catchability for commercial and research fishing. *Fishery Bulletin* 101(2):293-304.
- Hoenig, J.M., 1983. Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin*, 82(1), pp.898-903.
- Hulson P-J, D Hanselman, and T Quinn. 2012. Determining effective sample size in integrated age-structured assessment models. *ICES Journal of Marine Science*, 69:281-292.
- Lorenzen K. 2000. Allometry of natural mortality as a basis for assessing optimal release size in fish-stocking programmes. *Canadian Journal of Fisheries and Aquatic Sciences* 57(12):2374-2381.
- Methot RD and IG Taylor. 2011. Adjusting for bias due to variability of estimated recruitments in fishery assessment models. *Canadian Journal of Fisheries and Aquatic Sciences*, 68(10):1744-1760.
- Methot RD and CR Wetzel. 2013. Stock synthesis: a biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research* 142:86–99.
- Methot RD, CR Wetzel, IG Taylor and K Doering. 2020. Stock Synthesis User Manual Version 3.30.16. NOAA Fisheries, Seattle Washington. 225 pp.
- Quinn T and R Deriso. 1999. Quantitative fish dynamics. Oxford University Press, New York.
- Sauls B, T Cross, L Barbieri and M Guyas. 2019. Results from the first year of an exempted fishing permit (18-SERO-01) for state management of Red Snapper recreational harvest in Florida. Final report submitted to: NOAA Southeast Regional Office St. Petersburg, FL. Accessible at: https://media.fisheries.noaa.gov/dam-migration/2018_efp_fl_lchamp_final_report_508.pdf

Stephens A and A MacCall. 2004. A multispecies approach to subsetting logbook data for purposes of estimating CPUE. *Fisheries Research*, 70: 299-310.

Then AY, JM Hoenig, NG Hall, and DA Hewitt. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES Journal of Marine Science* 72(1):82-92.

Thorson JT, KF Johnson, RD Methot and IG Taylor. 2017. Model-based estimates of effective sample size in stock assessment models using the Dirichlet-multinomial distribution. *Fisheries Research* 192: 84–93. [doi:10.1016/j.fishres.2016.06.005](https://doi.org/10.1016/j.fishres.2016.06.005).

Taylor IG, KL Doering, KF Johnson, CR Wetzel and IJ Stewart, 2021. Beyond visualizing catch-at-age models: Lessons learned from the r4ss package about software to support stock assessments, *Fisheries Research* 239:105924. <https://doi.org/10.1016/j.fishres.2021.105924>.

Tables

Table 1. Conversion factors used to convert fork length (FL) in centimeters to whole weight (GW) in kilograms and to convert maximum total length (MaxTL) to fork length (FL) for Gulf of Mexico Gray Snapper males and females combined. Model fit criteria: linear regression models r^2 and non-linear regression models' residual square error (RSE).

| Model | N | Range | R2 or RSE value | Years |
|--|--------|---------------------|-----------------|-----------|
| $W = 1.43 \times 10^{-5} \times (FL^{3.02})$ | 10,954 | FL (cm): 9.6 – 71.5 | 0.130 | 1991-2015 |
| | | WW (kg): 1.8-6.6 | | |
| $FL = 0.36 + \max_TL \times 0.93$ | 3,050 | Max TL: 10.1 – 63.5 | 0.999 | 1991-2015 |
| | | FL: 9.6 – 60.5 | | |

Table 2. Growth parameters recommended for Gulf of Mexico Gray Snapper. The von Bertalanffy parameters (Linf, K, t0) were updated during SEDAR 75 Life History Topical Working Group using a larger sample of fish (SEDAR75-WP-05).

| Parameter | Value |
|-------------------------|-------|
| Linf (cm FL) | 61 |
| K (year ⁻¹) | 0.11 |
| t0 (year) | -1.47 |
| CV _{Amin} | 0.15 |
| CV _{Amax} | 0.19 |

Table 3. Age-specific natural mortality (per year) for Gulf of Mexico Gray Snapper. Female and male natural mortality were assumed equivalent.

| Age | Base M | Upper | Lower |
|-----|--------|-------|-------|
| 0 | 0.50 | 0.57 | 0.44 |
| 1 | 0.36 | 0.40 | 0.31 |
| 2 | 0.28 | 0.32 | 0.25 |
| 3 | 0.24 | 0.27 | 0.21 |
| 4 | 0.22 | 0.24 | 0.19 |
| 5 | 0.20 | 0.22 | 0.17 |
| 6 | 0.18 | 0.21 | 0.16 |
| 7 | 0.17 | 0.19 | 0.15 |
| 8 | 0.16 | 0.18 | 0.14 |
| 9 | 0.16 | 0.18 | 0.14 |
| 10 | 0.15 | 0.17 | 0.13 |
| 11 | 0.15 | 0.17 | 0.13 |
| 12 | 0.15 | 0.16 | 0.13 |
| 13 | 0.14 | 0.16 | 0.13 |
| 14 | 0.14 | 0.16 | 0.12 |
| 15 | 0.14 | 0.16 | 0.12 |
| 16 | 0.14 | 0.15 | 0.12 |
| 17 | 0.14 | 0.15 | 0.12 |
| 18 | 0.14 | 0.15 | 0.12 |
| 19 | 0.13 | 0.15 | 0.12 |
| 20 | 0.13 | 0.15 | 0.12 |
| 21 | 0.13 | 0.15 | 0.12 |

Table 3 Continued. Age-specific natural mortality (per year) for Gulf of Mexico Gray Snapper. Female and male natural mortality were assumed equivalent.

| Age | Base M | Upper | Lower |
|-----|--------|-------|-------|
| 22 | 0.13 | 0.15 | 0.12 |
| 23 | 0.13 | 0.15 | 0.12 |
| 24 | 0.13 | 0.15 | 0.12 |
| 25 | 0.13 | 0.15 | 0.11 |
| 26 | 0.13 | 0.15 | 0.11 |
| 27 | 0.13 | 0.15 | 0.11 |
| 28 | 0.13 | 0.15 | 0.11 |

Table 4. Gulf of Mexico Gray Snapper commercial landings in pounds gutted weight. Landings by “Other” gears were lumped into the Commercial Longline fleet for input into the stock assessment. In the absence of uncertainty estimates provided at the SEDAR75 DW, commercial landings were assigned a log-scale SE of 0.05.

| Year | Vertical Line | Longline | Nets & Traps | Total |
|------|---------------|----------|--------------|---------|
| 1962 | 321,700 | 0 | 15,000 | 338,000 |
| 1963 | 277,900 | 500 | 8,900 | 287,300 |
| 1964 | 297,200 | 0 | 13,800 | 311,000 |
| 1965 | 327,000 | 0 | 47,400 | 374,400 |
| 1966 | 279,100 | 0 | 30,900 | 310,000 |
| 1967 | 318,600 | 0 | 54,100 | 372,700 |
| 1968 | 398,500 | 0 | 72,200 | 470,700 |
| 1969 | 373,100 | 0 | 107,300 | 480,400 |
| 1970 | 335,800 | 0 | 104,400 | 440,200 |
| 1971 | 358,900 | 0 | 108,500 | 467,400 |
| 1972 | 402,400 | 0 | 126,500 | 528,900 |
| 1973 | 386,300 | 0 | 169,300 | 555,600 |
| 1974 | 392,700 | 0 | 194,300 | 587,000 |
| 1975 | 256,500 | 0 | 228,000 | 484,500 |
| 1976 | 473,000 | 0 | 125,300 | 598,300 |
| 1977 | 268,700 | 0 | 356,600 | 625,300 |
| 1978 | 233,500 | 0 | 432,800 | 666,300 |
| 1979 | 274,500 | 0 | 398,800 | 673,300 |
| 1980 | 508,121 | 19,744 | 178,153 | 706,018 |
| 1981 | 508,326 | 20,194 | 161,502 | 690,022 |
| 1982 | 676,569 | 47,193 | 174,049 | 897,811 |
| 1983 | 670,917 | 85,909 | 176,333 | 933,159 |

Table 4 Continued. Gulf of Mexico Gray Snapper commercial landings in pounds gutted weight. Landings by “Other” gears were lumped into the Commercial Longline fleet for input into the stock assessment. In the absence of uncertainty estimates provided at the SEDAR75 DW, commercial landings were assigned a log-scale SE of 0.05.

| Year | Vertical Line | Longline | Nets & Traps | Total |
|------|---------------|----------|--------------|---------|
| 1984 | 470,393 | 47,027 | 267,889 | 785,309 |
| 1985 | 373,337 | 34,739 | 218,295 | 626,371 |
| 1986 | 408,479 | 41,312 | 169,205 | 618,996 |
| 1987 | 495,832 | 45,597 | 137,059 | 678,488 |
| 1988 | 316,081 | 31,489 | 97,781 | 445,351 |
| 1989 | 380,372 | 46,158 | 119,517 | 546,047 |
| 1990 | 306,970 | 40,907 | 43,677 | 391,554 |
| 1991 | 380,636 | 43,384 | 44,050 | 468,070 |
| 1992 | 355,328 | 50,249 | 22,249 | 427,826 |
| 1993 | 429,768 | 78,857 | 13,088 | 521,713 |
| 1994 | 520,439 | 8,767 | 35,284 | 564,490 |
| 1995 | 414,993 | 9,176 | 18,324 | 442,493 |
| 1996 | 394,822 | 6,781 | 8,995 | 410,598 |
| 1997 | 378,117 | 7,477 | 9,868 | 395,462 |
| 1998 | 294,609 | 8,357 | 10,543 | 313,509 |
| 1999 | 264,401 | 12,812 | 7,638 | 284,851 |
| 2000 | 280,098 | 12,558 | 5,486 | 298,142 |
| 2001 | 299,159 | 10,984 | 4,766 | 314,909 |
| 2002 | 353,982 | 13,719 | 3,747 | 371,448 |
| 2003 | 310,329 | 9,862 | 4,790 | 324,981 |
| 2004 | 323,835 | 14,914 | 1,901 | 340,650 |
| 2005 | 290,736 | 12,932 | 1,532 | 305,200 |

Table 4 Continued. Gulf of Mexico Gray Snapper commercial landings in pounds gutted weight. Landings by “Other” gears were lumped into the Commercial Longline fleet for input into the stock assessment. In the absence of uncertainty estimates provided at the SEDAR75 DW, commercial landings were assigned a log-scale SE of 0.05.

| Year | Vertical Line | Longline | Nets & Traps | Total |
|------|---------------|----------|--------------|---------|
| 2006 | 258,306 | 12,824 | 724 | 271,854 |
| 2007 | 197,218 | 11,907 | 419 | 209,544 |
| 2008 | 205,342 | 14,883 | 972 | 221,197 |
| 2009 | 252,856 | 17,284 | 887 | 271,027 |
| 2010 | 207,739 | 6,815 | 88 | 214,642 |
| 2011 | 236,221 | 12,572 | 91 | 248,884 |
| 2012 | 245,632 | 13,376 | 372 | 259,380 |
| 2013 | 223,199 | 12,418 | 1,342 | 236,959 |
| 2014 | 278,837 | 18,145 | 571 | 297,553 |
| 2015 | 238,808 | 26,216 | 224 | 265,248 |
| 2016 | 241,072 | 25,401 | 178 | 266,651 |
| 2017 | 183,195 | 21,077 | 184 | 204,456 |
| 2018 | 198,231 | 12,239 | 81 | 210,551 |
| 2019 | 165,294 | 13,910 | 248 | 179,452 |
| 2020 | 134,251 | 10,927 | 368 | 145,546 |

Table 5. Gulf of Mexico Gray Snapper recreational landings in numbers. Landings from Private, Shore and Private/Shore (from LA Creel) were lumped into the Recreational Shore fleet for input into the stock assessment. Recreational landings were assigned a log-scale SE of 0.10.

| Year | Headboat | Charter | Private | Shore | Private/Shore |
|------|----------|---------|------------|-----------|---------------|
| 1981 | 74,647 | 111,197 | 2,333,365 | 1,261,095 | |
| 1982 | 227,421 | 349,426 | 1,835,937 | 1,415,808 | |
| 1983 | 164,577 | 236,802 | 535,256 | 1,183,441 | |
| 1984 | 47,379 | 60,960 | 8,387,582* | 1,210,392 | |
| 1985 | 51,490 | 50,554 | 974,810 | 597,227 | |
| 1986 | 79,308 | 111,134 | 842,745 | 1,025,696 | |
| 1987 | 49,452 | 76,059 | 1,772,352 | 1,169,326 | |
| 1988 | 47,366 | 18,910 | 1,326,650 | 1,222,560 | |
| 1989 | 73,169 | 61,904 | 2,421,003 | 2,311,290 | |
| 1990 | 61,181 | 27,903 | 1,953,384 | 1,117,930 | |
| 1991 | 55,480 | 173,700 | 1,750,093 | 3,147,714 | |
| 1992 | 54,229 | 161,549 | 1,318,498 | 958,933 | |
| 1993 | 67,384 | 57,563 | 1,731,418 | 973,239 | |
| 1994 | 75,008 | 89,189 | 1,424,975 | 667,381 | |
| 1995 | 57,153 | 52,342 | 1,803,451 | 624,697 | |
| 1996 | 44,271 | 74,482 | 1,322,923 | 941,554 | |
| 1997 | 46,448 | 24,914 | 1,309,017 | 468,198 | |
| 1998 | 61,575 | 129,692 | 1,830,662 | 451,372 | |
| 1999 | 66,851 | 91,085 | 1,574,755 | 356,449 | |
| 2000 | 40,178 | 73,275 | 1,365,867 | 498,612 | |

*The 1984 peak in Private landings was replaced with values from 1986 from west FL, wave 6, <= 10 miles. The new value is 2,578,761 fish.

Table 5 Continued. Gulf of Mexico Gray Snapper recreational landings in numbers. Landings from Private, Shore and Private/Shore (from LA Creel) were lumped into the Recreational Shore fleet for input into the stock assessment. Recreational landings were assigned a log-scale SE of 0.10.

| Year | Headboat | Charter | Private | Shore | Private/Shore |
|------|----------|---------|-----------|-----------|---------------|
| 2001 | 60,973 | 138,611 | 1,591,273 | 632,727 | |
| 2002 | 60,109 | 82,427 | 1,298,275 | 304,916 | |
| 2003 | 53,429 | 168,184 | 1,763,137 | 441,768 | |
| 2004 | 50,223 | 145,878 | 2,173,806 | 502,109 | |
| 2005 | 53,723 | 138,132 | 1,490,890 | 517,283 | |
| 2006 | 54,165 | 139,058 | 1,385,621 | 265,176 | |
| 2007 | 37,603 | 160,168 | 1,645,434 | 384,455 | |
| 2008 | 37,056 | 182,339 | 2,297,914 | 697,753 | |
| 2009 | 53,355 | 239,116 | 2,275,323 | 498,188 | |
| 2010 | 37,569 | 111,104 | 1,193,182 | 129,274 | |
| 2011 | 55,595 | 119,638 | 1,272,516 | 387,914 | |
| 2012 | 67,980 | 105,299 | 3,332,876 | 682,838 | |
| 2013 | 55,426 | 251,344 | 2,859,362 | 620,783 | |
| 2014 | 78,642 | 269,080 | 3,398,927 | 980,912 | 108,854 |
| 2015 | 69,075 | 201,111 | 2,665,812 | 642,119 | 137,871 |
| 2016 | 71,638 | 310,702 | 2,651,206 | 977,178 | 93,458 |
| 2017 | 66,658 | 259,867 | 1,798,646 | 1,239,436 | 62,868 |
| 2018 | 74,794 | 252,122 | 1,938,561 | 1,106,709 | 57,027 |
| 2019 | 71,629 | 315,047 | 1,744,558 | 1,602,805 | 68,591 |
| 2020 | 57,159 | 251,824 | 2,424,618 | 1,144,350 | 39,881 |

Table 6. Gulf of Mexico Gray Snapper commercial discards in numbers with associated log-scale standard errors (SE) input into the assessment model. Discards refer to the total number of fish discarded before applying the discard mortality rate.

| Year | Vertical Line | Vertical Line SE | Longline | Longline SE |
|-------|---------------|------------------|----------|-------------|
| 1,993 | 3,203 | 0.425 | 43 | 0.470 |
| 1,994 | 4,210 | 0.437 | 44 | 0.458 |
| 1,995 | 3,918 | 0.430 | 28 | 0.456 |
| 1,996 | 3,368 | 0.408 | 28 | 0.426 |
| 1,997 | 4,212 | 0.412 | 27 | 0.434 |
| 1,998 | 3,599 | 0.407 | 18 | 0.446 |
| 1,999 | 4,539 | 0.417 | 45 | 0.467 |
| 2,000 | 3,426 | 0.389 | 45 | 0.518 |
| 2,001 | 3,619 | 0.387 | 49 | 0.534 |
| 2,002 | 4,053 | 0.389 | 70 | 0.528 |
| 2,003 | 3,303 | 0.371 | 52 | 0.484 |
| 2,004 | 3,577 | 0.387 | 68 | 0.523 |
| 2,005 | 3,299 | 0.403 | 46 | 0.500 |
| 2,006 | 3,124 | 0.409 | 72 | 0.512 |
| 2,007 | 2,500 | 0.495 | 44 | 0.666 |
| 2,008 | 2,371 | 0.495 | 53 | 0.666 |
| 2,009 | 3,351 | 0.495 | 20 | 0.666 |
| 2,010 | 1,302 | 0.580 | 21 | 0.666 |
| 2,011 | 1,181 | 0.580 | 39 | 0.666 |
| 2,012 | 1,372 | 0.580 | 40 | 0.666 |
| 2,013 | 1,216 | 0.580 | 52 | 0.666 |
| 2,014 | 1,517 | 0.580 | 183 | 0.380 |
| 2,015 | 1,623 | 0.580 | 211 | 0.380 |

Table 6 Continued. Gulf of Mexico Gray Snapper commercial discards in numbers with associated log-scale standard errors (SE) input into the assessment model. Discards refer to the total number of fish discarded before applying the discard mortality rate.

| Year | Vertical Line | Vertical Line SE | Longline | Longline SE |
|-------|---------------|------------------|----------|-------------|
| 2,016 | 1,610 | 0.580 | 235 | 0.380 |
| 2,017 | 1,472 | 0.580 | 152 | 0.500 |
| 2,018 | 1,213 | 0.580 | 125 | 0.500 |
| 2,019 | 1,188 | 0.580 | 110 | 0.500 |
| 2,020 | 1,145 | 0.580 | 78 | 0.500 |

Table 7. Gulf of Mexico Gray Snapper recreational discards in numbers with associated log-scale standard errors (SE) input into the assessment model. Discards refer to the total number of fish discarded before applying the discard mortality rate.

| Year | Charter | Charter SE | Headboat | Headboat SE | Private | Private SE | Shore | Shore SE |
|------|---------|------------|----------|-------------|-----------|------------|-----------|----------|
| 1981 | 18,249 | 0.675 | 10,628 | 0.675 | 645,683 | 0.617 | 835,879 | 0.481 |
| 1982 | 113,815 | 0.506 | 71,184 | 0.506 | 608,777 | 0.538 | 446,852 | 0.421 |
| 1983 | 44,359 | 0.808 | 27,744 | 0.808 | 99,646 | 0.455 | 1,652,615 | 0.631 |
| 1984 | 6,071 | 0.472 | 960 | 0.833 | 660,573 | 0.646 | 295,909 | 0.403 |
| 1985 | 11,467 | 0.472 | 5,389 | 0.578 | 914,096 | 0.455 | 344,325 | 0.367 |
| 1986 | 12,678 | 0.385 | | | 561,141 | 0.455 | 118,816 | 0.412 |
| 1987 | 6,120 | 0.530 | | | 430,237 | 0.385 | 43,176 | 0.661 |
| 1988 | 3,712 | 0.438 | | | 816,772 | 0.312 | 381,218 | 0.331 |
| 1989 | 18,819 | 0.833 | | | 1,189,613 | 0.217 | 681,841 | 0.322 |
| 1990 | 48,228 | 0.827 | | | 2,222,315 | 0.340 | 1,273,208 | 0.331 |
| 1991 | 193,673 | 0.717 | | | 5,213,406 | 0.198 | 7,835,551 | 0.237 |
| 1992 | 59,066 | 0.358 | | | 4,896,293 | 0.110 | 3,915,744 | 0.284 |
| 1993 | 38,048 | 0.429 | | | 5,226,032 | 0.129 | 4,091,897 | 0.139 |
| 1994 | 56,953 | 0.376 | | | 3,996,091 | 0.129 | 3,380,181 | 0.159 |
| 1995 | 185,861 | 0.668 | | | 3,853,972 | 0.149 | 3,466,924 | 0.139 |
| 1996 | 27,416 | 0.472 | | | 4,119,728 | 0.179 | 4,601,138 | 0.275 |
| 1997 | 58,300 | 0.367 | | | 4,269,255 | 0.110 | 3,543,360 | 0.179 |
| 1998 | 146,962 | 0.303 | | | 5,878,807 | 0.120 | 3,455,676 | 0.159 |
| 1999 | 101,085 | 0.265 | | | 4,666,779 | 0.110 | 3,383,446 | 0.188 |
| 2000 | 117,022 | 0.421 | | | 4,815,223 | 0.129 | 3,500,946 | 0.217 |
| 2001 | 79,447 | 0.429 | | | 3,576,123 | 0.100 | 2,698,404 | 0.331 |

Table 7 Continued. Gulf of Mexico Gray Snapper recreational discards in numbers with associated log-scale standard errors (SE) input into the assessment model. Discards refer to the total number of fish discarded before applying the discard mortality rate.

| Year | Charter | Charter SE | Headboat | Headboat SE | Private | Private SE | Shore | Shore SE |
|------|---------|---------------|----------|----------------|------------|---------------|------------|-------------|
| 2002 | 94,140 | 0.217 | | | 4,307,791 | 0.090 | 2,298,504 | 0.188 |
| 2003 | 120,499 | 0.198 | | | 5,406,503 | 0.080 | 5,963,895 | 0.169 |
| 2004 | 92,490 | 0.208 | 5,452 | 0.198 | 4,735,098 | 0.129 | 4,780,642 | 0.198 |
| 2005 | 97,915 | 0.179 | 3,484 | 0.198 | 5,118,421 | 0.159 | 5,915,638 | 0.198 |
| 2006 | 91,967 | 0.275 | 5,413 | 0.198 | 4,339,792 | 0.159 | 2,957,106 | 0.237 |
| 2007 | 114,311 | 0.217 | 5,371 | 0.198 | 4,176,029 | 0.120 | 5,025,425 | 0.179 |
| 2008 | 170,558 | 0.169 | 4,710 | 0.198 | 7,288,358 | 0.110 | 7,291,040 | 0.217 |
| 2009 | 113,076 | 0.227 | 5,666 | 0.198 | 4,547,172 | 0.100 | 2,319,591 | 0.208 |
| 2010 | 136,030 | 0.198 | 5,849 | 0.198 | 3,371,951 | 0.169 | 1,621,169 | 0.217 |
| 2011 | 145,173 | 0.217 | 10,634 | 0.198 | 3,755,022 | 0.188 | 3,249,291 | 0.256 |
| 2012 | 144,093 | 0.159 | 8,636 | 0.198 | 6,307,317 | 0.149 | 4,346,403 | 0.188 |
| 2013 | 280,092 | 0.294 | 7,893 | 0.198 | 8,823,140 | 0.169 | 6,582,108 | 0.188 |
| 2014 | 258,945 | 0.188 | 13,017 | 0.198 | 11,593,216 | 0.110 | 5,884,124 | 0.208 |
| 2015 | 256,387 | 0.198 | 9,674 | 0.198 | 9,381,165 | 0.110 | 6,500,374 | 0.246 |
| 2016 | 254,007 | 0.208 | 7,883 | 0.198 | 6,361,109 | 0.090 | 6,992,583 | 0.169 |
| 2017 | 241,439 | 0.179 | 8,864 | 0.198 | 7,446,340 | 0.110 | 7,981,252 | 0.198 |
| 2018 | 356,050 | 0.188 | 10,795 | 0.198 | 5,970,615 | 0.120 | 7,807,543 | 0.227 |
| 2019 | 305,698 | 0.275 | 10,931 | 0.198 | 5,987,846 | 0.120 | 6,876,598 | 0.188 |
| 2020 | 487,968 | 0.294 | 11,016 | 0.198 | 10,440,887 | 0.129 | 12,175,650 | 0.169 |

Table 8. Standardized indices of relative abundance for Gulf of Mexico Gray Snapper. PriIndex = Private Index, ShrIndex = Shore Index, Trawl = SEAMAP Trawl Survey, CombVid = Combined Video Survey, RF = Reef fish visual survey.

| Year | CPUE PriInd | CPUE ShrInd | CPUE Age0 | CPUE Age1 | CPUE Trawl | CPUE CombVid Surv | CPUE RF Surv |
|------|----------------|----------------|--------------|--------------|---------------|-------------------------|-----------------|
| 1981 | 1.064 | 0.403 | | | | | |
| 1982 | 0.583 | 0.423 | | | | | |
| 1983 | 0.533 | 0.380 | | | | | |
| 1984 | 0.631 | 0.471 | | | | | |
| 1985 | 1.346 | 0.480 | | | | | |
| 1986 | 0.830 | 0.501 | | | | | |
| 1987 | 0.916 | 0.186 | | | | | |
| 1988 | 0.511 | 0.312 | | | | | |
| 1989 | 1.304 | 1.105 | | | | | |
| 1990 | 1.001 | 0.468 | | | | | |
| 1991 | 1.008 | 1.281 | | | | | |
| 1992 | 1.205 | 0.788 | | | | | |
| 1993 | 1.215 | 0.992 | | | | 0.720 | |
| 1994 | 1.180 | 0.760 | | | | 1.063 | |
| 1995 | 1.092 | 0.794 | | | | 0.695 | |
| 1996 | 0.830 | 1.082 | | 0.149 | | 0.645 | |
| 1997 | 1.034 | 1.063 | | 0.293 | | 1.029 | 0.371 |
| 1998 | 0.989 | 0.943 | 0.168 | 0.340 | | | 0.738 |
| 1999 | 0.762 | 0.701 | 0.927 | 0.280 | | | 0.755 |
| 2000 | 0.891 | 0.856 | 0.660 | 0.248 | | | 0.815 |
| 2001 | 0.892 | 0.586 | 1.343 | 0.226 | | | 0.958 |
| 2002 | 0.885 | 0.909 | 1.622 | 0.810 | | 0.863 | 1.053 |
| 2003 | 1.054 | 1.569 | 1.114 | 0.701 | | | 0.770 |

Table 8 Continued. Standardized indices of relative abundance for Gulf of Mexico Gray Snapper. PriIndex = Private Index, ShrIndex = Shore Index, Trawl = SEAMAP Trawl Survey, CombVid = Combined Video Survey, RF = Reef fish visual survey.

| Year | CPUE PriInd | CPUE ShrInd | CPUE Age0 | CPUE Age1 | CPUE Trawl | CPUE CombVid Surv | CPUE RF Surv |
|------|----------------|----------------|--------------|--------------|---------------|-------------------------|-----------------|
| 2004 | 0.679 | 1.094 | 1.725 | 0.716 | | 0.317 | 1.017 |
| 2005 | 0.826 | 1.670 | 0.446 | 0.535 | | 0.230 | 1.069 |
| 2006 | 0.990 | 1.072 | 1.302 | 0.221 | | 0.925 | 0.592 |
| 2007 | 1.107 | 1.864 | 0.970 | 1.897 | | 1.148 | 0.854 |
| 2008 | 1.026 | 1.229 | 0.470 | 0.955 | | 1.872 | 0.907 |
| 2009 | 0.919 | 0.790 | 1.320 | 0.747 | | 1.433 | 1.196 |
| 2010 | 0.572 | 0.586 | 0.867 | 0.274 | 1.101 | 0.744 | 0.935 |
| 2011 | 0.682 | 0.607 | 0.531 | 0.172 | 0.818 | 0.659 | 2.066 |
| 2012 | 0.787 | 0.984 | 1.395 | 0.512 | 0.822 | 0.880 | 1.167 |
| 2013 | 1.262 | 1.233 | 0.359 | 1.460 | 0.825 | 0.835 | |
| 2014 | 1.500 | 1.467 | 1.410 | 0.851 | 1.160 | 1.050 | 0.953 |
| 2015 | 1.330 | 1.323 | 0.615 | 2.117 | 0.955 | 1.124 | |
| 2016 | 1.220 | 1.454 | 1.639 | 2.220 | 1.099 | 1.096 | 0.910 |
| 2017 | 1.151 | 1.426 | 0.876 | 3.366 | 0.950 | 1.373 | |
| 2018 | 1.334 | 1.946 | 0.968 | 2.320 | 0.655 | 0.753 | 1.389 |
| 2019 | 1.257 | 1.839 | 1.046 | 1.948 | 1.616 | 1.204 | |
| 2020 | 1.601 | 2.363 | 1.228 | 1.642 | | 2.342 | |

Table 9. Log scale standard error associated with each standardized relative abundance index for Gulf of Mexico Gray Snapper. PriIndex = Private Index, ShrIndex = Shore Index, Trawl = SEAMAP Trawl Survey, CombVid = Combined Video Survey, RF = Reef fish visual survey.

| Year | SE PriInd | SE ShrInd | SE Age0 | SE Age1 | SE Trawl | SE CombVid Surv | SE RF Surv |
|------|-----------|-----------|---------|---------|----------|-----------------------|---------------|
| 1981 | 0.368 | 0.325 | | | | | |
| 1982 | 0.566 | 0.291 | | | | | |
| 1983 | 0.715 | 0.351 | | | | | |
| 1984 | 0.569 | 0.332 | | | | | |
| 1985 | 0.453 | 0.343 | | | | | |
| 1986 | 0.286 | 0.475 | | | | | |
| 1987 | 0.281 | 0.525 | | | | | |
| 1988 | 0.537 | 0.570 | | | | | |
| 1989 | 0.245 | 0.250 | | | | | |
| 1990 | 0.272 | 0.292 | | | | | |
| 1991 | 0.262 | 0.214 | | | | | |
| 1992 | 0.142 | 0.178 | | | | | |
| 1993 | 0.128 | 0.117 | | | | 0.315 | |
| 1994 | 0.127 | 0.128 | | | | 0.331 | |
| 1995 | 0.150 | 0.131 | | | | 0.369 | |
| 1996 | 0.137 | 0.145 | | 0.288 | | 0.264 | |
| 1997 | 0.129 | 0.140 | | 0.229 | | 0.272 | 0.137 |
| 1998 | 0.111 | 0.127 | 0.305 | 0.255 | | | 0.123 |
| 1999 | 0.110 | 0.132 | 0.233 | 0.230 | | | 0.106 |
| 2000 | 0.127 | 0.152 | 0.209 | 0.214 | | | 0.107 |
| 2001 | 0.123 | 0.163 | 0.188 | 0.249 | | | 0.118 |
| 2002 | 0.107 | 0.145 | 0.159 | 0.194 | | 0.280 | 0.121 |

Table 9 Continued. Log scale standard error associated with each standardized relative abundance index for Gulf of Mexico Gray Snapper. PriIndex = Private Index, ShrIndex = Shore Index, Trawl = SEAMAP Trawl Survey, CombVid = Combined Video Survey, RF = Reef fish visual survey.

| Year | SE PriInd | SE ShrInd | SE Age0 | SE Age1 | SE Trawl | SE CombVid Surv | SE RF Surv |
|------|-----------|-----------|---------|---------|----------|-----------------------|---------------|
| 2003 | 0.095 | 0.111 | 0.156 | 0.219 | | | 0.151 |
| 2004 | 0.127 | 0.142 | 0.159 | 0.211 | | 0.358 | 0.122 |
| 2005 | 0.116 | 0.122 | 0.200 | 0.237 | | 0.374 | 0.140 |
| 2006 | 0.120 | 0.172 | 0.147 | 0.239 | | 0.203 | 0.122 |
| 2007 | 0.111 | 0.114 | 0.165 | 0.191 | | 0.287 | 0.139 |
| 2008 | 0.103 | 0.130 | 0.190 | 0.201 | | 0.229 | 0.110 |
| 2009 | 0.106 | 0.138 | 0.150 | 0.203 | | 0.185 | 0.112 |
| 2010 | 0.168 | 0.167 | 0.163 | 0.234 | 0.227 | 0.185 | 0.098 |
| 2011 | 0.172 | 0.186 | 0.181 | 0.276 | 0.266 | 0.129 | 0.118 |
| 2012 | 0.145 | 0.146 | 0.173 | 0.182 | 0.228 | 0.138 | 0.095 |
| 2013 | 0.116 | 0.123 | 0.301 | 0.187 | 0.263 | 0.170 | |
| 2014 | 0.081 | 0.136 | 0.161 | 0.192 | 0.210 | 0.125 | 0.104 |
| 2015 | 0.087 | 0.147 | 0.190 | 0.180 | 0.249 | 0.134 | |
| 2016 | 0.090 | 0.127 | 0.149 | 0.188 | 0.218 | 0.119 | 0.102 |
| 2017 | 0.108 | 0.133 | 0.169 | 0.185 | 0.254 | 0.165 | |
| 2018 | 0.109 | 0.131 | 0.168 | 0.197 | 0.238 | 0.167 | 0.092 |
| 2019 | 0.104 | 0.139 | 0.164 | 0.182 | 0.244 | 0.167 | |
| 2020 | 0.095 | 0.111 | 0.167 | 0.197 | | 0.156 | |

Table 10. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|----------------------|----------|----------|-------|--------|-------|-------|
| L_at_Amin_Fem_GP_1 | 16.71 | (0,40) | 0.135 | 0.008 | | 3 |
| L_at_Amax_Fem_GP_1 | 60.8 | | | | | Fixed |
| VonBert_K_Fem_GP_1 | 0.113 | | | | | Fixed |
| CV_young_Fem_GP_1 | 0.151 | | | | | Fixed |
| CV_old_Fem_GP_1 | 0.192 | | | | | Fixed |
| Wtlen_1_Fem_GP_1 | 1.43e-05 | | | | | Fixed |
| Wtlen_2_Fem_GP_1 | 3.02 | | | | | Fixed |
| Mat50%_Fem_GP_1 | 27 | | | | | Fixed |
| Mat_slope_Fem_GP_1 | 0.02 | | | | | Fixed |
| Eggs_scalar_Fem_GP_1 | 1 | | | | | Fixed |
| Eggs_exp_wt_Fem_GP_1 | 1 | | | | | Fixed |
| CohortGrowDev | 1 | | | | | Fixed |
| FracFemale_GP_1 | 0.5 | | | | | Fixed |
| SR_LN(R0) | 10.05 | (5,15) | 0.046 | 0.005 | | 1 |
| SR_BH_steep | 0.99 | | | | | Fixed |
| SR_sigmaR | 0.376 | (0.01,2) | 0.046 | 0.123 | | 4 |
| SR_regime | 0.00e+00 | | | | | Fixed |
| SR_autocorr | 0.00e+00 | | | | | Fixed |
| Early_RecrDev_1951 | 0.015 | (-5,5) | 0.379 | 25.71 | | 6 |
| Early_RecrDev_1952 | 0.016 | (-5,5) | 0.379 | 23.23 | | 6 |
| Early_RecrDev_1953 | 0.018 | (-5,5) | 0.379 | 20.82 | | 6 |
| Early_RecrDev_1954 | 0.021 | (-5,5) | 0.38 | 18.35 | | 6 |
| Early_RecrDev_1955 | 0.022 | (-5,5) | 0.38 | 17.51 | | 6 |
| Early_RecrDev_1956 | 0.015 | (-5,5) | 0.379 | 25.1 | | 6 |
| Early_RecrDev_1957 | 0.009 | (-5,5) | 0.379 | 42.1 | | 6 |
| Early_RecrDev_1958 | 0.027 | (-5,5) | 0.383 | 14.31 | | 6 |
| Early_RecrDev_1959 | 0.013 | (-5,5) | 0.385 | 30.2 | | 6 |
| Early_RecrDev_1960 | -0.035 | (-5,5) | 0.377 | -10.8 | | 6 |
| Early_RecrDev_1961 | -0.021 | (-5,5) | 0.368 | -17.42 | | 6 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|--------------------|--------|--------|-------|--------|-------|-------|
| Early_RecrDev_1962 | 0.08 | (-5,5) | 0.359 | 4.52 | | 6 |
| Early_RecrDev_1963 | 0.045 | (-5,5) | 0.352 | 7.88 | | 6 |
| Early_RecrDev_1964 | -0.132 | (-5,5) | 0.343 | -2.59 | | 6 |
| Early_RecrDev_1965 | -0.16 | (-5,5) | 0.334 | -2.08 | | 6 |
| Early_RecrDev_1966 | -0.185 | (-5,5) | 0.328 | -1.77 | | 6 |
| Early_RecrDev_1967 | -0.178 | (-5,5) | 0.32 | -1.8 | | 6 |
| Early_RecrDev_1968 | -0.284 | (-5,5) | 0.317 | -1.12 | | 6 |
| Early_RecrDev_1969 | -0.309 | (-5,5) | 0.313 | -1.02 | | 6 |
| Early_RecrDev_1970 | -0.266 | (-5,5) | 0.31 | -1.17 | | 6 |
| Early_RecrDev_1971 | -0.163 | (-5,5) | 0.301 | -1.84 | | 6 |
| Early_RecrDev_1972 | -0.077 | (-5,5) | 0.291 | -3.8 | | 6 |
| Early_RecrDev_1973 | -0.066 | (-5,5) | 0.281 | -4.28 | | 6 |
| Early_RecrDev_1974 | -0.051 | (-5,5) | 0.267 | -5.24 | | 6 |
| Early_RecrDev_1975 | -0.173 | (-5,5) | 0.267 | -1.54 | | 6 |
| Early_RecrDev_1976 | -0.226 | (-5,5) | 0.258 | -1.14 | | 6 |
| Early_RecrDev_1977 | -0.33 | (-5,5) | 0.246 | -0.746 | | 6 |
| Early_RecrDev_1978 | -0.398 | (-5,5) | 0.234 | -0.588 | | 6 |
| Early_RecrDev_1979 | -0.53 | (-5,5) | 0.231 | -0.436 | | 6 |
| Early_RecrDev_1980 | -0.4 | (-5,5) | 0.188 | -0.469 | | 6 |
| Main_RecrDev_1981 | 0.01 | (-5,5) | 0.153 | 14.93 | | 3 |
| Main_RecrDev_1982 | -0.233 | (-5,5) | 0.204 | -0.877 | | 3 |
| Main_RecrDev_1983 | -0.118 | (-5,5) | 0.189 | -1.61 | | 3 |
| Main_RecrDev_1984 | -0.143 | (-5,5) | 0.184 | -1.29 | | 3 |
| Main_RecrDev_1985 | 0.037 | (-5,5) | 0.145 | 3.88 | | 3 |
| Main_RecrDev_1986 | -0.538 | (-5,5) | 0.17 | -0.317 | | 3 |
| Main_RecrDev_1987 | -0.329 | (-5,5) | 0.149 | -0.454 | | 3 |
| Main_RecrDev_1988 | 0.293 | (-5,5) | 0.098 | 0.335 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-------------------|--------|--------|-------|--------|-------|-------|
| Main_RecrDev_1989 | -0.786 | (-5,5) | 0.153 | -0.195 | | 3 |
| Main_RecrDev_1990 | 0.158 | (-5,5) | 0.084 | 0.532 | | 3 |
| Main_RecrDev_1991 | 0.184 | (-5,5) | 0.08 | 0.434 | | 3 |
| Main_RecrDev_1992 | -0.171 | (-5,5) | 0.093 | -0.543 | | 3 |
| Main_RecrDev_1993 | -0.529 | (-5,5) | 0.106 | -0.201 | | 3 |
| Main_RecrDev_1994 | -0.333 | (-5,5) | 0.088 | -0.265 | | 3 |
| Main_RecrDev_1995 | -0.434 | (-5,5) | 0.088 | -0.203 | | 3 |
| Main_RecrDev_1996 | -0.017 | (-5,5) | 0.065 | -3.71 | | 3 |
| Main_RecrDev_1997 | 0.181 | (-5,5) | 0.056 | 0.309 | | 3 |
| Main_RecrDev_1998 | -0.39 | (-5,5) | 0.077 | -0.197 | | 3 |
| Main_RecrDev_1999 | -0.158 | (-5,5) | 0.062 | -0.392 | | 3 |
| Main_RecrDev_2000 | -0.621 | (-5,5) | 0.076 | -0.123 | | 3 |
| Main_RecrDev_2001 | 0.082 | (-5,5) | 0.051 | 0.627 | | 3 |
| Main_RecrDev_2002 | -0.066 | (-5,5) | 0.053 | -0.792 | | 3 |
| Main_RecrDev_2003 | 0.06 | (-5,5) | 0.048 | 0.795 | | 3 |
| Main_RecrDev_2004 | 0.039 | (-5,5) | 0.048 | 1.22 | | 3 |
| Main_RecrDev_2005 | 0.09 | (-5,5) | 0.05 | 0.553 | | 3 |
| Main_RecrDev_2006 | 0.356 | (-5,5) | 0.043 | 0.12 | | 3 |
| Main_RecrDev_2007 | -0.164 | (-5,5) | 0.057 | -0.347 | | 3 |
| Main_RecrDev_2008 | -0.248 | (-5,5) | 0.063 | -0.254 | | 3 |
| Main_RecrDev_2009 | -0.166 | (-5,5) | 0.059 | -0.358 | | 3 |
| Main_RecrDev_2010 | 0.087 | (-5,5) | 0.058 | 0.667 | | 3 |
| Main_RecrDev_2011 | 0.469 | (-5,5) | 0.052 | 0.111 | | 3 |
| Main_RecrDev_2012 | 0.513 | (-5,5) | 0.053 | 0.103 | | 3 |
| Main_RecrDev_2013 | -0.317 | (-5,5) | 0.09 | -0.283 | | 3 |
| Main_RecrDev_2014 | 0.414 | (-5,5) | 0.059 | 0.142 | | 3 |
| Main_RecrDev_2015 | 0.071 | (-5,5) | 0.075 | 1.07 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|----------|---------|-------|-------|-------|-------|
| Main_RecrDev_2016 | 0.476 | (-5,5) | 0.067 | 0.142 | | 3 |
| Main_RecrDev_2017 | 0.412 | (-5,5) | 0.084 | 0.205 | | 3 |
| Main_RecrDev_2018 | 0.758 | (-5,5) | 0.097 | 0.127 | | 3 |
| Main_RecrDev_2019 | 0.71 | (-5,5) | 0.118 | 0.166 | | 3 |
| Main_RecrDev_2020 | 0.362 | (-5,5) | 0.143 | 0.396 | | 3 |
| ForeRecr_2021 | 0.00e+00 | | | | | Fixed |
| F_fleet_1_YR_1945_s_1 | 0.031 | (0,2.9) | 0.009 | 0.281 | | 3 |
| F_fleet_1_YR_1946_s_1 | 0.032 | (0,2.9) | 0.009 | 0.282 | | 3 |
| F_fleet_1_YR_1947_s_1 | 0.033 | (0,2.9) | 0.009 | 0.282 | | 3 |
| F_fleet_1_YR_1948_s_1 | 0.034 | (0,2.9) | 0.009 | 0.282 | | 3 |
| F_fleet_1_YR_1949_s_1 | 0.035 | (0,2.9) | 0.01 | 0.282 | | 3 |
| F_fleet_1_YR_1950_s_1 | 0.036 | (0,2.9) | 0.01 | 0.282 | | 3 |
| F_fleet_1_YR_1951_s_1 | 0.037 | (0,2.9) | 0.01 | 0.282 | | 3 |
| F_fleet_1_YR_1952_s_1 | 0.038 | (0,2.9) | 0.011 | 0.283 | | 3 |
| F_fleet_1_YR_1953_s_1 | 0.039 | (0,2.9) | 0.011 | 0.283 | | 3 |
| F_fleet_1_YR_1954_s_1 | 0.04 | (0,2.9) | 0.011 | 0.283 | | 3 |
| F_fleet_1_YR_1955_s_1 | 0.041 | (0,2.9) | 0.012 | 0.283 | | 3 |
| F_fleet_1_YR_1956_s_1 | 0.042 | (0,2.9) | 0.012 | 0.284 | | 3 |
| F_fleet_1_YR_1957_s_1 | 0.044 | (0,2.9) | 0.012 | 0.285 | | 3 |
| F_fleet_1_YR_1958_s_1 | 0.045 | (0,2.9) | 0.013 | 0.285 | | 3 |
| F_fleet_1_YR_1959_s_1 | 0.046 | (0,2.9) | 0.013 | 0.286 | | 3 |
| F_fleet_1_YR_1960_s_1 | 0.048 | (0,2.9) | 0.014 | 0.288 | | 3 |
| F_fleet_1_YR_1961_s_1 | 0.049 | (0,2.9) | 0.014 | 0.289 | | 3 |
| F_fleet_1_YR_1962_s_1 | 0.054 | (0,2.9) | 0.016 | 0.29 | | 3 |
| F_fleet_1_YR_1963_s_1 | 0.047 | (0,2.9) | 0.014 | 0.292 | | 3 |
| F_fleet_1_YR_1964_s_1 | 0.051 | (0,2.9) | 0.015 | 0.294 | | 3 |
| F_fleet_1_YR_1965_s_1 | 0.057 | (0,2.9) | 0.017 | 0.296 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|-------|-------|-------|-------|
| F_fleet_1_YR_1966_s_1 | 0.049 | (0,2.9) | 0.015 | 0.297 | | 3 |
| F_fleet_1_YR_1967_s_1 | 0.057 | (0,2.9) | 0.017 | 0.299 | | 3 |
| F_fleet_1_YR_1968_s_1 | 0.073 | (0,2.9) | 0.022 | 0.301 | | 3 |
| F_fleet_1_YR_1969_s_1 | 0.07 | (0,2.9) | 0.021 | 0.302 | | 3 |
| F_fleet_1_YR_1970_s_1 | 0.065 | (0,2.9) | 0.02 | 0.303 | | 3 |
| F_fleet_1_YR_1971_s_1 | 0.071 | (0,2.9) | 0.022 | 0.304 | | 3 |
| F_fleet_1_YR_1972_s_1 | 0.082 | (0,2.9) | 0.025 | 0.305 | | 3 |
| F_fleet_1_YR_1973_s_1 | 0.082 | (0,2.9) | 0.025 | 0.305 | | 3 |
| F_fleet_1_YR_1974_s_1 | 0.087 | (0,2.9) | 0.026 | 0.305 | | 3 |
| F_fleet_1_YR_1975_s_1 | 0.059 | (0,2.9) | 0.018 | 0.304 | | 3 |
| F_fleet_1_YR_1976_s_1 | 0.113 | (0,2.9) | 0.034 | 0.303 | | 3 |
| F_fleet_1_YR_1977_s_1 | 0.067 | (0,2.9) | 0.02 | 0.301 | | 3 |
| F_fleet_1_YR_1978_s_1 | 0.06 | (0,2.9) | 0.018 | 0.3 | | 3 |
| F_fleet_1_YR_1979_s_1 | 0.075 | (0,2.9) | 0.022 | 0.299 | | 3 |
| F_fleet_1_YR_1980_s_1 | 0.146 | (0,2.9) | 0.043 | 0.298 | | 3 |
| F_fleet_1_YR_1981_s_1 | 0.157 | (0,2.9) | 0.047 | 0.298 | | 3 |
| F_fleet_1_YR_1982_s_1 | 0.231 | (0,2.9) | 0.069 | 0.299 | | 3 |
| F_fleet_1_YR_1983_s_1 | 0.249 | (0,2.9) | 0.075 | 0.301 | | 3 |
| F_fleet_1_YR_1984_s_1 | 0.186 | (0,2.9) | 0.056 | 0.302 | | 3 |
| F_fleet_1_YR_1985_s_1 | 0.154 | (0,2.9) | 0.046 | 0.301 | | 3 |
| F_fleet_1_YR_1986_s_1 | 0.171 | (0,2.9) | 0.051 | 0.3 | | 3 |
| F_fleet_1_YR_1987_s_1 | 0.212 | (0,2.9) | 0.063 | 0.299 | | 3 |
| F_fleet_1_YR_1988_s_1 | 0.138 | (0,2.9) | 0.041 | 0.298 | | 3 |
| F_fleet_1_YR_1989_s_1 | 0.17 | (0,2.9) | 0.051 | 0.298 | | 3 |
| F_fleet_1_YR_1990_s_1 | 0.145 | (0,2.9) | 0.043 | 0.299 | | 3 |
| F_fleet_1_YR_1991_s_1 | 0.195 | (0,2.9) | 0.058 | 0.3 | | 3 |
| F_fleet_1_YR_1992_s_1 | 0.197 | (0,2.9) | 0.059 | 0.302 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|-------|-------|-------|-------|
| F_fleet_1_YR_1993_s_1 | 0.249 | (0,2.9) | 0.076 | 0.304 | | 3 |
| F_fleet_1_YR_1994_s_1 | 0.312 | (0,2.9) | 0.095 | 0.305 | | 3 |
| F_fleet_1_YR_1995_s_1 | 0.255 | (0,2.9) | 0.078 | 0.306 | | 3 |
| F_fleet_1_YR_1996_s_1 | 0.249 | (0,2.9) | 0.076 | 0.307 | | 3 |
| F_fleet_1_YR_1997_s_1 | 0.245 | (0,2.9) | 0.075 | 0.308 | | 3 |
| F_fleet_1_YR_1998_s_1 | 0.197 | (0,2.9) | 0.061 | 0.309 | | 3 |
| F_fleet_1_YR_1999_s_1 | 0.181 | (0,2.9) | 0.056 | 0.31 | | 3 |
| F_fleet_1_YR_2000_s_1 | 0.188 | (0,2.9) | 0.058 | 0.31 | | 3 |
| F_fleet_1_YR_2001_s_1 | 0.197 | (0,2.9) | 0.061 | 0.31 | | 3 |
| F_fleet_1_YR_2002_s_1 | 0.231 | (0,2.9) | 0.072 | 0.311 | | 3 |
| F_fleet_1_YR_2003_s_1 | 0.202 | (0,2.9) | 0.063 | 0.312 | | 3 |
| F_fleet_1_YR_2004_s_1 | 0.215 | (0,2.9) | 0.067 | 0.313 | | 3 |
| F_fleet_1_YR_2005_s_1 | 0.193 | (0,2.9) | 0.061 | 0.314 | | 3 |
| F_fleet_1_YR_2006_s_1 | 0.167 | (0,2.9) | 0.052 | 0.314 | | 3 |
| F_fleet_1_YR_2007_s_1 | 0.122 | (0,2.9) | 0.038 | 0.314 | | 3 |
| F_fleet_1_YR_2008_s_1 | 0.123 | (0,2.9) | 0.039 | 0.313 | | 3 |
| F_fleet_1_YR_2009_s_1 | 0.149 | (0,2.9) | 0.047 | 0.314 | | 3 |
| F_fleet_1_YR_2010_s_1 | 0.117 | (0,2.9) | 0.037 | 0.314 | | 3 |
| F_fleet_1_YR_2011_s_1 | 0.125 | (0,2.9) | 0.039 | 0.314 | | 3 |
| F_fleet_1_YR_2012_s_1 | 0.129 | (0,2.9) | 0.041 | 0.315 | | 3 |
| F_fleet_1_YR_2013_s_1 | 0.117 | (0,2.9) | 0.037 | 0.316 | | 3 |
| F_fleet_1_YR_2014_s_1 | 0.146 | (0,2.9) | 0.046 | 0.317 | | 3 |
| F_fleet_1_YR_2015_s_1 | 0.126 | (0,2.9) | 0.04 | 0.317 | | 3 |
| F_fleet_1_YR_2016_s_1 | 0.126 | (0,2.9) | 0.04 | 0.317 | | 3 |
| F_fleet_1_YR_2017_s_1 | 0.095 | (0,2.9) | 0.03 | 0.318 | | 3 |
| F_fleet_1_YR_2018_s_1 | 0.099 | (0,2.9) | 0.032 | 0.319 | | 3 |
| F_fleet_1_YR_2019_s_1 | 0.08 | (0,2.9) | 0.026 | 0.319 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|----------|---------|----------------------|-------|-------|-------|
| F_fleet_1_YR_2020_s_1 | 0.062 | (0,2.9) | 0.02 | 0.319 | | 3 |
| F_fleet_2_YR_1963_s_1 | 2.72e-05 | (0,2.9) | 4.05e- ^{^4} | 0.149 | | 3 |
| F_fleet_2_YR_1980_s_1 | 0.002 | (0,2.9) | 3.33e- ^{^4} | 0.184 | | 3 |
| F_fleet_2_YR_1981_s_1 | 0.002 | (0,2.9) | 3.67e- ^{^4} | 0.184 | | 3 |
| F_fleet_2_YR_1982_s_1 | 0.005 | (0,2.9) | 9.51e- ^{^4} | 0.186 | | 3 |
| F_fleet_2_YR_1983_s_1 | 0.01 | (0,2.9) | 0.002 | 0.189 | | 3 |
| F_fleet_2_YR_1984_s_1 | 0.006 | (0,2.9) | 0.001 | 0.19 | | 3 |
| F_fleet_2_YR_1985_s_1 | 0.005 | (0,2.9) | 8.56e- ^{^4} | 0.189 | | 3 |
| F_fleet_2_YR_1986_s_1 | 0.006 | (0,2.9) | 0.001 | 0.187 | | 3 |
| F_fleet_2_YR_1987_s_1 | 0.006 | (0,2.9) | 0.001 | 0.186 | | 3 |
| F_fleet_2_YR_1988_s_1 | 0.005 | (0,2.9) | 8.34e- ^{^4} | 0.183 | | 3 |
| F_fleet_2_YR_1989_s_1 | 0.007 | (0,2.9) | 0.001 | 0.181 | | 3 |
| F_fleet_2_YR_1990_s_1 | 0.007 | (0,2.9) | 0.001 | 0.18 | | 3 |
| F_fleet_2_YR_1991_s_1 | 0.007 | (0,2.9) | 0.001 | 0.181 | | 3 |
| F_fleet_2_YR_1992_s_1 | 0.009 | (0,2.9) | 0.002 | 0.182 | | 3 |
| F_fleet_2_YR_1993_s_1 | 0.013 | (0,2.9) | 0.002 | 0.181 | | 3 |
| F_fleet_2_YR_1994_s_1 | 0.002 | (0,2.9) | 3.30e- ^{^4} | 0.184 | | 3 |
| F_fleet_2_YR_1995_s_1 | 0.002 | (0,2.9) | 3.61e- ^{^4} | 0.185 | | 3 |
| F_fleet_2_YR_1996_s_1 | 0.002 | (0,2.9) | 2.79e- ^{^4} | 0.186 | | 3 |
| F_fleet_2_YR_1997_s_1 | 0.002 | (0,2.9) | 3.15e- ^{^4} | 0.186 | | 3 |
| F_fleet_2_YR_1998_s_1 | 0.002 | (0,2.9) | 3.64e- ^{^4} | 0.186 | | 3 |
| F_fleet_2_YR_1999_s_1 | 0.003 | (0,2.9) | 5.76e- ^{^4} | 0.187 | | 3 |
| F_fleet_2_YR_2000_s_1 | 0.003 | (0,2.9) | 5.65e- ^{^4} | 0.187 | | 3 |
| F_fleet_2_YR_2001_s_1 | 0.003 | (0,2.9) | 4.93e- ^{^4} | 0.187 | | 3 |
| F_fleet_2_YR_2002_s_1 | 0.003 | (0,2.9) | 6.19e- ^{^4} | 0.188 | | 3 |
| F_fleet_2_YR_2003_s_1 | 0.002 | (0,2.9) | 4.47e- ^{^4} | 0.189 | | 3 |
| F_fleet_2_YR_2004_s_1 | 0.004 | (0,2.9) | 6.93e- ^{^4} | 0.191 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|----------|---------|--------|-------|-------|-------|
| F_fleet_2_YR_2005_s_1 | 0.003 | (0,2.9) | 6.07e- | 0.193 | | 3 |
| F_fleet_2_YR_2006_s_1 | 0.003 | (0,2.9) | 5.90e- | 0.193 | | 3 |
| F_fleet_2_YR_2007_s_1 | 0.003 | (0,2.9) | 5.29e- | 0.192 | | 3 |
| F_fleet_2_YR_2008_s_1 | 0.003 | (0,2.9) | 6.45e- | 0.192 | | 3 |
| F_fleet_2_YR_2009_s_1 | 0.004 | (0,2.9) | 7.05e- | 0.192 | | 3 |
| F_fleet_2_YR_2010_s_1 | 0.001 | (0,2.9) | 2.86e- | 0.194 | | 3 |
| F_fleet_2_YR_2011_s_1 | 0.003 | (0,2.9) | 4.95e- | 0.192 | | 3 |
| F_fleet_2_YR_2012_s_1 | 0.003 | (0,2.9) | 5.15e- | 0.193 | | 3 |
| F_fleet_2_YR_2013_s_1 | 0.002 | (0,2.9) | 4.82e- | 0.195 | | 3 |
| F_fleet_2_YR_2014_s_1 | 0.004 | (0,2.9) | 7.04e- | 0.197 | | 3 |
| F_fleet_2_YR_2015_s_1 | 0.005 | (0,2.9) | 0.001 | 0.198 | | 3 |
| F_fleet_2_YR_2016_s_1 | 0.005 | (0,2.9) | 9.88e- | 0.2 | | 3 |
| F_fleet_2_YR_2017_s_1 | 0.004 | (0,2.9) | 8.18e- | 0.203 | | 3 |
| F_fleet_2_YR_2018_s_1 | 0.002 | (0,2.9) | 4.66e- | 0.204 | | 3 |
| F_fleet_2_YR_2019_s_1 | 0.003 | (0,2.9) | 5.18e- | 0.204 | | 3 |
| F_fleet_2_YR_2020_s_1 | 0.002 | (0,2.9) | 3.97e- | 0.205 | | 3 |
| F_fleet_3_YR_1945_s_1 | 4.61e-04 | (0,2.9) | 3.86e- | 0.084 | | 3 |
| F_fleet_3_YR_1946_s_1 | 4.72e-04 | (0,2.9) | 3.96e- | 0.084 | | 3 |
| F_fleet_3_YR_1947_s_1 | 4.84e-04 | (0,2.9) | 4.06e- | 0.084 | | 3 |
| F_fleet_3_YR_1948_s_1 | 4.97e-04 | (0,2.9) | 4.17e- | 0.084 | | 3 |
| F_fleet_3_YR_1949_s_1 | 5.11e-04 | (0,2.9) | 4.29e- | 0.084 | | 3 |
| F_fleet_3_YR_1950_s_1 | 5.25e-04 | (0,2.9) | 4.42e- | 0.084 | | 3 |
| F_fleet_3_YR_1951_s_1 | 5.41e-04 | (0,2.9) | 4.56e- | 0.084 | | 3 |
| F_fleet_3_YR_1952_s_1 | 5.57e-04 | (0,2.9) | 4.71e- | 0.085 | | 3 |
| F_fleet_3_YR_1953_s_1 | 5.73e-04 | (0,2.9) | 4.88e- | 0.085 | | 3 |
| F_fleet_3_YR_1954_s_1 | 5.90e-04 | (0,2.9) | 5.28e- | 0.089 | | 3 |
| F_fleet_3_YR_1955_s_1 | 6.07e-04 | (0,2.9) | 6.11e- | 0.101 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|----------|---------|--------|-------|-------|-------|
| F_fleet_3_YR_1956_s_1 | 6.22e-04 | (0,2.9) | 7.09e- | 0.114 | | 3 |
| F_fleet_3_YR_1957_s_1 | 6.37e-04 | (0,2.9) | 7.95e- | 0.125 | | 3 |
| F_fleet_3_YR_1958_s_1 | 6.52e-04 | (0,2.9) | 8.63e- | 0.132 | | 3 |
| F_fleet_3_YR_1959_s_1 | 6.68e-04 | (0,2.9) | 9.18e- | 0.137 | | 3 |
| F_fleet_3_YR_1960_s_1 | 6.85e-04 | (0,2.9) | 9.64e- | 0.141 | | 3 |
| F_fleet_3_YR_1961_s_1 | 7.03e-04 | (0,2.9) | 1.01e- | 0.143 | | 3 |
| F_fleet_3_YR_1962_s_1 | 4.67e-04 | (0,2.9) | 6.76e- | 0.145 | | 3 |
| F_fleet_3_YR_1963_s_1 | 2.81e-04 | (0,2.9) | 4.10e- | 0.146 | | 3 |
| F_fleet_3_YR_1964_s_1 | 4.40e-04 | (0,2.9) | 6.42e- | 0.146 | | 3 |
| F_fleet_3_YR_1965_s_1 | 0.002 | (0,2.9) | 2.21e- | 0.145 | | 3 |
| F_fleet_3_YR_1966_s_1 | 0.001 | (0,2.9) | 1.45e- | 0.145 | | 3 |
| F_fleet_3_YR_1967_s_1 | 0.002 | (0,2.9) | 2.58e- | 0.145 | | 3 |
| F_fleet_3_YR_1968_s_1 | 0.002 | (0,2.9) | 3.52e- | 0.144 | | 3 |
| F_fleet_3_YR_1969_s_1 | 0.004 | (0,2.9) | 5.35e- | 0.141 | | 3 |
| F_fleet_3_YR_1970_s_1 | 0.004 | (0,2.9) | 5.34e- | 0.139 | | 3 |
| F_fleet_3_YR_1971_s_1 | 0.004 | (0,2.9) | 5.74e- | 0.137 | | 3 |
| F_fleet_3_YR_1972_s_1 | 0.005 | (0,2.9) | 6.97e- | 0.135 | | 3 |
| F_fleet_3_YR_1973_s_1 | 0.007 | (0,2.9) | 9.68e- | 0.134 | | 3 |
| F_fleet_3_YR_1974_s_1 | 0.009 | (0,2.9) | 0.001 | 0.132 | | 3 |
| F_fleet_3_YR_1975_s_1 | 0.01 | (0,2.9) | 0.001 | 0.13 | | 3 |
| F_fleet_3_YR_1976_s_1 | 0.006 | (0,2.9) | 7.44e- | 0.129 | | 3 |
| F_fleet_3_YR_1977_s_1 | 0.017 | (0,2.9) | 0.002 | 0.127 | | 3 |
| F_fleet_3_YR_1978_s_1 | 0.021 | (0,2.9) | 0.003 | 0.125 | | 3 |
| F_fleet_3_YR_1979_s_1 | 0.02 | (0,2.9) | 0.002 | 0.122 | | 3 |
| F_fleet_3_YR_1980_s_1 | 0.009 | (0,2.9) | 0.001 | 0.119 | | 3 |
| F_fleet_3_YR_1981_s_1 | 0.009 | (0,2.9) | 0.001 | 0.117 | | 3 |
| F_fleet_3_YR_1982_s_1 | 0.011 | (0,2.9) | 0.001 | 0.117 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|----------|---------|----------------------|-------|-------|-------|
| F_fleet_3_YR_1983_s_1 | 0.012 | (0,2.9) | 0.001 | 0.117 | | 3 |
| F_fleet_3_YR_1984_s_1 | 0.019 | (0,2.9) | 0.002 | 0.114 | | 3 |
| F_fleet_3_YR_1985_s_1 | 0.015 | (0,2.9) | 0.002 | 0.11 | | 3 |
| F_fleet_3_YR_1986_s_1 | 0.011 | (0,2.9) | 0.001 | 0.106 | | 3 |
| F_fleet_3_YR_1987_s_1 | 0.009 | (0,2.9) | 9.07e- ^{^4} | 0.104 | | 3 |
| F_fleet_3_YR_1988_s_1 | 0.006 | (0,2.9) | 6.14e- ^{^4} | 0.101 | | 3 |
| F_fleet_3_YR_1989_s_1 | 0.008 | (0,2.9) | 7.53e- ^{^4} | 0.1 | | 3 |
| F_fleet_3_YR_1990_s_1 | 0.003 | (0,2.9) | 2.91e- ^{^4} | 0.1 | | 3 |
| F_fleet_3_YR_1991_s_1 | 0.003 | (0,2.9) | 3.17e- ^{^4} | 0.1 | | 3 |
| F_fleet_3_YR_1992_s_1 | 0.002 | (0,2.9) | 1.77e- ^{^4} | 0.101 | | 3 |
| F_fleet_3_YR_1993_s_1 | 0.001 | (0,2.9) | 1.02e- ^{^4} | 0.099 | | 3 |
| F_fleet_3_YR_1994_s_1 | 0.003 | (0,2.9) | 2.53e- ^{^4} | 0.096 | | 3 |
| F_fleet_3_YR_1995_s_1 | 0.001 | (0,2.9) | 1.26e- ^{^4} | 0.095 | | 3 |
| F_fleet_3_YR_1996_s_1 | 6.73e-04 | (0,2.9) | 6.44e- ^{^5} | 0.096 | | 3 |
| F_fleet_3_YR_1997_s_1 | 7.74e-04 | (0,2.9) | 7.53e- ^{^5} | 0.097 | | 3 |
| F_fleet_3_YR_1998_s_1 | 8.58e-04 | (0,2.9) | 8.45e- ^{^5} | 0.099 | | 3 |
| F_fleet_3_YR_1999_s_1 | 6.12e-04 | (0,2.9) | 6.03e- ^{^5} | 0.099 | | 3 |
| F_fleet_3_YR_2000_s_1 | 4.08e-04 | (0,2.9) | 3.95e- ^{^5} | 0.097 | | 3 |
| F_fleet_3_YR_2001_s_1 | 3.39e-04 | (0,2.9) | 3.26e- ^{^5} | 0.096 | | 3 |
| F_fleet_3_YR_2002_s_1 | 2.65e-04 | (0,2.9) | 2.57e- ^{^5} | 0.097 | | 3 |
| F_fleet_3_YR_2003_s_1 | 3.44e-04 | (0,2.9) | 3.39e- ^{^5} | 0.098 | | 3 |
| F_fleet_3_YR_2004_s_1 | 1.39e-04 | (0,2.9) | 1.40e- ^{^5} | 0.1 | | 3 |
| F_fleet_3_YR_2005_s_1 | 1.11e-04 | (0,2.9) | 1.12e- ^{^5} | 0.102 | | 3 |
| F_fleet_3_YR_2006_s_1 | 5.00e-05 | (0,2.9) | 5.06e- ^{^6} | 0.101 | | 3 |
| F_fleet_3_YR_2007_s_1 | 2.69e-05 | (0,2.9) | 2.70e- ^{^6} | 0.1 | | 3 |
| F_fleet_3_YR_2008_s_1 | 5.89e-05 | (0,2.9) | 5.92e- ^{^6} | 0.101 | | 3 |
| F_fleet_3_YR_2009_s_1 | 5.17e-05 | (0,2.9) | 5.27e- ^{^6} | 0.102 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|----------|---------|--------|-------|-------|-------|
| F_fleet_3_YR_2010_s_1 | 5.04e-06 | (0,2.9) | 5.18e- | 0.103 | | 3 |
| F_fleet_3_YR_2011_s_1 | 4.96e-06 | (0,2.9) | 5.09e- | 0.103 | | 3 |
| F_fleet_3_YR_2012_s_1 | 2.18e-05 | (0,2.9) | 2.28e- | 0.105 | | 3 |
| F_fleet_3_YR_2013_s_1 | 8.09e-05 | (0,2.9) | 8.69e- | 0.107 | | 3 |
| F_fleet_3_YR_2014_s_1 | 3.34e-05 | (0,2.9) | 3.59e- | 0.108 | | 3 |
| F_fleet_3_YR_2015_s_1 | 1.23e-05 | (0,2.9) | 1.32e- | 0.108 | | 3 |
| F_fleet_3_YR_2016_s_1 | 9.72e-06 | (0,2.9) | 1.07e- | 0.11 | | 3 |
| F_fleet_3_YR_2017_s_1 | 9.72e-06 | (0,2.9) | 1.10e- | 0.113 | | 3 |
| F_fleet_3_YR_2018_s_1 | 4.76e-06 | (0,2.9) | 5.46e- | 0.115 | | 3 |
| F_fleet_3_YR_2019_s_1 | 1.25e-05 | (0,2.9) | 1.44e- | 0.115 | | 3 |
| F_fleet_3_YR_2020_s_1 | 1.79e-05 | (0,2.9) | 2.07e- | 0.115 | | 3 |
| F_fleet_4_YR_1946_s_1 | 0.018 | (0,2.9) | 0.003 | 0.147 | | 3 |
| F_fleet_4_YR_1947_s_1 | 0.037 | (0,2.9) | 0.005 | 0.148 | | 3 |
| F_fleet_4_YR_1948_s_1 | 0.056 | (0,2.9) | 0.008 | 0.148 | | 3 |
| F_fleet_4_YR_1949_s_1 | 0.075 | (0,2.9) | 0.011 | 0.148 | | 3 |
| F_fleet_4_YR_1950_s_1 | 0.094 | (0,2.9) | 0.014 | 0.148 | | 3 |
| F_fleet_4_YR_1951_s_1 | 0.114 | (0,2.9) | 0.017 | 0.148 | | 3 |
| F_fleet_4_YR_1952_s_1 | 0.133 | (0,2.9) | 0.022 | 0.164 | | 3 |
| F_fleet_4_YR_1953_s_1 | 0.153 | (0,2.9) | 0.028 | 0.184 | | 3 |
| F_fleet_4_YR_1954_s_1 | 0.173 | (0,2.9) | 0.034 | 0.194 | | 3 |
| F_fleet_4_YR_1955_s_1 | 0.146 | (0,2.9) | 0.029 | 0.199 | | 3 |
| F_fleet_4_YR_1956_s_1 | 0.162 | (0,2.9) | 0.033 | 0.201 | | 3 |
| F_fleet_4_YR_1957_s_1 | 0.179 | (0,2.9) | 0.036 | 0.202 | | 3 |
| F_fleet_4_YR_1958_s_1 | 0.196 | (0,2.9) | 0.04 | 0.202 | | 3 |
| F_fleet_4_YR_1959_s_1 | 0.212 | (0,2.9) | 0.043 | 0.203 | | 3 |
| F_fleet_4_YR_1960_s_1 | 0.23 | (0,2.9) | 0.047 | 0.204 | | 3 |
| F_fleet_4_YR_1961_s_1 | 0.241 | (0,2.9) | 0.049 | 0.202 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|-------|-------|-------|-------|
| F_fleet_4_YR_1962_s_1 | 0.253 | (0,2.9) | 0.051 | 0.2 | | 3 |
| F_fleet_4_YR_1963_s_1 | 0.259 | (0,2.9) | 0.051 | 0.199 | | 3 |
| F_fleet_4_YR_1964_s_1 | 0.265 | (0,2.9) | 0.052 | 0.198 | | 3 |
| F_fleet_4_YR_1965_s_1 | 0.282 | (0,2.9) | 0.055 | 0.194 | | 3 |
| F_fleet_4_YR_1966_s_1 | 0.305 | (0,2.9) | 0.058 | 0.189 | | 3 |
| F_fleet_4_YR_1967_s_1 | 0.329 | (0,2.9) | 0.061 | 0.186 | | 3 |
| F_fleet_4_YR_1968_s_1 | 0.351 | (0,2.9) | 0.064 | 0.184 | | 3 |
| F_fleet_4_YR_1969_s_1 | 0.378 | (0,2.9) | 0.069 | 0.182 | | 3 |
| F_fleet_4_YR_1970_s_1 | 0.409 | (0,2.9) | 0.074 | 0.181 | | 3 |
| F_fleet_4_YR_1971_s_1 | 0.463 | (0,2.9) | 0.084 | 0.181 | | 3 |
| F_fleet_4_YR_1972_s_1 | 0.505 | (0,2.9) | 0.092 | 0.182 | | 3 |
| F_fleet_4_YR_1973_s_1 | 0.535 | (0,2.9) | 0.097 | 0.182 | | 3 |
| F_fleet_4_YR_1974_s_1 | 0.563 | (0,2.9) | 0.102 | 0.181 | | 3 |
| F_fleet_4_YR_1975_s_1 | 0.594 | (0,2.9) | 0.105 | 0.177 | | 3 |
| F_fleet_4_YR_1976_s_1 | 0.61 | (0,2.9) | 0.105 | 0.172 | | 3 |
| F_fleet_4_YR_1977_s_1 | 0.64 | (0,2.9) | 0.108 | 0.168 | | 3 |
| F_fleet_4_YR_1978_s_1 | 0.684 | (0,2.9) | 0.113 | 0.165 | | 3 |
| F_fleet_4_YR_1979_s_1 | 0.74 | (0,2.9) | 0.12 | 0.161 | | 3 |
| F_fleet_4_YR_1980_s_1 | 0.815 | (0,2.9) | 0.13 | 0.159 | | 3 |
| F_fleet_4_YR_1981_s_1 | 1.44 | (0,2.9) | 0.222 | 0.154 | | 3 |
| F_fleet_4_YR_1982_s_1 | 1.07 | (0,2.9) | 0.161 | 0.15 | | 3 |
| F_fleet_4_YR_1983_s_1 | 0.259 | (0,2.9) | 0.04 | 0.154 | | 3 |
| F_fleet_4_YR_1984_s_1 | 1.34 | (0,2.9) | 0.196 | 0.147 | | 3 |
| F_fleet_4_YR_1985_s_1 | 0.573 | (0,2.9) | 0.09 | 0.157 | | 3 |
| F_fleet_4_YR_1986_s_1 | 0.454 | (0,2.9) | 0.07 | 0.154 | | 3 |
| F_fleet_4_YR_1987_s_1 | 0.947 | (0,2.9) | 0.138 | 0.146 | | 3 |
| F_fleet_4_YR_1988_s_1 | 0.82 | (0,2.9) | 0.125 | 0.152 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|----------------------------|----------------------------|-------|-------|
| F_fleet_4_YR_1989_s_1 | 1.29 | (0,2.9) | 0.19 | 0.147 | | 3 |
| F_fleet_4_YR_1990_s_1 | 2.05 | (0,2.9) | 0.287 | 0.14 | | 3 |
| F_fleet_4_YR_1991_s_1 | 2.12 | (0,2.9) | 0.309 | 0.146 | | 3 |
| F_fleet_4_YR_1992_s_1 | 1.6 | (0,2.9) | 0.235 | 0.147 | | 3 |
| F_fleet_4_YR_1993_s_1 | 1.89 | (0,2.9) | 0.278 | 0.147 | | 3 |
| F_fleet_4_YR_1994_s_1 | 1.58 | (0,2.9) | 0.234 | 0.149 | | 3 |
| F_fleet_4_YR_1995_s_1 | 2.2 | (0,2.9) | 0.323 | 0.147 | | 3 |
| F_fleet_4_YR_1996_s_1 | 1.8 | (0,2.9) | 0.269 | 0.149 | | 3 |
| F_fleet_4_YR_1997_s_1 | 1.84 | (0,2.9) | 0.272 | 0.148 | | 3 |
| F_fleet_4_YR_1998_s_1 | 2.36 | (0,2.9) | 0.342 | 0.145 | | 3 |
| F_fleet_4_YR_1999_s_1 | 1.76 | (0,2.9) | 0.257 | 0.146 | | 3 |
| F_fleet_4_YR_2000_s_1 | 1.46 | (0,2.9) | 0.214 | 0.147 | | 3 |
| F_fleet_4_YR_2001_s_1 | 1.73 | (0,2.9) | 0.252 | 0.145 | | 3 |
| F_fleet_4_YR_2002_s_1 | 1.54 | (0,2.9) | 0.226 | 0.147 | | 3 |
| F_fleet_4_YR_2003_s_1 | 2.1 | (0,2.9) | 0.307 | 0.146 | | 3 |
| F_fleet_4_YR_2004_s_1 | 2.62 | (0,2.9) | 0.372 | 0.142 | | 3 |
| F_fleet_4_YR_2005_s_1 | 1.76 | (0,2.9) | 0.259 | 0.147 | | 3 |
| F_fleet_4_YR_2006_s_1 | 1.51 | (0,2.9) | 0.221 | 0.147 | | 3 |
| F_fleet_4_YR_2007_s_1 | 1.6 | (0,2.9) | 0.231 | 0.144 | | 3 |
| F_fleet_4_YR_2008_s_1 | 2.18 | (0,2.9) | 0.321 | 0.147 | | 3 |
| F_fleet_4_YR_2009_s_1 | 2.3 | (0,2.9) | 0.337 | 0.146 | | 3 |
| F_fleet_4_YR_2010_s_1 | 1.22 | (0,2.9) | 0.182 | 0.149 | | 3 |
| F_fleet_4_YR_2011_s_1 | 1.28 | (0,2.9) | 0.19 | 0.148 | | 3 |
| F_fleet_4_YR_2012_s_1 | 2.9 | (0,2.9) | 6.29e- $\hat{\alpha}_1$ | 2.17e- $\hat{\alpha}_1$ | | 3 |
| F_fleet_4_YR_2013_s_1 | 2.31 | (0,2.9) | 0.328 | 0.142 | | 3 |
| F_fleet_4_YR_2014_s_1 | 2.52 | (0,2.9) | 0.361 | 0.143 | | 3 |
| F_fleet_4_YR_2015_s_1 | 2.28 | (0,2.9) | 0.327 | 0.144 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|----------------------|-------|-------|-------|
| F_fleet_4_YR_2016_s_1 | 2.37 | (0,2.9) | 0.34 | 0.143 | | 3 |
| F_fleet_4_YR_2017_s_1 | 1.64 | (0,2.9) | 0.243 | 0.148 | | 3 |
| F_fleet_4_YR_2018_s_1 | 1.62 | (0,2.9) | 0.239 | 0.147 | | 3 |
| F_fleet_4_YR_2019_s_1 | 1.34 | (0,2.9) | 0.198 | 0.148 | | 3 |
| F_fleet_4_YR_2020_s_1 | 1.59 | (0,2.9) | 0.243 | 0.152 | | 3 |
| F_fleet_5_YR_1946_s_1 | 0.002 | (0,2.9) | 2.53e- ^{^4} | 0.121 | | 3 |
| F_fleet_5_YR_1947_s_1 | 0.004 | (0,2.9) | 5.07e- ^{^4} | 0.121 | | 3 |
| F_fleet_5_YR_1948_s_1 | 0.006 | (0,2.9) | 7.64e- ^{^4} | 0.121 | | 3 |
| F_fleet_5_YR_1949_s_1 | 0.008 | (0,2.9) | 0.001 | 0.121 | | 3 |
| F_fleet_5_YR_1950_s_1 | 0.011 | (0,2.9) | 0.001 | 0.121 | | 3 |
| F_fleet_5_YR_1951_s_1 | 0.013 | (0,2.9) | 0.002 | 0.121 | | 3 |
| F_fleet_5_YR_1952_s_1 | 0.015 | (0,2.9) | 0.003 | 0.218 | | 3 |
| F_fleet_5_YR_1953_s_1 | 0.017 | (0,2.9) | 0.004 | 0.24 | | 3 |
| F_fleet_5_YR_1954_s_1 | 0.019 | (0,2.9) | 0.005 | 0.246 | | 3 |
| F_fleet_5_YR_1955_s_1 | 0.016 | (0,2.9) | 0.004 | 0.248 | | 3 |
| F_fleet_5_YR_1956_s_1 | 0.018 | (0,2.9) | 0.004 | 0.248 | | 3 |
| F_fleet_5_YR_1957_s_1 | 0.02 | (0,2.9) | 0.005 | 0.248 | | 3 |
| F_fleet_5_YR_1958_s_1 | 0.021 | (0,2.9) | 0.005 | 0.248 | | 3 |
| F_fleet_5_YR_1959_s_1 | 0.023 | (0,2.9) | 0.006 | 0.25 | | 3 |
| F_fleet_5_YR_1960_s_1 | 0.025 | (0,2.9) | 0.006 | 0.251 | | 3 |
| F_fleet_5_YR_1961_s_1 | 0.027 | (0,2.9) | 0.007 | 0.245 | | 3 |
| F_fleet_5_YR_1962_s_1 | 0.028 | (0,2.9) | 0.007 | 0.241 | | 3 |
| F_fleet_5_YR_1963_s_1 | 0.027 | (0,2.9) | 0.007 | 0.244 | | 3 |
| F_fleet_5_YR_1964_s_1 | 0.028 | (0,2.9) | 0.007 | 0.238 | | 3 |
| F_fleet_5_YR_1965_s_1 | 0.032 | (0,2.9) | 0.007 | 0.226 | | 3 |
| F_fleet_5_YR_1966_s_1 | 0.035 | (0,2.9) | 0.008 | 0.221 | | 3 |
| F_fleet_5_YR_1967_s_1 | 0.037 | (0,2.9) | 0.008 | 0.219 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|-------|-------|-------|-------|
| F_fleet_5_YR_1968_s_1 | 0.04 | (0,2.9) | 0.009 | 0.218 | | 3 |
| F_fleet_5_YR_1969_s_1 | 0.043 | (0,2.9) | 0.009 | 0.214 | | 3 |
| F_fleet_5_YR_1970_s_1 | 0.047 | (0,2.9) | 0.01 | 0.214 | | 3 |
| F_fleet_5_YR_1971_s_1 | 0.052 | (0,2.9) | 0.011 | 0.216 | | 3 |
| F_fleet_5_YR_1972_s_1 | 0.054 | (0,2.9) | 0.012 | 0.216 | | 3 |
| F_fleet_5_YR_1973_s_1 | 0.054 | (0,2.9) | 0.012 | 0.212 | | 3 |
| F_fleet_5_YR_1974_s_1 | 0.057 | (0,2.9) | 0.012 | 0.204 | | 3 |
| F_fleet_5_YR_1975_s_1 | 0.059 | (0,2.9) | 0.011 | 0.193 | | 3 |
| F_fleet_5_YR_1976_s_1 | 0.063 | (0,2.9) | 0.012 | 0.185 | | 3 |
| F_fleet_5_YR_1977_s_1 | 0.067 | (0,2.9) | 0.012 | 0.179 | | 3 |
| F_fleet_5_YR_1978_s_1 | 0.074 | (0,2.9) | 0.013 | 0.172 | | 3 |
| F_fleet_5_YR_1979_s_1 | 0.081 | (0,2.9) | 0.013 | 0.167 | | 3 |
| F_fleet_5_YR_1980_s_1 | 0.091 | (0,2.9) | 0.015 | 0.163 | | 3 |
| F_fleet_5_YR_1981_s_1 | 0.129 | (0,2.9) | 0.021 | 0.16 | | 3 |
| F_fleet_5_YR_1982_s_1 | 0.116 | (0,2.9) | 0.018 | 0.152 | | 3 |
| F_fleet_5_YR_1983_s_1 | 0.095 | (0,2.9) | 0.014 | 0.151 | | 3 |
| F_fleet_5_YR_1984_s_1 | 0.095 | (0,2.9) | 0.014 | 0.146 | | 3 |
| F_fleet_5_YR_1985_s_1 | 0.046 | (0,2.9) | 0.007 | 0.148 | | 3 |
| F_fleet_5_YR_1986_s_1 | 0.069 | (0,2.9) | 0.009 | 0.128 | | 3 |
| F_fleet_5_YR_1987_s_1 | 0.075 | (0,2.9) | 0.009 | 0.124 | | 3 |
| F_fleet_5_YR_1988_s_1 | 0.109 | (0,2.9) | 0.015 | 0.136 | | 3 |
| F_fleet_5_YR_1989_s_1 | 0.152 | (0,2.9) | 0.019 | 0.123 | | 3 |
| F_fleet_5_YR_1990_s_1 | 0.309 | (0,2.9) | 0.03 | 0.096 | | 3 |
| F_fleet_5_YR_1991_s_1 | 1.14 | (0,2.9) | 0.11 | 0.097 | | 3 |
| F_fleet_5_YR_1992_s_1 | 0.398 | (0,2.9) | 0.042 | 0.106 | | 3 |
| F_fleet_5_YR_1993_s_1 | 0.379 | (0,2.9) | 0.043 | 0.113 | | 3 |
| F_fleet_5_YR_1994_s_1 | 0.278 | (0,2.9) | 0.034 | 0.123 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|--------|-------|-------|-------|
| F_fleet_5_YR_1995_s_1 | 0.298 | (0,2.9) | 0.037 | 0.124 | | 3 |
| F_fleet_5_YR_1996_s_1 | 0.499 | (0,2.9) | 0.06 | 0.12 | | 3 |
| F_fleet_5_YR_1997_s_1 | 0.257 | (0,2.9) | 0.032 | 0.124 | | 3 |
| F_fleet_5_YR_1998_s_1 | 0.218 | (0,2.9) | 0.026 | 0.121 | | 3 |
| F_fleet_5_YR_1999_s_1 | 0.151 | (0,2.9) | 0.019 | 0.127 | | 3 |
| F_fleet_5_YR_2000_s_1 | 0.207 | (0,2.9) | 0.026 | 0.125 | | 3 |
| F_fleet_5_YR_2001_s_1 | 0.271 | (0,2.9) | 0.033 | 0.122 | | 3 |
| F_fleet_5_YR_2002_s_1 | 0.147 | (0,2.9) | 0.018 | 0.126 | | 3 |
| F_fleet_5_YR_2003_s_1 | 0.208 | (0,2.9) | 0.027 | 0.13 | | 3 |
| F_fleet_5_YR_2004_s_1 | 0.226 | (0,2.9) | 0.029 | 0.129 | | 3 |
| F_fleet_5_YR_2005_s_1 | 0.222 | (0,2.9) | 0.029 | 0.132 | | 3 |
| F_fleet_5_YR_2006_s_1 | 0.104 | (0,2.9) | 0.014 | 0.131 | | 3 |
| F_fleet_5_YR_2007_s_1 | 0.138 | (0,2.9) | 0.018 | 0.13 | | 3 |
| F_fleet_5_YR_2008_s_1 | 0.237 | (0,2.9) | 0.031 | 0.131 | | 3 |
| F_fleet_5_YR_2009_s_1 | 0.177 | (0,2.9) | 0.023 | 0.127 | | 3 |
| F_fleet_5_YR_2010_s_1 | 0.051 | (0,2.9) | 0.007 | 0.135 | | 3 |
| F_fleet_5_YR_2011_s_1 | 0.157 | (0,2.9) | 0.02 | 0.13 | | 3 |
| F_fleet_5_YR_2012_s_1 | 0.247 | (0,2.9) | 0.03 | 0.121 | | 3 |
| F_fleet_5_YR_2013_s_1 | 0.199 | (0,2.9) | 0.025 | 0.128 | | 3 |
| F_fleet_5_YR_2014_s_1 | 0.28 | (0,2.9) | 0.036 | 0.127 | | 3 |
| F_fleet_5_YR_2015_s_1 | 0.209 | (0,2.9) | 0.028 | 0.133 | | 3 |
| F_fleet_5_YR_2016_s_1 | 0.327 | (0,2.9) | 0.044 | 0.133 | | 3 |
| F_fleet_5_YR_2017_s_1 | 0.404 | (0,2.9) | 0.053 | 0.13 | | 3 |
| F_fleet_5_YR_2018_s_1 | 0.334 | (0,2.9) | 0.044 | 0.133 | | 3 |
| F_fleet_5_YR_2019_s_1 | 0.383 | (0,2.9) | 0.047 | 0.123 | | 3 |
| F_fleet_5_YR_2020_s_1 | 0.268 | (0,2.9) | 0.037 | 0.139 | | 3 |
| F_fleet_6_YR_1946_s_1 | 0.001 | (0,2.9) | 3.32e- | 0.326 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|--------|-------|-------|-------|
| F_fleet_6_YR_1947_s_1 | 0.002 | (0,2.9) | 6.67e- | 0.327 | | 3 |
| F_fleet_6_YR_1948_s_1 | 0.003 | (0,2.9) | 0.001 | 0.327 | | 3 |
| F_fleet_6_YR_1949_s_1 | 0.004 | (0,2.9) | 0.001 | 0.327 | | 3 |
| F_fleet_6_YR_1950_s_1 | 0.005 | (0,2.9) | 0.002 | 0.327 | | 3 |
| F_fleet_6_YR_1951_s_1 | 0.006 | (0,2.9) | 0.002 | 0.327 | | 3 |
| F_fleet_6_YR_1952_s_1 | 0.007 | (0,2.9) | 0.002 | 0.327 | | 3 |
| F_fleet_6_YR_1953_s_1 | 0.009 | (0,2.9) | 0.003 | 0.329 | | 3 |
| F_fleet_6_YR_1954_s_1 | 0.01 | (0,2.9) | 0.003 | 0.333 | | 3 |
| F_fleet_6_YR_1955_s_1 | 0.008 | (0,2.9) | 0.003 | 0.337 | | 3 |
| F_fleet_6_YR_1956_s_1 | 0.009 | (0,2.9) | 0.003 | 0.34 | | 3 |
| F_fleet_6_YR_1957_s_1 | 0.01 | (0,2.9) | 0.003 | 0.342 | | 3 |
| F_fleet_6_YR_1958_s_1 | 0.011 | (0,2.9) | 0.004 | 0.343 | | 3 |
| F_fleet_6_YR_1959_s_1 | 0.012 | (0,2.9) | 0.004 | 0.345 | | 3 |
| F_fleet_6_YR_1960_s_1 | 0.013 | (0,2.9) | 0.005 | 0.347 | | 3 |
| F_fleet_6_YR_1961_s_1 | 0.014 | (0,2.9) | 0.005 | 0.347 | | 3 |
| F_fleet_6_YR_1962_s_1 | 0.014 | (0,2.9) | 0.005 | 0.347 | | 3 |
| F_fleet_6_YR_1963_s_1 | 0.015 | (0,2.9) | 0.005 | 0.348 | | 3 |
| F_fleet_6_YR_1964_s_1 | 0.015 | (0,2.9) | 0.005 | 0.35 | | 3 |
| F_fleet_6_YR_1965_s_1 | 0.016 | (0,2.9) | 0.006 | 0.352 | | 3 |
| F_fleet_6_YR_1966_s_1 | 0.017 | (0,2.9) | 0.006 | 0.353 | | 3 |
| F_fleet_6_YR_1967_s_1 | 0.018 | (0,2.9) | 0.006 | 0.353 | | 3 |
| F_fleet_6_YR_1968_s_1 | 0.019 | (0,2.9) | 0.007 | 0.351 | | 3 |
| F_fleet_6_YR_1969_s_1 | 0.02 | (0,2.9) | 0.007 | 0.349 | | 3 |
| F_fleet_6_YR_1970_s_1 | 0.022 | (0,2.9) | 0.008 | 0.348 | | 3 |
| F_fleet_6_YR_1971_s_1 | 0.025 | (0,2.9) | 0.009 | 0.347 | | 3 |
| F_fleet_6_YR_1972_s_1 | 0.028 | (0,2.9) | 0.01 | 0.347 | | 3 |
| F_fleet_6_YR_1973_s_1 | 0.031 | (0,2.9) | 0.011 | 0.348 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-----------------------|-------|---------|-------|-------|-------|-------|
| F_fleet_6_YR_1974_s_1 | 0.034 | (0,2.9) | 0.012 | 0.35 | | 3 |
| F_fleet_6_YR_1975_s_1 | 0.037 | (0,2.9) | 0.013 | 0.352 | | 3 |
| F_fleet_6_YR_1976_s_1 | 0.037 | (0,2.9) | 0.013 | 0.354 | | 3 |
| F_fleet_6_YR_1977_s_1 | 0.038 | (0,2.9) | 0.014 | 0.355 | | 3 |
| F_fleet_6_YR_1978_s_1 | 0.04 | (0,2.9) | 0.014 | 0.355 | | 3 |
| F_fleet_6_YR_1979_s_1 | 0.042 | (0,2.9) | 0.015 | 0.355 | | 3 |
| F_fleet_6_YR_1980_s_1 | 0.045 | (0,2.9) | 0.016 | 0.354 | | 3 |
| F_fleet_6_YR_1981_s_1 | 0.043 | (0,2.9) | 0.015 | 0.354 | | 3 |
| F_fleet_6_YR_1982_s_1 | 0.145 | (0,2.9) | 0.052 | 0.356 | | 3 |
| F_fleet_6_YR_1983_s_1 | 0.104 | (0,2.9) | 0.037 | 0.357 | | 3 |
| F_fleet_6_YR_1984_s_1 | 0.025 | (0,2.9) | 0.009 | 0.355 | | 3 |
| F_fleet_6_YR_1985_s_1 | 0.026 | (0,2.9) | 0.009 | 0.356 | | 3 |
| F_fleet_6_YR_1986_s_1 | 0.041 | (0,2.9) | 0.015 | 0.353 | | 3 |
| F_fleet_6_YR_1987_s_1 | 0.025 | (0,2.9) | 0.009 | 0.35 | | 3 |
| F_fleet_6_YR_1988_s_1 | 0.015 | (0,2.9) | 0.005 | 0.35 | | 3 |
| F_fleet_6_YR_1989_s_1 | 0.034 | (0,2.9) | 0.012 | 0.351 | | 3 |
| F_fleet_6_YR_1990_s_1 | 0.029 | (0,2.9) | 0.01 | 0.351 | | 3 |
| F_fleet_6_YR_1991_s_1 | 0.079 | (0,2.9) | 0.028 | 0.352 | | 3 |
| F_fleet_6_YR_1992_s_1 | 0.07 | (0,2.9) | 0.024 | 0.349 | | 3 |
| F_fleet_6_YR_1993_s_1 | 0.039 | (0,2.9) | 0.014 | 0.348 | | 3 |
| F_fleet_6_YR_1994_s_1 | 0.053 | (0,2.9) | 0.018 | 0.35 | | 3 |
| F_fleet_6_YR_1995_s_1 | 0.038 | (0,2.9) | 0.013 | 0.355 | | 3 |
| F_fleet_6_YR_1996_s_1 | 0.039 | (0,2.9) | 0.014 | 0.35 | | 3 |
| F_fleet_6_YR_1997_s_1 | 0.028 | (0,2.9) | 0.01 | 0.356 | | 3 |
| F_fleet_6_YR_1998_s_1 | 0.075 | (0,2.9) | 0.027 | 0.356 | | 3 |
| F_fleet_6_YR_1999_s_1 | 0.058 | (0,2.9) | 0.021 | 0.355 | | 3 |
| F_fleet_6_YR_2000_s_1 | 0.039 | (0,2.9) | 0.014 | 0.356 | | 3 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|-------------------------------|--------|----------|-------|-------|-------|-------|
| F_fleet_6_YR_2001_s_1 | 0.066 | (0,2.9) | 0.023 | 0.353 | | 3 |
| F_fleet_6_YR_2002_s_1 | 0.049 | (0,2.9) | 0.018 | 0.355 | | 3 |
| F_fleet_6_YR_2003_s_1 | 0.077 | (0,2.9) | 0.027 | 0.354 | | 3 |
| F_fleet_6_YR_2004_s_1 | 0.067 | (0,2.9) | 0.024 | 0.354 | | 3 |
| F_fleet_6_YR_2005_s_1 | 0.065 | (0,2.9) | 0.023 | 0.356 | | 3 |
| F_fleet_6_YR_2006_s_1 | 0.061 | (0,2.9) | 0.022 | 0.356 | | 3 |
| F_fleet_6_YR_2007_s_1 | 0.06 | (0,2.9) | 0.021 | 0.356 | | 3 |
| F_fleet_6_YR_2008_s_1 | 0.063 | (0,2.9) | 0.023 | 0.357 | | 3 |
| F_fleet_6_YR_2009_s_1 | 0.08 | (0,2.9) | 0.028 | 0.356 | | 3 |
| F_fleet_6_YR_2010_s_1 | 0.042 | (0,2.9) | 0.015 | 0.359 | | 3 |
| F_fleet_6_YR_2011_s_1 | 0.049 | (0,2.9) | 0.018 | 0.357 | | 3 |
| F_fleet_6_YR_2012_s_1 | 0.05 | (0,2.9) | 0.018 | 0.356 | | 3 |
| F_fleet_6_YR_2013_s_1 | 0.088 | (0,2.9) | 0.031 | 0.357 | | 3 |
| F_fleet_6_YR_2014_s_1 | 0.092 | (0,2.9) | 0.033 | 0.357 | | 3 |
| F_fleet_6_YR_2015_s_1 | 0.07 | (0,2.9) | 0.025 | 0.358 | | 3 |
| F_fleet_6_YR_2016_s_1 | 0.1 | (0,2.9) | 0.036 | 0.358 | | 3 |
| F_fleet_6_YR_2017_s_1 | 0.087 | (0,2.9) | 0.031 | 0.361 | | 3 |
| F_fleet_6_YR_2018_s_1 | 0.085 | (0,2.9) | 0.031 | 0.362 | | 3 |
| F_fleet_6_YR_2019_s_1 | 0.094 | (0,2.9) | 0.034 | 0.361 | | 3 |
| F_fleet_6_YR_2020_s_1 | 0.069 | (0,2.9) | 0.025 | 0.363 | | 3 |
| LnQ_base_Index_Private_7(7) | -8.73 | (-25,25) | | | | Float |
| LnQ_base_Index_Shore_8(8) | -9.69 | (-25,25) | | | | Float |
| LnQ_base_FWRI_Age0_9(9) | -10.15 | (-25,25) | | | | Float |
| LnQ_base_FWRI_Age1_10(10) | -11.19 | (-25,25) | | | | Float |
| LnQ_base_SEAMAP_Trawl_11(11) | -10.16 | (-25,25) | | | | Float |
| LnQ_base_Combo_Video_12(12) | -10.01 | (-25,25) | | | | Float |
| LnQ_base_Visual_Survey_13(13) | -10.46 | (-25,25) | | | | Float |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|--------------------------------------|----------|------------|-------|--------|-------|-------|
| Size_DblN_peak_Com_VL_1(1) | 29.04 | (13,60) | 0.042 | 0.001 | | 2 |
| Size_DblN_top_logit_Com_VL_1(1) | -11 | (-15,15) | 60.19 | -5.47 | | 2 |
| Size_DblN_ascend_se_Com_VL_1(1) | -8.66 | (-25,10) | 110.6 | -12.78 | | 3 |
| Size_DblN_descend_se_Com_VL_1(1) | 4.16 | (0,15) | 0.234 | 0.056 | | 3 |
| Size_DblN_start_logit_Com_VL_1(1) | -1.55 | (-999,999) | 0.197 | -0.127 | | 2 |
| Size_DblN_end_logit_Com_VL_1(1) | -2.15 | (-999,999) | 0.313 | -0.146 | | 2 |
| Retain_L_infl_Com_VL_1(1) | 12.8 | (10,80) | 12.87 | 1.01 | | 4 |
| Retain_L_width_Com_VL_1(1) | 1.36 | (1e-04,15) | 2.41 | 1.77 | | 4 |
| Retain_L_asymptote_logit_Com_VL_1(1) | 10 | | | | | Fixed |
| Retain_L_maleoffset_Com_VL_1(1) | 0.00e+00 | | | | | Fixed |
| DiscMort_L_infl_Com_VL_1(1) | -5 | | | | | Fixed |
| DiscMort_L_width_Com_VL_1(1) | 1 | | | | | Fixed |
| DiscMort_L_level_old_Com_VL_1(1) | 0.14 | | | | | Fixed |
| DiscMort_L_male_offset_Com_VL_1(1) | 0.00e+00 | | | | | Fixed |
| Size_DblN_peak_Com_LL_2(2) | 41.56 | (13,60) | 1.28 | 0.031 | | 2 |
| Size_DblN_top_logit_Com_LL_2(2) | -10.89 | (-15,15) | 61.37 | -5.63 | | 2 |
| Size_DblN_ascend_se_Com_LL_2(2) | 3.29 | (-25,10) | 0.655 | 0.199 | | 3 |
| Size_DblN_descend_se_Com_LL_2(2) | 5.5 | (0,15) | 0.189 | 0.034 | | 3 |
| Size_DblN_start_logit_Com_LL_2(2) | -0.641 | (-999,999) | 0.256 | -0.399 | | 2 |
| Size_DblN_end_logit_Com_LL_2(2) | -999 | | | | | Fixed |
| Retain_L_infl_Com_LL_2(2) | 12.53 | (10,80) | 54.58 | 4.36 | | 4 |
| Retain_L_width_Com_LL_2(2) | 0.916 | (1e-04,5) | 19.82 | 21.65 | | 4 |
| Retain_L_asymptote_logit_Com_LL_2(2) | 10 | | | | | Fixed |
| Retain_L_maleoffset_Com_LL_2(2) | 0.00e+00 | | | | | Fixed |
| DiscMort_L_infl_Com_LL_2(2) | -5 | | | | | Fixed |
| DiscMort_L_width_Com_LL_2(2) | 1 | | | | | Fixed |
| DiscMort_L_level_old_Com_LL_2(2) | 0.14 | | | | | Fixed |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|--------------------------------------|----------|----------|-------|--------|-------|-------|
| DiscMort_L_male_offset_Com_LL_2(2) | 0.00e+00 | | | | | Fixed |
| Size_DblN_peak_Com_NT_3(3) | 33.9 | | | | | Fixed |
| Size_DblN_top_logit_Com_NT_3(3) | -12.36 | (-15,15) | 45.3 | -3.66 | | 2 |
| Size_DblN_ascend_se_Com_NT_3(3) | 3.03 | (-25,20) | 0.079 | 0.026 | | 3 |
| Size_DblN_descend_se_Com_NT_3(3) | 4.37 | (-20,15) | 0.159 | 0.036 | | 3 |
| Size_DblN_start_logit_Com_NT_3(3) | -6.79 | (-15,15) | 0.713 | -0.105 | | 2 |
| Size_DblN_end_logit_Com_NT_3(3) | -2.71 | (-15,15) | 0.351 | -0.13 | | 2 |
| Size_DblN_peak_Rec_PR_4(4) | 13 | | | | | Fixed |
| Size_DblN_top_logit_Rec_PR_4(4) | -11.78 | (-15,15) | 51.89 | -4.4 | | 2 |
| Size_DblN_ascend_se_Rec_PR_4(4) | 0.444 | (-20,20) | 0.415 | 0.936 | | 3 |
| Size_DblN_descend_se_Rec_PR_4(4) | 6.18 | (-20,15) | 0.074 | 0.012 | | 3 |
| Size_DblN_start_logit_Rec_PR_4(4) | -999 | | | | | Fixed |
| Size_DblN_end_logit_Rec_PR_4(4) | -999 | | | | | Fixed |
| Retain_L_infl_Rec_PR_4(4) | 19.25 | (6,79) | 0.404 | 0.021 | | 3 |
| Retain_L_width_Rec_PR_4(4) | 0.903 | (-1,30) | 0.237 | 0.262 | | 3 |
| Retain_L_asymptote_logit_Rec_PR_4(4) | 10 | | | | | Fixed |
| Retain_L_maleoffset_Rec_PR_4(4) | 0.00e+00 | | | | | Fixed |
| DiscMort_L_infl_Rec_PR_4(4) | -5 | | | | | Fixed |
| DiscMort_L_width_Rec_PR_4(4) | 1 | | | | | Fixed |
| DiscMort_L_level_old_Rec_PR_4(4) | 0.069 | | | | | Fixed |
| DiscMort_L_male_offset_Rec_PR_4(4) | 0.00e+00 | | | | | Fixed |
| Size_DblN_peak_Rec_Shore_5(5) | 13 | | | | | Fixed |
| Size_DblN_top_logit_Rec_Shore_5(5) | -12.71 | (-15,15) | 41.11 | -3.23 | | 2 |
| Size_DblN_ascend_se_Rec_Shore_5(5) | -0.229 | (-20,20) | 0.228 | -0.994 | | 3 |
| Size_DblN_descend_se_Rec_Shore_5(5) | 4.98 | (-20,15) | 0.038 | 0.008 | | 3 |
| Size_DblN_start_logit_Rec_Shore_5(5) | -999 | | | | | Fixed |
| Size_DblN_end_logit_Rec_Shore_5(5) | -999 | | | | | Fixed |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|---|----------|------------|-------|-------|-------|-------|
| Retain_L_infl_Rec_Shore_5(5) | 14.1 | (10,79) | 0.758 | 0.054 | | 3 |
| Retain_L_width_Rec_Shore_5(5) | 1.67 | (-1,30) | 0.549 | 0.329 | | 3 |
| Retain_L_asymptote_logit_Rec_Shore_5(5) | 10 | | | | | Fixed |
| Retain_L_maleoffset_Rec_Shore_5(5) | 0.00e+00 | | | | | Fixed |
| DiscMort_L_infl_Rec_Shore_5(5) | -5 | | | | | Fixed |
| DiscMort_L_width_Rec_Shore_5(5) | 1 | | | | | Fixed |
| DiscMort_L_level_old_Rec_Shore_5(5) | 0.069 | | | | | Fixed |
| DiscMort_L_male_offset_Rec_Shore_5(5) | 0.00e+00 | | | | | Fixed |
| Size_DblN_peak_Rec_HB_CBT_6(6) | 29.04 | (13,60) | 0.511 | 0.018 | | 2 |
| Size_DblN_top_logit_Rec_HB_CBT_6(6) | -11.92 | (-15,15) | 50.34 | -4.22 | | 2 |
| Size_DblN_ascend_se_Rec_HB_CBT_6(6) | 3.38 | (-20,20) | 0.202 | 0.06 | | 3 |
| Size_DblN_descend_se_Rec_HB_CBT_6(6) | 6.38 | (-20,15) | 0.155 | 0.024 | | 3 |
| Size_DblN_start_logit_Rec_HB_CBT_6(6) | -999 | | | | | Fixed |
| Size_DblN_end_logit_Rec_HB_CBT_6(6) | -999 | | | | | Fixed |
| Retain_L_infl_Rec_HB_CBT_6(6) | 23.68 | (6,80) | 0.283 | 0.012 | | 4 |
| Retain_L_width_Rec_HB_CBT_6(6) | 1.39 | (1e-04,10) | 0.212 | 0.152 | | 4 |
| Retain_L_asymptote_logit_Rec_HB_CBT_6(6) | 10 | | | | | Fixed |
| Retain_L_maleoffset_Rec_HB_CBT_6(6) | 0.00e+00 | | | | | Fixed |
| DiscMort_L_infl_Rec_HB_CBT_6(6) | -5 | | | | | Fixed |
| DiscMort_L_width_Rec_HB_CBT_6(6) | 1 | | | | | Fixed |
| DiscMort_L_level_old_Rec_HB_CBT_6(6) | 0.069 | | | | | Fixed |
| DiscMort_L_male_offset_Rec_HB_CBT_6(6) | 0.00e+00 | | | | | Fixed |
| Size_DblN_peak_SEAMAP_Trawl_11(11) | 23.99 | (13,73) | 0.396 | 0.017 | | 2 |
| Size_DblN_top_logit_SEAMAP_Trawl_11(11) | -13.05 | (-15,15) | 36.78 | -2.82 | | 2 |
| Size_DblN_ascend_se_SEAMAP_Trawl_11(11) | 3.04 | (-25,20) | 0.145 | 0.048 | | 3 |
| Size_DblN_descend_se_SEAMAP_Trawl_11(11) | 5.27 | (-20,15) | 0.118 | 0.022 | | 3 |
| Size_DblN_start_logit_SEAMAP_Trawl_11(11) | -999 | | | | | Fixed |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|--|--------|----------|-------|--------|----------------|-------|
| Size_DblN_end_logit_SEAMAP_Trawl_11(1) | -999 | | | | | Fixed |
| Size_inflection_Combo_Video_12(12) | 24.75 | (10,85) | 0.411 | 0.017 | | 2 |
| Size_95%width_Combo_Video_12(12) | 6.27 | (0,50) | 0.512 | 0.082 | | 2 |
| Size_DblN_peak_Visual_Survey_13(13) | 13.69 | (13,73) | 0.418 | 0.03 | | 2 |
| Size_DblN_top_logit_Visual_Survey_13(13) | -10.71 | (-15,15) | 63.24 | -5.91 | | 2 |
| Size_DblN_ascend_se_Visual_Survey_13(13) | 0.695 | (-25,20) | 0.319 | 0.459 | | 3 |
| Size_DblN_descend_se_Visual_Survey_13(13) | 7.66 | (-20,15) | 0.211 | 0.028 | | 3 |
| Size_DblN_start_logit_Visual_Survey_13(13) | -999 | | | | | Fixed |
| Size_DblN_end_logit_Visual_Survey_13(13) | -999 | | | | | Fixed |
| Age_DblN_peak_Com_VL_1(1) | 19.89 | | | | | Fixed |
| Age_DblN_top_logit_Com_VL_1(1) | -0.26 | (-8,3) | 1.36 | -5.23 | Sym_Beta(0.05) | 3 |
| Age_DblN_ascend_se_Com_VL_1(1) | 4.41 | (-4,12) | 0.092 | 0.021 | Sym_Beta(0.05) | 3 |
| Age_DblN_descend_se_Com_VL_1(1) | 0.255 | (-2,6) | 5.2 | 20.34 | Sym_Beta(0.05) | 3 |
| Age_DblN_start_logit_Com_VL_1(1) | -11.27 | (-15,5) | 8.15 | -0.723 | Sym_Beta(0.05) | 2 |
| Age_DblN_end_logit_Com_VL_1(1) | -4.11 | (-5,5) | 2.94 | -0.716 | Sym_Beta(0.05) | 2 |
| Age_DblN_peak_Com_LL_2(2) | 18 | (0,19.8) | 0.619 | 0.034 | Sym_Beta(0.05) | 2 |
| Age_DblN_top_logit_Com_LL_2(2) | -4.89 | (-8,3) | 7.84 | -1.6 | Sym_Beta(0.05) | 3 |
| Age_DblN_ascend_se_Com_LL_2(2) | 3.82 | (-4,12) | 0.099 | 0.026 | Sym_Beta(0.05) | 3 |
| Age_DblN_descend_se_Com_LL_2(2) | 2.16 | (-2,6) | 3.28 | 1.52 | Sym_Beta(0.05) | 3 |
| Age_DblN_start_logit_Com_LL_2(2) | -10.16 | (-15,5) | 10.11 | -0.996 | Sym_Beta(0.05) | 2 |
| Age_DblN_end_logit_Com_LL_2(2) | -0.588 | (-5,5) | 2.47 | -4.2 | Sym_Beta(0.05) | 2 |
| Age_DblN_peak_Rec_PR_4(4) | 21.21 | (0,25) | 1.42 | 0.067 | Sym_Beta(0.05) | 2 |
| Age_DblN_top_logit_Rec_PR_4(4) | -2.18 | (-8,3) | 20.16 | -9.25 | Sym_Beta(0.05) | 3 |
| Age_DblN_ascend_se_Rec_PR_4(4) | 3.29 | (-4,12) | 0.389 | 0.118 | Sym_Beta(0.05) | 3 |
| Age_DblN_descend_se_Rec_PR_4(4) | 2.34 | (-2,6) | 11.37 | 4.86 | Sym_Beta(0.05) | 3 |
| Age_DblN_start_logit_Rec_PR_4(4) | -2.12 | (-15,5) | 0.122 | -0.057 | Sym_Beta(0.05) | 2 |
| Age_DblN_end_logit_Rec_PR_4(4) | 3.86 | (-5,5) | 3.61 | 0.936 | Sym_Beta(0.05) | 2 |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|--|--------|---------|-------|--------|---------------|-------|
| Age_DblN_peak_Rec_HB_CBT_6(6) | 21.42 | (0,25) | 2.82 | 0.131 | Sym_Beta(0.05 | 2 |
| Age_DblN_top_logit_Rec_HB_CBT_6(6) | -0.267 | (-8,3) | 3.05 | -11.43 | Sym_Beta(0.05 | 3 |
| Age_DblN_ascend_se_Rec_HB_CBT_6(6) | 3.85 | (-4,12) | 0.498 | 0.13 | Sym_Beta(0.05 | 3 |
| Age_DblN_descend_se_Rec_HB_CBT_6(6) | -0.017 | (-2,6) | 7.08 | -415.4 | Sym_Beta(0.05 | 3 |
| Age_DblN_start_logit_Rec_HB_CBT_6(6) | -1.13 | (-15,5) | 0.42 | -0.371 | Sym_Beta(0.05 | 2 |
| Age_DblN_end_logit_Rec_HB_CBT_6(6) | -4.04 | (-5,5) | 3.13 | -0.774 | Sym_Beta(0.05 | 2 |
| ln(DM_theta)_1 | 0.178 | (-5,10) | 0.219 | 1.23 | Normal(0,1) | 6 |
| ln(DM_theta)_2 | 3.75 | (-5,10) | 0.549 | 0.146 | Normal(0,1) | 6 |
| ln(DM_theta)_3 | -1.24 | (-5,10) | 0.13 | -0.105 | Normal(0,1) | 6 |
| ln(DM_theta)_4 | 2.02 | (-5,10) | 0.466 | 0.231 | Normal(0,1) | 6 |
| ln(DM_theta)_5 | 3.86 | (-5,10) | 0.357 | 0.092 | Normal(0,1) | 6 |
| ln(DM_theta)_6 | 1.22 | (-5,10) | 0.443 | 0.362 | Normal(0,1) | 6 |
| ln(DM_theta)_7 | 2.44 | (-5,10) | 0.542 | 0.222 | Normal(0,1) | 6 |
| ln(DM_theta)_8 | 1.88 | (-5,10) | 0.473 | 0.252 | Normal(0,1) | 6 |
| ln(DM_theta)_9 | -1.4 | (-5,10) | 0.096 | -0.068 | Normal(0,1) | 6 |
| ln(DM_theta)_10 | -0.721 | (-5,10) | 0.094 | -0.131 | Normal(0,1) | 6 |
| ln(DM_theta)_11 | 2.44 | (-5,10) | 0.468 | 0.191 | Normal(0,1) | 6 |
| ln(DM_theta)_12 | -0.364 | (-5,10) | 0.149 | -0.408 | Normal(0,1) | 6 |
| ln(DM_theta)_13 | -0.683 | (-5,10) | 0.088 | -0.128 | Normal(0,1) | 6 |
| Retain_L_infl_Com_VL_1(1)_BLK1repl_199 | 24.54 | (5,80) | 0.436 | 0.018 | | 3 |
| Retain_L_width_Com_VL_1(1)_BLK1repl_1 | 1 | | | | | Fixed |
| Retain_L_infl_Com_LL_2(2)_BLK1repl_199 | 28.8 | | | | | Fixed |
| Retain_L_width_Com_LL_2(2)_BLK1repl_19 | 1 | | | | | Fixed |
| Retain_L_infl_Rec_PR_4(4)_BLK1repl_1990 | 25 | | | | | Fixed |
| Retain_L_width_Rec_PR_4(4)_BLK1repl_19 | 1 | | | | | Fixed |
| Retain_L_infl_Rec_Shore_5(5)_BLK1repl_19 | 25.88 | | | | | Fixed |
| Retain_L_width_Rec_Shore_5(5)_BLK1repl_1 | 1 | | | | | Fixed |

Table 10 Continued. List of Stock Synthesis parameters for Gulf of Mexico Gray Snapper. The list includes predicted parameter values, lower and upper bounds of the parameters, associated standard errors and coefficients of variation, the prior type and densities (value, SE) assigned to the parameters as applicable, and phases (negative identifies parameters that were fixed). Parameters designated as fixed were held at their initial values and have no associated range or SE.

| Label | Value | Range | SE | CV | Prior | Phase |
|--|-------|--------|-------|-------|-------|-------|
| Retain_L_infl_Rec_HB_CBT_6(6)_BLK1repl | 28.23 | (6,79) | 0.217 | 0.008 | | 3 |
| Retain_L_width_Rec_HB_CBT_6(6)_BLK1re | 1 | | | | | Fixed |

Table 11. Estimates of annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) combined across all fleets for Gulf of Mexico Gray Snapper, which was used as the proxy for annual fishing mortality rate. Estimates are provided for SEDAR75 and SEDAR51.

| Year | SEDAR75 | SEDAR51 |
|------|---------|---------|
| 1945 | 0.002 | 0.001 |
| 1946 | 0.003 | 0.005 |
| 1947 | 0.004 | 0.008 |
| 1948 | 0.005 | 0.012 |
| 1949 | 0.006 | 0.017 |
| 1950 | 0.007 | 0.020 |
| 1951 | 0.008 | 0.024 |
| 1952 | 0.008 | 0.028 |
| 1953 | 0.009 | 0.031 |
| 1954 | 0.010 | 0.036 |
| 1955 | 0.009 | 0.038 |
| 1956 | 0.010 | 0.041 |
| 1957 | 0.011 | 0.044 |
| 1958 | 0.012 | 0.048 |
| 1959 | 0.013 | 0.051 |
| 1960 | 0.014 | 0.052 |
| 1961 | 0.014 | 0.053 |
| 1962 | 0.015 | 0.054 |
| 1963 | 0.015 | 0.055 |
| 1964 | 0.016 | 0.059 |
| 1965 | 0.017 | 0.059 |
| 1966 | 0.017 | 0.062 |
| 1967 | 0.019 | 0.067 |
| 1968 | 0.021 | 0.069 |
| 1969 | 0.022 | 0.070 |
| 1970 | 0.023 | 0.074 |

Table 11 Continued. Estimates of annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) combined across all fleets for Gulf of Mexico Gray Snapper, which was used as the proxy for annual fishing mortality rate. Estimates are provided for SEDAR75 and SEDAR51.

| Year | SEDAR75 | SEDAR51 |
|------|---------|---------|
| 1971 | 0.026 | 0.087 |
| 1972 | 0.029 | 0.089 |
| 1973 | 0.031 | 0.097 |
| 1974 | 0.034 | 0.105 |
| 1975 | 0.035 | 0.124 |
| 1976 | 0.038 | 0.140 |
| 1977 | 0.040 | 0.165 |
| 1978 | 0.043 | 0.191 |
| 1979 | 0.046 | 0.189 |
| 1980 | 0.049 | 0.172 |
| 1981 | 0.069 | 0.266 |
| 1982 | 0.083 | 0.171 |
| 1983 | 0.053 | 0.296 |
| 1984 | 0.073 | 0.229 |
| 1985 | 0.042 | 0.158 |
| 1986 | 0.045 | 0.266 |
| 1987 | 0.059 | 0.349 |
| 1988 | 0.051 | 0.297 |
| 1989 | 0.082 | 0.160 |
| 1990 | 0.084 | 0.317 |
| 1991 | 0.133 | 0.168 |
| 1992 | 0.088 | 0.191 |
| 1993 | 0.097 | 0.197 |
| 1994 | 0.088 | 0.167 |
| 1995 | 0.101 | 0.141 |
| 1996 | 0.097 | 0.134 |
| 1997 | 0.084 | 0.155 |

Table 11 Continued. Estimates of annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) combined across all fleets for Gulf of Mexico Gray Snapper, which was used as the proxy for annual fishing mortality rate. Estimates are provided for SEDAR75 and SEDAR51.

| Year | SEDAR75 | SEDAR51 |
|------|---------|---------|
| 1998 | 0.104 | 0.116 |
| 1999 | 0.082 | 0.173 |
| 2000 | 0.075 | 0.165 |
| 2001 | 0.090 | 0.153 |
| 2002 | 0.076 | 0.182 |
| 2003 | 0.098 | 0.224 |
| 2004 | 0.114 | 0.180 |
| 2005 | 0.089 | 0.105 |
| 2006 | 0.073 | 0.154 |
| 2007 | 0.078 | 0.203 |
| 2008 | 0.102 | 0.219 |
| 2009 | 0.105 | 0.102 |
| 2010 | 0.055 | 0.088 |
| 2011 | 0.063 | 0.153 |
| 2012 | 0.115 | 0.136 |
| 2013 | 0.104 | 0.144 |
| 2014 | 0.118 | 0.135 |
| 2015 | 0.105 | 0.132 |
| 2016 | 0.117 | |
| 2017 | 0.098 | |
| 2018 | 0.093 | |
| 2019 | 0.091 | |
| 2020 | 0.088 | |

Table 12. Estimates of annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) by fleet for Gulf of Mexico Gray Snapper.

| Year | ComVL | ComLL | ComNT | Pri | Shr | Ch/Hbt | Total |
|------|-------|-------|-------|-------|-------|--------|-------|
| 1945 | 0.002 | 0 | 0.000 | 0.000 | 0.000 | 0.000 | 0.002 |
| 1946 | 0.002 | 0 | 0.000 | 0.001 | 0.000 | 0.000 | 0.003 |
| 1947 | 0.002 | 0 | 0.000 | 0.001 | 0.000 | 0.000 | 0.004 |
| 1948 | 0.002 | 0 | 0.000 | 0.002 | 0.000 | 0.001 | 0.005 |
| 1949 | 0.002 | 0 | 0.000 | 0.002 | 0.001 | 0.001 | 0.006 |
| 1950 | 0.002 | 0 | 0.000 | 0.003 | 0.001 | 0.001 | 0.007 |
| 1951 | 0.002 | 0 | 0.000 | 0.003 | 0.001 | 0.001 | 0.008 |
| 1952 | 0.002 | 0 | 0.000 | 0.004 | 0.001 | 0.001 | 0.008 |
| 1953 | 0.002 | 0 | 0.000 | 0.005 | 0.001 | 0.002 | 0.009 |
| 1954 | 0.002 | 0 | 0.000 | 0.005 | 0.001 | 0.002 | 0.010 |
| 1955 | 0.002 | 0 | 0.000 | 0.004 | 0.001 | 0.001 | 0.009 |
| 1956 | 0.002 | 0 | 0.000 | 0.005 | 0.001 | 0.002 | 0.010 |
| 1957 | 0.002 | 0 | 0.000 | 0.005 | 0.001 | 0.002 | 0.011 |
| 1958 | 0.003 | 0 | 0.000 | 0.006 | 0.001 | 0.002 | 0.012 |
| 1959 | 0.003 | 0 | 0.000 | 0.006 | 0.002 | 0.002 | 0.013 |
| 1960 | 0.003 | 0 | 0.000 | 0.007 | 0.002 | 0.002 | 0.014 |
| 1961 | 0.003 | 0 | 0.000 | 0.007 | 0.002 | 0.002 | 0.014 |
| 1962 | 0.003 | 0 | 0.000 | 0.008 | 0.002 | 0.003 | 0.015 |
| 1963 | 0.003 | 0 | 0.000 | 0.008 | 0.002 | 0.003 | 0.015 |
| 1964 | 0.003 | 0 | 0.000 | 0.008 | 0.002 | 0.003 | 0.016 |
| 1965 | 0.003 | 0 | 0.000 | 0.008 | 0.002 | 0.003 | 0.017 |
| 1966 | 0.003 | 0 | 0.000 | 0.009 | 0.002 | 0.003 | 0.017 |
| 1967 | 0.003 | 0 | 0.001 | 0.010 | 0.002 | 0.003 | 0.019 |
| 1968 | 0.004 | 0 | 0.001 | 0.010 | 0.002 | 0.003 | 0.021 |
| 1969 | 0.004 | 0 | 0.001 | 0.011 | 0.003 | 0.004 | 0.022 |
| 1970 | 0.004 | 0 | 0.001 | 0.012 | 0.003 | 0.004 | 0.023 |

Table 12 Continued. Estimates of annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) by fleet for Gulf of Mexico Gray Snapper.

| Year | ComVL | ComLL | ComNT | Pri | Shr | Ch/Hbt | Total |
|------|-------|-------|-------|-------|-------|--------|-------|
| 1971 | 0.004 | 0.000 | 0.001 | 0.013 | 0.003 | 0.004 | 0.026 |
| 1972 | 0.005 | 0.000 | 0.001 | 0.014 | 0.003 | 0.005 | 0.029 |
| 1973 | 0.005 | 0.000 | 0.002 | 0.015 | 0.004 | 0.005 | 0.031 |
| 1974 | 0.005 | 0.000 | 0.002 | 0.017 | 0.004 | 0.006 | 0.034 |
| 1975 | 0.003 | 0.000 | 0.003 | 0.018 | 0.005 | 0.006 | 0.035 |
| 1976 | 0.006 | 0.000 | 0.002 | 0.019 | 0.005 | 0.007 | 0.038 |
| 1977 | 0.004 | 0.000 | 0.005 | 0.020 | 0.005 | 0.007 | 0.040 |
| 1978 | 0.003 | 0.000 | 0.006 | 0.021 | 0.005 | 0.007 | 0.043 |
| 1979 | 0.004 | 0.000 | 0.006 | 0.022 | 0.006 | 0.008 | 0.046 |
| 1980 | 0.008 | 0.000 | 0.003 | 0.024 | 0.006 | 0.008 | 0.049 |
| 1981 | 0.009 | 0.000 | 0.003 | 0.042 | 0.009 | 0.007 | 0.069 |
| 1982 | 0.012 | 0.001 | 0.003 | 0.032 | 0.009 | 0.025 | 0.083 |
| 1983 | 0.013 | 0.002 | 0.003 | 0.008 | 0.009 | 0.018 | 0.053 |
| 1984 | 0.010 | 0.001 | 0.005 | 0.043 | 0.009 | 0.004 | 0.073 |
| 1985 | 0.008 | 0.001 | 0.005 | 0.019 | 0.005 | 0.005 | 0.042 |
| 1986 | 0.009 | 0.001 | 0.004 | 0.016 | 0.008 | 0.007 | 0.045 |
| 1987 | 0.010 | 0.001 | 0.003 | 0.033 | 0.008 | 0.004 | 0.059 |
| 1988 | 0.007 | 0.001 | 0.002 | 0.028 | 0.011 | 0.003 | 0.051 |
| 1989 | 0.008 | 0.001 | 0.003 | 0.046 | 0.018 | 0.006 | 0.082 |
| 1990 | 0.007 | 0.001 | 0.001 | 0.057 | 0.014 | 0.005 | 0.084 |
| 1991 | 0.009 | 0.001 | 0.001 | 0.058 | 0.051 | 0.013 | 0.133 |
| 1992 | 0.009 | 0.001 | 0.001 | 0.046 | 0.020 | 0.011 | 0.088 |
| 1993 | 0.011 | 0.002 | 0.000 | 0.056 | 0.021 | 0.006 | 0.097 |
| 1994 | 0.014 | 0.000 | 0.001 | 0.048 | 0.016 | 0.009 | 0.088 |
| 1995 | 0.012 | 0.000 | 0.001 | 0.066 | 0.016 | 0.006 | 0.101 |
| 1996 | 0.011 | 0.000 | 0.000 | 0.053 | 0.026 | 0.007 | 0.097 |
| 1997 | 0.011 | 0.000 | 0.000 | 0.054 | 0.013 | 0.005 | 0.084 |

Table 12 Continued. Estimates of annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) by fleet for Gulf of Mexico Gray Snapper.

| Year | ComVL | ComLL | ComNT | Pri | Shr | Ch/Hbt | Total |
|------|-------|-------|-------|-------|-------|--------|-------|
| 1998 | 0.009 | 0.000 | 0 | 0.070 | 0.012 | 0.012 | 0.104 |
| 1999 | 0.008 | 0.000 | 0 | 0.054 | 0.009 | 0.010 | 0.082 |
| 2000 | 0.008 | 0.000 | 0 | 0.047 | 0.013 | 0.007 | 0.075 |
| 2001 | 0.009 | 0.000 | 0 | 0.054 | 0.016 | 0.011 | 0.090 |
| 2002 | 0.010 | 0.000 | 0 | 0.048 | 0.009 | 0.009 | 0.076 |
| 2003 | 0.009 | 0.000 | 0 | 0.064 | 0.012 | 0.013 | 0.098 |
| 2004 | 0.009 | 0.000 | 0 | 0.080 | 0.014 | 0.011 | 0.114 |
| 2005 | 0.008 | 0.000 | 0 | 0.055 | 0.014 | 0.011 | 0.089 |
| 2006 | 0.007 | 0.000 | 0 | 0.049 | 0.007 | 0.010 | 0.073 |
| 2007 | 0.005 | 0.000 | 0 | 0.053 | 0.010 | 0.010 | 0.078 |
| 2008 | 0.005 | 0.000 | 0 | 0.070 | 0.016 | 0.011 | 0.102 |
| 2009 | 0.006 | 0.000 | 0 | 0.073 | 0.011 | 0.014 | 0.105 |
| 2010 | 0.005 | 0.000 | 0 | 0.039 | 0.003 | 0.007 | 0.055 |
| 2011 | 0.006 | 0.000 | 0 | 0.040 | 0.009 | 0.009 | 0.063 |
| 2012 | 0.006 | 0.000 | 0 | 0.087 | 0.014 | 0.008 | 0.115 |
| 2013 | 0.005 | 0.000 | 0 | 0.071 | 0.013 | 0.014 | 0.104 |
| 2014 | 0.006 | 0.000 | 0 | 0.078 | 0.019 | 0.015 | 0.118 |
| 2015 | 0.005 | 0.001 | 0 | 0.073 | 0.014 | 0.012 | 0.105 |
| 2016 | 0.005 | 0.001 | 0 | 0.074 | 0.021 | 0.017 | 0.117 |
| 2017 | 0.004 | 0.000 | 0 | 0.052 | 0.026 | 0.015 | 0.098 |
| 2018 | 0.004 | 0.000 | 0 | 0.052 | 0.022 | 0.014 | 0.093 |
| 2019 | 0.003 | 0.000 | 0 | 0.044 | 0.028 | 0.016 | 0.091 |
| 2020 | 0.003 | 0.000 | 0 | 0.054 | 0.020 | 0.011 | 0.088 |

Table 13. Expected biomass (metric tons) for all Gray Snapper and exploited Gray Snapper (2+ years), spawning stock biomass (SSB, metric tons), exploited numbers (2+years, 1,000s of fish), age-0 recruits (1,000s of fish), and SSB ratio (SSB/SSB₀) where SSB₀ = 21,718 metric tons for Gulf of Mexico Gray Snapper.

| Year | Biomass (all) | Biomass (exploited) | SSB | Abundance (exploited) | Recruits | SSB ratio |
|------|------------------|------------------------|--------|--------------------------|----------|--------------|
| 1945 | 56,913 | 55,844 | 21,718 | 54,482 | 23,190 | 1.00 |
| 1946 | 56,780 | 55,711 | 21,669 | 54,407 | 23,190 | 1.00 |
| 1947 | 56,584 | 55,514 | 21,595 | 54,247 | 23,190 | 0.99 |
| 1948 | 56,325 | 55,256 | 21,496 | 54,019 | 23,190 | 0.99 |
| 1949 | 56,005 | 54,935 | 21,373 | 53,735 | 23,189 | 0.98 |
| 1950 | 55,625 | 54,555 | 21,227 | 53,404 | 23,189 | 0.98 |
| 1951 | 55,187 | 54,118 | 21,059 | 53,035 | 23,532 | 0.97 |
| 1952 | 54,712 | 53,627 | 20,871 | 52,631 | 23,569 | 0.96 |
| 1953 | 54,196 | 53,109 | 20,676 | 52,345 | 23,614 | 0.95 |
| 1954 | 53,644 | 52,555 | 20,469 | 52,014 | 23,671 | 0.94 |
| 1955 | 53,060 | 51,968 | 20,251 | 51,654 | 23,695 | 0.93 |
| 1956 | 52,604 | 51,512 | 20,086 | 51,500 | 23,538 | 0.92 |
| 1957 | 52,125 | 51,040 | 19,917 | 51,291 | 23,394 | 0.92 |
| 1958 | 51,618 | 50,539 | 19,734 | 50,964 | 23,813 | 0.91 |
| 1959 | 51,104 | 50,006 | 19,538 | 50,555 | 23,481 | 0.90 |
| 1960 | 50,560 | 49,478 | 19,346 | 50,316 | 22,383 | 0.89 |
| 1961 | 49,943 | 48,911 | 19,137 | 49,885 | 22,644 | 0.88 |
| 1962 | 49,315 | 48,270 | 18,889 | 49,028 | 24,986 | 0.87 |
| 1963 | 48,773 | 47,621 | 18,640 | 48,418 | 24,078 | 0.86 |
| 1964 | 48,278 | 47,170 | 18,489 | 48,886 | 20,132 | 0.85 |
| 1965 | 47,624 | 46,696 | 18,324 | 48,768 | 19,528 | 0.84 |
| 1966 | 46,826 | 45,926 | 18,009 | 46,928 | 19,007 | 0.83 |
| 1967 | 45,948 | 45,072 | 17,653 | 45,249 | 19,106 | 0.81 |
| 1968 | 44,953 | 44,073 | 17,236 | 43,640 | 17,140 | 0.79 |
| 1969 | 43,756 | 42,966 | 16,784 | 42,316 | 16,688 | 0.77 |
| 1970 | 42,447 | 41,677 | 16,249 | 40,346 | 17,377 | 0.75 |
| 1971 | 41,118 | 40,316 | 15,689 | 38,552 | 19,211 | 0.72 |

Table 13 Continued. Expected biomass (metric tons) for all Gray Snapper and exploited Gray Snapper (2+ years), spawning stock biomass (SSB, metric tons), exploited numbers (2+years, 1,000s of fish), age-0 recruits (1,000s of fish), and SSB ratio (SSB/SSB₀) where SSB₀ = 21,718 metric tons for Gulf of Mexico Gray Snapper.

| Year | Biomass (all) | Biomass (exploited) | SSB | Abundance (exploited) | Recruits | SSB ratio |
|------|------------------|------------------------|--------|--------------------------|----------|--------------|
| 1972 | 39,764 | 38,879 | 15,112 | 37,242 | 20,902 | 0.70 |
| 1973 | 38,450 | 37,487 | 14,578 | 36,757 | 21,080 | 0.67 |
| 1974 | 37,198 | 36,227 | 14,119 | 36,846 | 21,347 | 0.65 |
| 1975 | 36,011 | 35,029 | 13,692 | 36,752 | 18,841 | 0.63 |
| 1976 | 34,836 | 33,969 | 13,321 | 36,619 | 17,844 | 0.61 |
| 1977 | 33,594 | 32,772 | 12,879 | 35,384 | 16,040 | 0.59 |
| 1978 | 32,282 | 31,543 | 12,410 | 33,927 | 14,948 | 0.57 |
| 1979 | 30,862 | 30,174 | 11,870 | 31,964 | 13,069 | 0.55 |
| 1980 | 29,300 | 28,698 | 11,281 | 29,935 | 14,851 | 0.52 |
| 1981 | 27,713 | 27,026 | 10,608 | 27,531 | 22,324 | 0.49 |
| 1982 | 25,815 | 24,790 | 9,724 | 25,325 | 17,458 | 0.45 |
| 1983 | 23,826 | 23,022 | 9,102 | 26,515 | 19,542 | 0.42 |
| 1984 | 23,197 | 22,297 | 8,868 | 26,740 | 19,009 | 0.41 |
| 1985 | 22,066 | 21,190 | 8,478 | 26,413 | 22,706 | 0.39 |
| 1986 | 22,205 | 21,163 | 8,529 | 27,605 | 12,743 | 0.39 |
| 1987 | 22,034 | 21,446 | 8,723 | 29,806 | 15,682 | 0.40 |
| 1988 | 21,473 | 20,746 | 8,431 | 26,720 | 29,132 | 0.39 |
| 1989 | 21,742 | 20,408 | 8,291 | 25,809 | 9,880 | 0.38 |
| 1990 | 20,638 | 20,178 | 8,272 | 28,819 | 25,334 | 0.38 |
| 1991 | 19,994 | 18,827 | 7,697 | 24,499 | 25,948 | 0.35 |
| 1992 | 18,584 | 17,395 | 7,158 | 25,569 | 18,182 | 0.33 |
| 1993 | 18,270 | 17,435 | 7,273 | 28,626 | 12,703 | 0.34 |
| 1994 | 17,597 | 17,011 | 7,131 | 27,355 | 15,460 | 0.33 |
| 1995 | 17,186 | 16,475 | 6,897 | 24,709 | 13,967 | 0.32 |
| 1996 | 16,438 | 15,793 | 6,612 | 23,586 | 21,179 | 0.30 |
| 1997 | 16,114 | 15,138 | 6,324 | 22,122 | 25,813 | 0.29 |
| 1998 | 16,426 | 15,241 | 6,415 | 24,546 | 14,591 | 0.30 |

Table 13 Continued. Expected biomass (metric tons) for all Gray Snapper and exploited Gray Snapper (2+ years), spawning stock biomass (SSB, metric tons), exploited numbers (2+years, 1,000s of fish), age-0 recruits (1,000s of fish), and SSB ratio (SSB/SSB₀) where SSB₀ = 21,718 metric tons for Gulf of Mexico Gray Snapper.

| Year | Biomass (all) | Biomass (exploited) | SSB | Abundance (exploited) | Recruits | SSB ratio |
|------|------------------|------------------------|--------|--------------------------|----------|--------------|
| 1999 | 16,125 | 15,452 | 6,582 | 27,689 | 18,392 | 0.30 |
| 2000 | 16,365 | 15,521 | 6,609 | 25,922 | 11,582 | 0.30 |
| 2001 | 16,430 | 15,892 | 6,773 | 26,419 | 23,387 | 0.31 |
| 2002 | 16,566 | 15,490 | 6,561 | 23,571 | 20,163 | 0.30 |
| 2003 | 16,972 | 16,042 | 6,834 | 26,928 | 22,897 | 0.32 |
| 2004 | 17,116 | 16,062 | 6,859 | 27,292 | 22,416 | 0.32 |
| 2005 | 17,024 | 15,992 | 6,856 | 28,146 | 23,589 | 0.32 |
| 2006 | 17,547 | 16,458 | 7,075 | 29,274 | 30,795 | 0.33 |
| 2007 | 18,738 | 17,324 | 7,470 | 31,180 | 18,304 | 0.34 |
| 2008 | 19,435 | 18,593 | 8,066 | 35,358 | 16,840 | 0.37 |
| 2009 | 19,323 | 18,547 | 8,004 | 32,217 | 18,280 | 0.37 |
| 2010 | 19,020 | 18,176 | 7,794 | 29,499 | 23,547 | 0.36 |
| 2011 | 19,951 | 18,861 | 8,053 | 29,829 | 34,497 | 0.37 |
| 2012 | 21,195 | 19,606 | 8,366 | 31,899 | 36,054 | 0.38 |
| 2013 | 21,505 | 19,854 | 8,532 | 35,921 | 15,718 | 0.39 |
| 2014 | 21,448 | 20,718 | 8,973 | 39,850 | 32,658 | 0.41 |
| 2015 | 21,482 | 19,982 | 8,587 | 33,622 | 23,169 | 0.40 |
| 2016 | 21,568 | 20,497 | 8,840 | 36,890 | 34,747 | 0.41 |
| 2017 | 21,747 | 20,149 | 8,663 | 34,716 | 32,587 | 0.40 |
| 2018 | 22,482 | 20,976 | 9,055 | 38,496 | 46,234 | 0.42 |
| 2019 | 23,986 | 21,859 | 9,458 | 40,514 | 44,247 | 0.44 |
| 2020 | 25,746 | 23,714 | 10,345 | 47,564 | 31,341 | 0.48 |

Table 14. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Vertical Line fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1944 | 0.05 | 0.00 | 0.00 | 0 | |
| 1945 | 0.05 | 0.22 | 0.22 | 73 | 3.0 |
| 1946 | 0.05 | 0.22 | 0.22 | 74 | 3.0 |
| 1947 | 0.05 | 0.23 | 0.23 | 76 | 3.0 |
| 1948 | 0.05 | 0.23 | 0.23 | 78 | 3.0 |
| 1949 | 0.05 | 0.24 | 0.24 | 79 | 3.0 |
| 1950 | 0.05 | 0.24 | 0.24 | 81 | 3.0 |
| 1951 | 0.05 | 0.25 | 0.25 | 83 | 3.0 |
| 1952 | 0.05 | 0.25 | 0.25 | 84 | 3.0 |
| 1953 | 0.05 | 0.26 | 0.26 | 86 | 3.0 |
| 1954 | 0.05 | 0.26 | 0.26 | 88 | 3.0 |
| 1955 | 0.05 | 0.27 | 0.27 | 89 | 3.0 |
| 1956 | 0.05 | 0.27 | 0.27 | 91 | 3.0 |
| 1957 | 0.05 | 0.28 | 0.28 | 93 | 3.0 |
| 1958 | 0.05 | 0.28 | 0.28 | 95 | 3.0 |
| 1959 | 0.05 | 0.29 | 0.29 | 96 | 3.0 |
| 1960 | 0.05 | 0.29 | 0.29 | 98 | 3.0 |
| 1961 | 0.05 | 0.30 | 0.30 | 100 | 3.0 |
| 1962 | 0.05 | 0.32 | 0.32 | 109 | 2.9 |
| 1963 | 0.05 | 0.28 | 0.28 | 95 | 2.9 |
| 1964 | 0.05 | 0.30 | 0.30 | 101 | 2.9 |
| 1965 | 0.05 | 0.33 | 0.33 | 112 | 2.9 |
| 1966 | 0.05 | 0.28 | 0.28 | 95 | 2.9 |
| 1967 | 0.05 | 0.32 | 0.32 | 109 | 2.9 |
| 1968 | 0.05 | 0.40 | 0.40 | 136 | 2.9 |
| 1969 | 0.05 | 0.37 | 0.37 | 126 | 3.0 |
| 1970 | 0.05 | 0.34 | 0.34 | 113 | 3.0 |

Table 14 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Vertical Line fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1971 | 0.05 | 0.36 | 0.36 | 120 | 3.0 |
| 1972 | 0.05 | 0.40 | 0.40 | 134 | 3.0 |
| 1973 | 0.05 | 0.39 | 0.39 | 128 | 3.0 |
| 1974 | 0.05 | 0.39 | 0.39 | 130 | 3.0 |
| 1975 | 0.05 | 0.26 | 0.26 | 85 | 3.0 |
| 1976 | 0.05 | 0.47 | 0.47 | 158 | 3.0 |
| 1977 | 0.05 | 0.27 | 0.27 | 91 | 3.0 |
| 1978 | 0.05 | 0.23 | 0.23 | 79 | 2.9 |
| 1979 | 0.05 | 0.27 | 0.28 | 94 | 2.9 |
| 1980 | 0.05 | 0.51 | 0.51 | 174 | 2.9 |
| 1981 | 0.05 | 0.51 | 0.51 | 174 | 2.9 |
| 1982 | 0.05 | 0.68 | 0.68 | 233 | 2.9 |
| 1983 | 0.05 | 0.67 | 0.67 | 233 | 2.9 |
| 1984 | 0.05 | 0.47 | 0.47 | 166 | 2.8 |
| 1985 | 0.05 | 0.37 | 0.37 | 134 | 2.8 |
| 1986 | 0.05 | 0.41 | 0.41 | 150 | 2.7 |
| 1987 | 0.05 | 0.50 | 0.50 | 186 | 2.7 |
| 1988 | 0.05 | 0.32 | 0.32 | 121 | 2.6 |
| 1989 | 0.05 | 0.38 | 0.38 | 147 | 2.6 |
| 1990 | 0.05 | 0.31 | 0.31 | 116 | 2.6 |
| 1991 | 0.05 | 0.38 | 0.38 | 145 | 2.6 |
| 1992 | 0.05 | 0.35 | 0.35 | 135 | 2.6 |
| 1993 | 0.05 | 0.43 | 0.43 | 167 | 2.6 |
| 1994 | 0.05 | 0.52 | 0.52 | 210 | 2.5 |
| 1995 | 0.05 | 0.41 | 0.42 | 172 | 2.4 |
| 1996 | 0.05 | 0.40 | 0.40 | 164 | 2.4 |
| 1997 | 0.05 | 0.38 | 0.38 | 157 | 2.4 |

Table 14 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Vertical Line fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1998 | 0.05 | 0.29 | 0.30 | 122 | 2.4 |
| 1999 | 0.05 | 0.26 | 0.27 | 111 | 2.4 |
| 2000 | 0.05 | 0.28 | 0.28 | 121 | 2.3 |
| 2001 | 0.05 | 0.30 | 0.30 | 130 | 2.3 |
| 2002 | 0.05 | 0.35 | 0.36 | 154 | 2.3 |
| 2003 | 0.05 | 0.31 | 0.31 | 135 | 2.3 |
| 2004 | 0.05 | 0.32 | 0.33 | 141 | 2.3 |
| 2005 | 0.05 | 0.29 | 0.29 | 127 | 2.3 |
| 2006 | 0.05 | 0.26 | 0.26 | 114 | 2.3 |
| 2007 | 0.05 | 0.20 | 0.20 | 88 | 2.2 |
| 2008 | 0.05 | 0.20 | 0.21 | 93 | 2.2 |
| 2009 | 0.05 | 0.25 | 0.25 | 115 | 2.2 |
| 2010 | 0.05 | 0.21 | 0.21 | 94 | 2.2 |
| 2011 | 0.05 | 0.24 | 0.23 | 104 | 2.2 |
| 2012 | 0.05 | 0.25 | 0.24 | 106 | 2.3 |
| 2013 | 0.05 | 0.22 | 0.22 | 94 | 2.3 |
| 2014 | 0.05 | 0.28 | 0.27 | 119 | 2.3 |
| 2015 | 0.05 | 0.24 | 0.24 | 105 | 2.2 |
| 2016 | 0.05 | 0.24 | 0.24 | 106 | 2.2 |
| 2017 | 0.05 | 0.18 | 0.18 | 81 | 2.3 |
| 2018 | 0.05 | 0.20 | 0.20 | 86 | 2.3 |
| 2019 | 0.05 | 0.16 | 0.16 | 72 | 2.3 |
| 2020 | 0.05 | 0.13 | 0.13 | 60 | 2.2 |

Table 15. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Longline fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1944 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1945 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1946 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1947 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1948 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1949 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1950 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1951 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1952 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1953 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1954 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1955 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1956 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1957 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1958 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1959 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1960 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1961 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1962 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1963 | 0.05 | 0.001 | 0.001 | 0.13 | 3.9 |
| 1964 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1965 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1966 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1967 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1968 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1969 | 0.05 | 0.000 | 0.000 | 0.00 | |
| 1970 | 0.05 | 0.000 | 0.000 | 0.00 | |

Table 15 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Longline fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1971 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1972 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1973 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1974 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1975 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1976 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1977 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1978 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1979 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1980 | 0.05 | 0.020 | 0.020 | 5.0 | 3.9 |
| 1981 | 0.05 | 0.020 | 0.020 | 5.1 | 3.9 |
| 1982 | 0.05 | 0.047 | 0.047 | 12.1 | 3.9 |
| 1983 | 0.05 | 0.086 | 0.086 | 22.0 | 3.9 |
| 1984 | 0.05 | 0.047 | 0.047 | 12.1 | 3.9 |
| 1985 | 0.05 | 0.035 | 0.035 | 8.9 | 3.9 |
| 1986 | 0.05 | 0.041 | 0.041 | 10.6 | 3.9 |
| 1987 | 0.05 | 0.046 | 0.046 | 11.8 | 3.9 |
| 1988 | 0.05 | 0.031 | 0.031 | 8.2 | 3.9 |
| 1989 | 0.05 | 0.046 | 0.046 | 12.1 | 3.8 |
| 1990 | 0.05 | 0.041 | 0.041 | 10.6 | 3.9 |
| 1991 | 0.05 | 0.043 | 0.043 | 11.3 | 3.8 |
| 1992 | 0.05 | 0.050 | 0.050 | 13.1 | 3.8 |
| 1993 | 0.05 | 0.079 | 0.068 | 17.9 | 3.8 |
| 1994 | 0.05 | 0.009 | 0.009 | 2.3 | 3.8 |
| 1995 | 0.05 | 0.009 | 0.009 | 2.4 | 3.8 |
| 1996 | 0.05 | 0.007 | 0.007 | 1.8 | 3.8 |
| 1997 | 0.05 | 0.007 | 0.008 | 2.0 | 3.8 |

Table 15 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Longline fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1998 | 0.05 | 0.008 | 0.008 | 2.2 | 3.7 |
| 1999 | 0.05 | 0.013 | 0.013 | 3.5 | 3.7 |
| 2000 | 0.05 | 0.013 | 0.013 | 3.4 | 3.7 |
| 2001 | 0.05 | 0.011 | 0.011 | 3.0 | 3.7 |
| 2002 | 0.05 | 0.014 | 0.014 | 3.8 | 3.7 |
| 2003 | 0.05 | 0.010 | 0.010 | 2.7 | 3.7 |
| 2004 | 0.05 | 0.015 | 0.015 | 4.1 | 3.7 |
| 2005 | 0.05 | 0.013 | 0.013 | 3.6 | 3.7 |
| 2006 | 0.05 | 0.013 | 0.013 | 3.5 | 3.7 |
| 2007 | 0.05 | 0.012 | 0.012 | 3.3 | 3.7 |
| 2008 | 0.05 | 0.015 | 0.015 | 4.1 | 3.6 |
| 2009 | 0.05 | 0.017 | 0.017 | 4.5 | 3.6 |
| 2010 | 0.05 | 0.007 | 0.007 | 1.9 | 3.6 |
| 2011 | 0.05 | 0.013 | 0.013 | 3.5 | 3.6 |
| 2012 | 0.05 | 0.013 | 0.013 | 3.7 | 3.6 |
| 2013 | 0.05 | 0.012 | 0.013 | 3.5 | 3.6 |
| 2014 | 0.05 | 0.018 | 0.018 | 5.0 | 3.6 |
| 2015 | 0.05 | 0.026 | 0.026 | 7.2 | 3.6 |
| 2016 | 0.05 | 0.025 | 0.026 | 7.0 | 3.7 |
| 2017 | 0.05 | 0.021 | 0.021 | 5.8 | 3.7 |
| 2018 | 0.05 | 0.012 | 0.012 | 3.4 | 3.7 |
| 2019 | 0.05 | 0.014 | 0.014 | 3.8 | 3.7 |
| 2020 | 0.05 | 0.011 | 0.011 | 3.0 | 3.7 |

Table 16. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Nets & Traps fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1944 | 0.05 | 0.000 | 0.000 | 0.0 | |
| 1945 | 0.05 | 0.017 | 0.017 | 8.8 | 1.9 |
| 1946 | 0.05 | 0.017 | 0.017 | 9.0 | 1.9 |
| 1947 | 0.05 | 0.018 | 0.018 | 9.2 | 1.9 |
| 1948 | 0.05 | 0.018 | 0.018 | 9.4 | 1.9 |
| 1949 | 0.05 | 0.018 | 0.018 | 9.6 | 1.9 |
| 1950 | 0.05 | 0.019 | 0.019 | 9.8 | 1.9 |
| 1951 | 0.05 | 0.019 | 0.019 | 10.0 | 1.9 |
| 1952 | 0.05 | 0.019 | 0.019 | 10.2 | 1.9 |
| 1953 | 0.05 | 0.020 | 0.020 | 10.4 | 1.9 |
| 1954 | 0.05 | 0.020 | 0.020 | 10.6 | 1.9 |
| 1955 | 0.05 | 0.021 | 0.021 | 10.8 | 1.9 |
| 1956 | 0.05 | 0.021 | 0.021 | 11.1 | 1.9 |
| 1957 | 0.05 | 0.021 | 0.021 | 11.3 | 1.9 |
| 1958 | 0.05 | 0.022 | 0.022 | 11.5 | 1.9 |
| 1959 | 0.05 | 0.022 | 0.022 | 11.7 | 1.9 |
| 1960 | 0.05 | 0.022 | 0.022 | 11.9 | 1.9 |
| 1961 | 0.05 | 0.023 | 0.023 | 12.1 | 1.9 |
| 1962 | 0.05 | 0.015 | 0.015 | 8.0 | 1.9 |
| 1963 | 0.05 | 0.009 | 0.009 | 4.7 | 1.9 |
| 1964 | 0.05 | 0.014 | 0.014 | 7.3 | 1.9 |
| 1965 | 0.05 | 0.047 | 0.047 | 25.3 | 1.9 |
| 1966 | 0.05 | 0.031 | 0.031 | 16.5 | 1.9 |
| 1967 | 0.05 | 0.054 | 0.054 | 28.7 | 1.9 |
| 1968 | 0.05 | 0.072 | 0.072 | 38.0 | 1.9 |
| 1969 | 0.05 | 0.107 | 0.107 | 56.1 | 1.9 |
| 1970 | 0.05 | 0.104 | 0.104 | 54.4 | 1.9 |

Table 16 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Nets & Traps fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1971 | 0.05 | 0.108 | 0.108 | 56.2 | 1.9 |
| 1972 | 0.05 | 0.127 | 0.127 | 65.4 | 1.9 |
| 1973 | 0.05 | 0.169 | 0.169 | 87.8 | 1.9 |
| 1974 | 0.05 | 0.194 | 0.194 | 102.0 | 1.9 |
| 1975 | 0.05 | 0.228 | 0.228 | 121.5 | 1.9 |
| 1976 | 0.05 | 0.125 | 0.125 | 67.7 | 1.9 |
| 1977 | 0.05 | 0.357 | 0.357 | 194.1 | 1.8 |
| 1978 | 0.05 | 0.433 | 0.433 | 235.4 | 1.8 |
| 1979 | 0.05 | 0.399 | 0.399 | 215.5 | 1.9 |
| 1980 | 0.05 | 0.178 | 0.178 | 95.5 | 1.9 |
| 1981 | 0.05 | 0.162 | 0.162 | 85.7 | 1.9 |
| 1982 | 0.05 | 0.174 | 0.174 | 91.6 | 1.9 |
| 1983 | 0.05 | 0.176 | 0.176 | 94.3 | 1.9 |
| 1984 | 0.05 | 0.268 | 0.268 | 148.2 | 1.8 |
| 1985 | 0.05 | 0.218 | 0.218 | 123.5 | 1.8 |
| 1986 | 0.05 | 0.169 | 0.169 | 97.4 | 1.7 |
| 1987 | 0.05 | 0.137 | 0.137 | 79.8 | 1.7 |
| 1988 | 0.05 | 0.098 | 0.098 | 56.9 | 1.7 |
| 1989 | 0.05 | 0.120 | 0.120 | 68.2 | 1.8 |
| 1990 | 0.05 | 0.044 | 0.044 | 24.8 | 1.8 |
| 1991 | 0.05 | 0.044 | 0.044 | 25.2 | 1.7 |
| 1992 | 0.05 | 0.022 | 0.022 | 12.8 | 1.7 |
| 1993 | 0.05 | 0.013 | 0.013 | 7.8 | 1.7 |
| 1994 | 0.05 | 0.035 | 0.035 | 21.7 | 1.6 |
| 1995 | 0.05 | 0.018 | 0.018 | 11.2 | 1.6 |
| 1996 | 0.05 | 0.009 | 0.009 | 5.3 | 1.7 |
| 1997 | 0.05 | 0.010 | 0.010 | 5.8 | 1.7 |

Table 16 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Commercial Nets & Traps fleet in weight (B, million pounds whole weight) and number (1,000s of fish) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input B SE | Input B | Exp B | Exp N | MW |
|------|------------|---------|-------|-------|-----|
| 1998 | 0.05 | 0.011 | 0.011 | 6.228 | 1.7 |
| 1999 | 0.05 | 0.008 | 0.008 | 4.655 | 1.6 |
| 2000 | 0.05 | 0.005 | 0.005 | 3.410 | 1.6 |
| 2001 | 0.05 | 0.005 | 0.005 | 2.925 | 1.6 |
| 2002 | 0.05 | 0.004 | 0.004 | 2.259 | 1.7 |
| 2003 | 0.05 | 0.005 | 0.005 | 2.859 | 1.7 |
| 2004 | 0.05 | 0.002 | 0.002 | 1.147 | 1.7 |
| 2005 | 0.05 | 0.002 | 0.002 | 0.934 | 1.6 |
| 2006 | 0.05 | 0.001 | 0.001 | 0.452 | 1.6 |
| 2007 | 0.05 | 0.000 | 0.000 | 0.261 | 1.6 |
| 2008 | 0.05 | 0.001 | 0.001 | 0.607 | 1.6 |
| 2009 | 0.05 | 0.001 | 0.001 | 0.549 | 1.6 |
| 2010 | 0.05 | 0.000 | 0.000 | 0.054 | 1.6 |
| 2011 | 0.05 | 0.000 | 0.000 | 0.053 | 1.7 |
| 2012 | 0.05 | 0.000 | 0.000 | 0.222 | 1.7 |
| 2013 | 0.05 | 0.001 | 0.001 | 0.812 | 1.7 |
| 2014 | 0.05 | 0.001 | 0.001 | 0.358 | 1.6 |
| 2015 | 0.05 | 0.000 | 0.000 | 0.138 | 1.6 |
| 2016 | 0.05 | 0.000 | 0.000 | 0.109 | 1.6 |
| 2017 | 0.05 | 0.000 | 0.000 | 0.108 | 1.6 |
| 2018 | 0.05 | 0.000 | 0.000 | 0.054 | 1.6 |
| 2019 | 0.05 | 0.000 | 0.000 | 0.151 | 1.6 |
| 2020 | 0.05 | 0.000 | 0.000 | 0.238 | 1.6 |

Table 17. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Private fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1945 | 0.1 | 0 | 0 | 0.000 | |
| 1946 | 0.1 | 57 | 57 | 0.067 | 1.2 |
| 1947 | 0.1 | 115 | 115 | 0.133 | 1.2 |
| 1948 | 0.1 | 172 | 172 | 0.200 | 1.2 |
| 1949 | 0.1 | 229 | 229 | 0.266 | 1.2 |
| 1950 | 0.1 | 286 | 286 | 0.332 | 1.2 |
| 1951 | 0.1 | 344 | 344 | 0.398 | 1.2 |
| 1952 | 0.1 | 401 | 401 | 0.462 | 1.2 |
| 1953 | 0.1 | 458 | 458 | 0.526 | 1.1 |
| 1954 | 0.1 | 516 | 516 | 0.589 | 1.1 |
| 1955 | 0.1 | 436 | 436 | 0.496 | 1.1 |
| 1956 | 0.1 | 483 | 483 | 0.546 | 1.1 |
| 1957 | 0.1 | 530 | 529 | 0.597 | 1.1 |
| 1958 | 0.1 | 576 | 576 | 0.649 | 1.1 |
| 1959 | 0.1 | 623 | 623 | 0.698 | 1.1 |
| 1960 | 0.1 | 669 | 669 | 0.748 | 1.1 |
| 1961 | 0.1 | 692 | 691 | 0.776 | 1.1 |
| 1962 | 0.1 | 714 | 714 | 0.801 | 1.1 |
| 1963 | 0.1 | 736 | 736 | 0.813 | 1.1 |
| 1964 | 0.1 | 759 | 758 | 0.828 | 1.1 |
| 1965 | 0.1 | 781 | 781 | 0.868 | 1.1 |
| 1966 | 0.1 | 805 | 805 | 0.914 | 1.1 |
| 1967 | 0.1 | 829 | 829 | 0.955 | 1.2 |
| 1968 | 0.1 | 853 | 853 | 0.987 | 1.2 |
| 1969 | 0.1 | 877 | 878 | 1.026 | 1.2 |
| 1970 | 0.1 | 901 | 902 | 1.065 | 1.2 |
| 1971 | 0.1 | 984 | 987 | 1.160 | 1.2 |

Table 17 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Private fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1972 | 0.1 | 1,068 | 1,071 | 1.23 | 1.1 |
| 1973 | 0.1 | 1,151 | 1,154 | 1.27 | 1.1 |
| 1974 | 0.1 | 1,235 | 1,238 | 1.32 | 1.1 |
| 1975 | 0.1 | 1,318 | 1,321 | 1.38 | 1.0 |
| 1976 | 0.1 | 1,324 | 1,326 | 1.39 | 1.0 |
| 1977 | 0.1 | 1,329 | 1,332 | 1.42 | 1.1 |
| 1978 | 0.1 | 1,334 | 1,339 | 1.45 | 1.1 |
| 1979 | 0.1 | 1,340 | 1,346 | 1.48 | 1.1 |
| 1980 | 0.1 | 1,345 | 1,354 | 1.52 | 1.1 |
| 1981 | 0.1 | 2,333 | 2,217 | 2.48 | 1.1 |
| 1982 | 0.1 | 1,836 | 1,726 | 1.73 | 1.0 |
| 1983 | 0.1 | 535 | 432 | 0.41 | 1.0 |
| 1984 | 0.1 | 2,579 | 2,285 | 2.10 | 0.9 |
| 1985 | 0.1 | 975 | 998 | 0.90 | 0.9 |
| 1986 | 0.1 | 843 | 862 | 0.74 | 0.9 |
| 1987 | 0.1 | 1,772 | 1,676 | 1.53 | 0.9 |
| 1988 | 0.1 | 1,327 | 1,343 | 1.28 | 1.0 |
| 1989 | 0.1 | 2,421 | 2,370 | 2.03 | 0.9 |
| 1990 | 0.1 | 1,953 | 1,861 | 2.48 | 1.3 |
| 1991 | 0.1 | 1,750 | 1,760 | 2.35 | 1.3 |
| 1992 | 0.1 | 1,318 | 1,338 | 1.69 | 1.3 |
| 1993 | 0.1 | 1,731 | 1,751 | 2.08 | 1.2 |
| 1994 | 0.1 | 1,425 | 1,470 | 1.75 | 1.2 |
| 1995 | 0.1 | 1,803 | 1,897 | 2.35 | 1.2 |
| 1996 | 0.1 | 1,323 | 1,422 | 1.80 | 1.3 |
| 1997 | 0.1 | 1,309 | 1,398 | 1.76 | 1.3 |
| 1998 | 0.1 | 1,831 | 1,903 | 2.27 | 1.2 |

Table 17 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Private fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1999 | 0.1 | 1,575 | 1,578 | 1.8 | 1.1 |
| 2000 | 0.1 | 1,366 | 1,337 | 1.6 | 1.2 |
| 2001 | 0.1 | 1,591 | 1,542 | 1.8 | 1.2 |
| 2002 | 0.1 | 1,298 | 1,305 | 1.6 | 1.2 |
| 2003 | 0.1 | 1,763 | 1,841 | 2.2 | 1.2 |
| 2004 | 0.1 | 2,174 | 2,365 | 2.7 | 1.2 |
| 2005 | 0.1 | 1,491 | 1,660 | 1.9 | 1.1 |
| 2006 | 0.1 | 1,386 | 1,514 | 1.7 | 1.1 |
| 2007 | 0.1 | 1,645 | 1,738 | 1.9 | 1.1 |
| 2008 | 0.1 | 2,298 | 2,520 | 2.8 | 1.1 |
| 2009 | 0.1 | 2,275 | 2,549 | 2.9 | 1.2 |
| 2010 | 0.1 | 1,193 | 1,274 | 1.5 | 1.2 |
| 2011 | 0.1 | 1,273 | 1,324 | 1.6 | 1.2 |
| 2012 | 0.1 | 3,333 | 3,054 | 3.6 | 1.2 |
| 2013 | 0.1 | 2,859 | 2,686 | 3.0 | 1.1 |
| 2014 | 0.1 | 3,508 | 3,182 | 3.5 | 1.1 |
| 2015 | 0.1 | 2,804 | 2,744 | 3.1 | 1.1 |
| 2016 | 0.1 | 2,745 | 2,831 | 3.2 | 1.1 |
| 2017 | 0.1 | 1,862 | 1,963 | 2.2 | 1.1 |
| 2018 | 0.1 | 1,996 | 2,070 | 2.3 | 1.1 |
| 2019 | 0.1 | 1,813 | 1,866 | 2.0 | 1.1 |
| 2020 | 0.1 | 2,464 | 2,537 | 2.7 | 1.1 |

Table 18. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Shore fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1945 | 0.1 | 0 | 0 | 0.000 | |
| 1946 | 0.1 | 39 | 39 | 0.016 | 0.4 |
| 1947 | 0.1 | 79 | 79 | 0.032 | 0.4 |
| 1948 | 0.1 | 118 | 118 | 0.048 | 0.4 |
| 1949 | 0.1 | 157 | 157 | 0.064 | 0.4 |
| 1950 | 0.1 | 197 | 197 | 0.080 | 0.4 |
| 1951 | 0.1 | 236 | 236 | 0.096 | 0.4 |
| 1952 | 0.1 | 275 | 275 | 0.112 | 0.4 |
| 1953 | 0.1 | 315 | 315 | 0.128 | 0.4 |
| 1954 | 0.1 | 354 | 354 | 0.143 | 0.4 |
| 1955 | 0.1 | 299 | 299 | 0.121 | 0.4 |
| 1956 | 0.1 | 331 | 331 | 0.134 | 0.4 |
| 1957 | 0.1 | 363 | 363 | 0.147 | 0.4 |
| 1958 | 0.1 | 395 | 395 | 0.160 | 0.4 |
| 1959 | 0.1 | 427 | 427 | 0.172 | 0.4 |
| 1960 | 0.1 | 459 | 459 | 0.185 | 0.4 |
| 1961 | 0.1 | 475 | 475 | 0.193 | 0.4 |
| 1962 | 0.1 | 490 | 490 | 0.198 | 0.4 |
| 1963 | 0.1 | 505 | 505 | 0.200 | 0.4 |
| 1964 | 0.1 | 521 | 521 | 0.207 | 0.4 |
| 1965 | 0.1 | 536 | 536 | 0.221 | 0.4 |
| 1966 | 0.1 | 552 | 553 | 0.230 | 0.4 |
| 1967 | 0.1 | 569 | 569 | 0.236 | 0.4 |
| 1968 | 0.1 | 585 | 586 | 0.240 | 0.4 |
| 1969 | 0.1 | 602 | 603 | 0.249 | 0.4 |
| 1970 | 0.1 | 618 | 619 | 0.256 | 0.4 |
| 1971 | 0.1 | 675 | 677 | 0.274 | 0.4 |

Table 18 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Shore fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1972 | 0.1 | 733 | 734 | 0.29 | 0.4 |
| 1973 | 0.1 | 790 | 791 | 0.30 | 0.4 |
| 1974 | 0.1 | 847 | 848 | 0.32 | 0.4 |
| 1975 | 0.1 | 905 | 906 | 0.35 | 0.4 |
| 1976 | 0.1 | 908 | 910 | 0.36 | 0.4 |
| 1977 | 0.1 | 912 | 914 | 0.37 | 0.4 |
| 1978 | 0.1 | 916 | 918 | 0.37 | 0.4 |
| 1979 | 0.1 | 919 | 923 | 0.38 | 0.4 |
| 1980 | 0.1 | 923 | 929 | 0.38 | 0.4 |
| 1981 | 0.1 | 1,261 | 1,294 | 0.51 | 0.4 |
| 1982 | 0.1 | 1,416 | 1,451 | 0.51 | 0.4 |
| 1983 | 0.1 | 1,183 | 1,180 | 0.44 | 0.4 |
| 1984 | 0.1 | 1,210 | 1,225 | 0.46 | 0.4 |
| 1985 | 0.1 | 597 | 606 | 0.23 | 0.4 |
| 1986 | 0.1 | 1,026 | 1,028 | 0.38 | 0.4 |
| 1987 | 0.1 | 1,169 | 891 | 0.36 | 0.4 |
| 1988 | 0.1 | 1,223 | 1,237 | 0.49 | 0.4 |
| 1989 | 0.1 | 2,311 | 2,318 | 0.81 | 0.4 |
| 1990 | 0.1 | 1,118 | 668 | 0.54 | 0.8 |
| 1991 | 0.1 | 3,148 | 2,240 | 1.84 | 0.8 |
| 1992 | 0.1 | 959 | 823 | 0.65 | 0.8 |
| 1993 | 0.1 | 973 | 916 | 0.72 | 0.8 |
| 1994 | 0.1 | 667 | 674 | 0.54 | 0.8 |
| 1995 | 0.1 | 625 | 638 | 0.53 | 0.8 |
| 1996 | 0.1 | 942 | 959 | 0.80 | 0.8 |
| 1997 | 0.1 | 468 | 477 | 0.39 | 0.8 |
| 1998 | 0.1 | 451 | 451 | 0.36 | 0.8 |

Table 18 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Shore fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1999 | 0.1 | 356 | 363 | 0.28 | 0.8 |
| 2000 | 0.1 | 499 | 498 | 0.40 | 0.8 |
| 2001 | 0.1 | 633 | 616 | 0.51 | 0.8 |
| 2002 | 0.1 | 305 | 309 | 0.26 | 0.8 |
| 2003 | 0.1 | 442 | 464 | 0.37 | 0.8 |
| 2004 | 0.1 | 502 | 536 | 0.43 | 0.8 |
| 2005 | 0.1 | 517 | 556 | 0.44 | 0.8 |
| 2006 | 0.1 | 265 | 278 | 0.22 | 0.8 |
| 2007 | 0.1 | 384 | 400 | 0.32 | 0.8 |
| 2008 | 0.1 | 698 | 736 | 0.59 | 0.8 |
| 2009 | 0.1 | 498 | 514 | 0.42 | 0.8 |
| 2010 | 0.1 | 129 | 134 | 0.11 | 0.8 |
| 2011 | 0.1 | 388 | 402 | 0.33 | 0.8 |
| 2012 | 0.1 | 683 | 664 | 0.54 | 0.8 |
| 2013 | 0.1 | 621 | 622 | 0.48 | 0.8 |
| 2014 | 0.1 | 981 | 969 | 0.76 | 0.8 |
| 2015 | 0.1 | 642 | 664 | 0.54 | 0.8 |
| 2016 | 0.1 | 977 | 1,026 | 0.83 | 0.8 |
| 2017 | 0.1 | 1,239 | 1,267 | 1.02 | 0.8 |
| 2018 | 0.1 | 1,107 | 1,134 | 0.90 | 0.8 |
| 2019 | 0.1 | 1,603 | 1,445 | 1.14 | 0.8 |
| 2020 | 0.1 | 1,144 | 1,178 | 0.92 | 0.8 |

Table 19. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Charter & Headboat fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1945 | 0.1 | 0.0 | 0.0 | 0.000 | |
| 1946 | 0.1 | 9.5 | 9.5 | 0.022 | 2.3 |
| 1947 | 0.1 | 19.1 | 19.1 | 0.044 | 2.3 |
| 1948 | 0.1 | 28.6 | 28.6 | 0.067 | 2.3 |
| 1949 | 0.1 | 38.1 | 38.1 | 0.089 | 2.3 |
| 1950 | 0.1 | 47.7 | 47.7 | 0.111 | 2.3 |
| 1951 | 0.1 | 57.2 | 57.2 | 0.133 | 2.3 |
| 1952 | 0.1 | 66.8 | 66.8 | 0.155 | 2.3 |
| 1953 | 0.1 | 76.3 | 76.3 | 0.177 | 2.3 |
| 1954 | 0.1 | 85.8 | 85.8 | 0.199 | 2.3 |
| 1955 | 0.1 | 72.6 | 72.6 | 0.168 | 2.3 |
| 1956 | 0.1 | 80.4 | 80.4 | 0.185 | 2.3 |
| 1957 | 0.1 | 88.1 | 88.1 | 0.202 | 2.3 |
| 1958 | 0.1 | 95.9 | 95.9 | 0.220 | 2.3 |
| 1959 | 0.1 | 103.6 | 103.6 | 0.237 | 2.3 |
| 1960 | 0.1 | 111.4 | 111.4 | 0.254 | 2.3 |
| 1961 | 0.1 | 115.1 | 115.1 | 0.262 | 2.3 |
| 1962 | 0.1 | 118.8 | 118.8 | 0.271 | 2.3 |
| 1963 | 0.1 | 122.5 | 122.5 | 0.279 | 2.3 |
| 1964 | 0.1 | 126.3 | 126.3 | 0.286 | 2.3 |
| 1965 | 0.1 | 130.0 | 130.0 | 0.293 | 2.3 |
| 1966 | 0.1 | 134.0 | 133.9 | 0.303 | 2.3 |
| 1967 | 0.1 | 137.9 | 137.9 | 0.316 | 2.3 |
| 1968 | 0.1 | 141.9 | 141.9 | 0.328 | 2.3 |
| 1969 | 0.1 | 145.9 | 145.9 | 0.340 | 2.3 |
| 1970 | 0.1 | 149.9 | 149.9 | 0.353 | 2.4 |
| 1971 | 0.1 | 163.8 | 163.8 | 0.389 | 2.4 |

Table 19 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Charter & Headboat fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1972 | 0.1 | 178 | 178 | 0.42 | 2.4 |
| 1973 | 0.1 | 192 | 192 | 0.45 | 2.4 |
| 1974 | 0.1 | 206 | 206 | 0.48 | 2.3 |
| 1975 | 0.1 | 219 | 220 | 0.50 | 2.3 |
| 1976 | 0.1 | 220 | 220 | 0.49 | 2.2 |
| 1977 | 0.1 | 221 | 221 | 0.49 | 2.2 |
| 1978 | 0.1 | 222 | 222 | 0.49 | 2.2 |
| 1979 | 0.1 | 223 | 223 | 0.50 | 2.2 |
| 1980 | 0.1 | 224 | 224 | 0.51 | 2.3 |
| 1981 | 0.1 | 186 | 192 | 0.44 | 2.3 |
| 1982 | 0.1 | 577 | 595 | 1.36 | 2.3 |
| 1983 | 0.1 | 401 | 417 | 0.91 | 2.2 |
| 1984 | 0.1 | 108 | 102 | 0.21 | 2.1 |
| 1985 | 0.1 | 102 | 105 | 0.21 | 2.0 |
| 1986 | 0.1 | 190 | 176 | 0.34 | 2.0 |
| 1987 | 0.1 | 126 | 110 | 0.21 | 1.9 |
| 1988 | 0.1 | 66 | 63 | 0.12 | 1.9 |
| 1989 | 0.1 | 135 | 139 | 0.27 | 2.0 |
| 1990 | 0.1 | 89 | 92 | 0.21 | 2.2 |
| 1991 | 0.1 | 229 | 237 | 0.53 | 2.2 |
| 1992 | 0.1 | 216 | 194 | 0.43 | 2.2 |
| 1993 | 0.1 | 125 | 113 | 0.24 | 2.1 |
| 1994 | 0.1 | 164 | 159 | 0.32 | 2.0 |
| 1995 | 0.1 | 109 | 112 | 0.23 | 2.0 |
| 1996 | 0.1 | 119 | 110 | 0.23 | 2.1 |
| 1997 | 0.1 | 71 | 74 | 0.16 | 2.1 |
| 1998 | 0.1 | 191 | 198 | 0.41 | 2.1 |

Table 19 Continued. Input (with log-scale standard errors, SE) and expected (Exp) landings for the Recreational Charter & Headboat fleet in numbers (N, 1,000s of fish) and weight (B, million pounds whole weight) for Gulf of Mexico Gray Snapper. The mean body weight (MW, gutted pounds per fish) was determined by dividing the expected landings in weights by the expected landings in numbers .

| Year | Input N SE | Input N | Exp N | Exp B | MW |
|------|------------|---------|-------|-------|-----|
| 1999 | 0.1 | 158 | 161 | 0.32 | 2.0 |
| 2000 | 0.1 | 113 | 117 | 0.23 | 1.9 |
| 2001 | 0.1 | 200 | 199 | 0.39 | 2.0 |
| 2002 | 0.1 | 143 | 146 | 0.29 | 2.0 |
| 2003 | 0.1 | 222 | 225 | 0.45 | 2.0 |
| 2004 | 0.1 | 196 | 197 | 0.39 | 2.0 |
| 2005 | 0.1 | 192 | 194 | 0.38 | 2.0 |
| 2006 | 0.1 | 193 | 195 | 0.38 | 1.9 |
| 2007 | 0.1 | 198 | 202 | 0.38 | 1.9 |
| 2008 | 0.1 | 219 | 227 | 0.43 | 1.9 |
| 2009 | 0.1 | 292 | 292 | 0.55 | 1.9 |
| 2010 | 0.1 | 149 | 154 | 0.30 | 1.9 |
| 2011 | 0.1 | 175 | 181 | 0.36 | 2.0 |
| 2012 | 0.1 | 173 | 179 | 0.36 | 2.0 |
| 2013 | 0.1 | 307 | 315 | 0.63 | 2.0 |
| 2014 | 0.1 | 348 | 353 | 0.68 | 1.9 |
| 2015 | 0.1 | 270 | 276 | 0.52 | 1.9 |
| 2016 | 0.1 | 382 | 390 | 0.75 | 1.9 |
| 2017 | 0.1 | 327 | 336 | 0.65 | 1.9 |
| 2018 | 0.1 | 327 | 338 | 0.65 | 1.9 |
| 2019 | 0.1 | 387 | 397 | 0.76 | 1.9 |
| 2020 | 0.1 | 309 | 318 | 0.59 | 1.9 |

Table 20. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Vertical Line fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.25), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1945 | | | 0.028 | 0.004 | 0.013 | 0.002 | 0.5 |
| 1946 | | | 0.029 | 0.004 | 0.013 | 0.002 | 0.5 |
| 1947 | | | 0.029 | 0.004 | 0.013 | 0.002 | 0.4 |
| 1948 | | | 0.030 | 0.004 | 0.013 | 0.002 | 0.4 |
| 1949 | | | 0.031 | 0.004 | 0.013 | 0.002 | 0.4 |
| 1950 | | | 0.032 | 0.004 | 0.013 | 0.000 | 0.4 |
| 1951 | | | 0.032 | 0.005 | 0.015 | 0.002 | 0.5 |
| 1952 | | | 0.034 | 0.005 | 0.015 | 0.002 | 0.5 |
| 1953 | | | 0.035 | 0.005 | 0.015 | 0.002 | 0.4 |
| 1954 | | | 0.036 | 0.005 | 0.015 | 0.002 | 0.4 |
| 1955 | | | 0.037 | 0.005 | 0.015 | 0.002 | 0.4 |
| 1956 | | | 0.038 | 0.005 | 0.015 | 0.002 | 0.4 |
| 1957 | | | 0.039 | 0.005 | 0.015 | 0.002 | 0.4 |
| 1958 | | | 0.040 | 0.006 | 0.018 | 0.004 | 0.4 |
| 1959 | | | 0.041 | 0.006 | 0.018 | 0.002 | 0.4 |
| 1960 | | | 0.042 | 0.006 | 0.018 | 0.002 | 0.4 |
| 1961 | | | 0.041 | 0.006 | 0.018 | 0.002 | 0.4 |
| 1962 | | | 0.046 | 0.007 | 0.020 | 0.002 | 0.4 |
| 1963 | | | 0.042 | 0.006 | 0.015 | 0.002 | 0.4 |
| 1964 | | | 0.045 | 0.006 | 0.020 | 0.002 | 0.4 |
| 1965 | | | 0.045 | 0.006 | 0.020 | 0.004 | 0.4 |
| 1966 | | | 0.037 | 0.005 | 0.018 | 0.002 | 0.5 |
| 1967 | | | 0.042 | 0.006 | 0.020 | 0.004 | 0.5 |
| 1968 | | | 0.052 | 0.007 | 0.022 | 0.002 | 0.4 |
| 1969 | | | 0.046 | 0.006 | 0.022 | 0.004 | 0.5 |
| 1970 | | | 0.042 | 0.006 | 0.020 | 0.002 | 0.5 |
| 1971 | | | 0.046 | 0.006 | 0.020 | 0.002 | 0.4 |

Table 20 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Vertical Line fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.25), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1972 | | | 0.056 | 0.007 | 0.024 | 0.002 | 0.4 |
| 1973 | | | 0.059 | 0.008 | 0.024 | 0.004 | 0.4 |
| 1974 | | | 0.063 | 0.008 | 0.024 | 0.002 | 0.4 |
| 1975 | | | 0.043 | 0.006 | 0.015 | 0.002 | 0.4 |
| 1976 | | | 0.077 | 0.011 | 0.029 | 0.002 | 0.4 |
| 1977 | | | 0.043 | 0.006 | 0.018 | 0.002 | 0.4 |
| 1978 | | | 0.035 | 0.005 | 0.015 | 0.002 | 0.4 |
| 1979 | | | 0.040 | 0.006 | 0.018 | 0.002 | 0.4 |
| 1980 | | | 0.071 | 0.010 | 0.031 | 0.004 | 0.4 |
| 1981 | | | 0.079 | 0.011 | 0.031 | 0.004 | 0.4 |
| 1982 | | | 0.149 | 0.021 | 0.046 | 0.007 | 0.3 |
| 1983 | | | 0.149 | 0.021 | 0.046 | 0.007 | 0.3 |
| 1984 | | | 0.115 | 0.016 | 0.033 | 0.004 | 0.3 |
| 1985 | | | 0.097 | 0.014 | 0.026 | 0.002 | 0.3 |
| 1986 | | | 0.120 | 0.017 | 0.033 | 0.004 | 0.3 |
| 1987 | | | 0.109 | 0.015 | 0.035 | 0.004 | 0.3 |
| 1988 | | | 0.073 | 0.010 | 0.022 | 0.004 | 0.3 |
| 1989 | | | 0.131 | 0.019 | 0.033 | 0.004 | 0.3 |
| 1990 | | | 3.342 | 0.468 | 1.217 | 0.170 | 0.4 |
| 1991 | | | 4.662 | 0.652 | 1.581 | 0.220 | 0.3 |
| 1992 | | | 5.470 | 0.766 | 1.812 | 0.254 | 0.3 |
| 1993 | 0.3 | 1.1 | 6.987 | 0.978 | 2.432 | 0.340 | 0.3 |
| 1994 | 0.3 | 1.4 | 7.566 | 1.059 | 2.758 | 0.386 | 0.4 |
| 1995 | 0.3 | 1.3 | 5.582 | 0.782 | 2.006 | 0.280 | 0.4 |
| 1996 | 0.3 | 1.2 | 5.095 | 0.713 | 1.821 | 0.256 | 0.4 |
| 1997 | 0.3 | 1.5 | 5.555 | 0.778 | 1.885 | 0.265 | 0.3 |
| 1998 | 0.3 | 1.3 | 5.391 | 0.754 | 1.788 | 0.251 | 0.3 |

Table 20 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Vertical Line fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.25), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1999 | 0.3 | 1.58 | 4.8 | 0.67 | 1.7 | 0.24 | 0.4 |
| 2000 | 0.3 | 1.26 | 4.8 | 0.67 | 1.7 | 0.24 | 0.4 |
| 2001 | 0.3 | 1.32 | 4.4 | 0.62 | 1.6 | 0.23 | 0.4 |
| 2002 | 0.3 | 1.47 | 5.6 | 0.79 | 1.9 | 0.27 | 0.3 |
| 2003 | 0.3 | 1.25 | 5.4 | 0.75 | 1.9 | 0.26 | 0.3 |
| 2004 | 0.3 | 1.31 | 6.0 | 0.84 | 2.1 | 0.29 | 0.3 |
| 2005 | 0.3 | 1.15 | 5.6 | 0.79 | 1.9 | 0.27 | 0.3 |
| 2006 | 0.3 | 1.08 | 5.1 | 0.72 | 1.8 | 0.25 | 0.3 |
| 2007 | 0.3 | 0.84 | 4.3 | 0.60 | 1.4 | 0.20 | 0.3 |
| 2008 | 0.3 | 0.82 | 4.1 | 0.57 | 1.5 | 0.20 | 0.4 |
| 2009 | 0.3 | 1.10 | 4.3 | 0.60 | 1.6 | 0.22 | 0.4 |
| 2010 | 0.3 | 0.50 | 3.2 | 0.44 | 1.1 | 0.16 | 0.4 |
| 2011 | 0.3 | 0.45 | 3.6 | 0.51 | 1.3 | 0.18 | 0.3 |
| 2012 | 0.3 | 0.52 | 4.5 | 0.63 | 1.5 | 0.21 | 0.3 |
| 2013 | 0.3 | 0.46 | 4.8 | 0.67 | 1.6 | 0.22 | 0.3 |
| 2014 | 0.3 | 0.57 | 5.2 | 0.73 | 1.9 | 0.27 | 0.4 |
| 2015 | 0.3 | 0.61 | 4.5 | 0.63 | 1.5 | 0.22 | 0.3 |
| 2016 | 0.3 | 0.61 | 4.4 | 0.62 | 1.5 | 0.22 | 0.4 |
| 2017 | 0.3 | 0.56 | 3.6 | 0.50 | 1.2 | 0.17 | 0.3 |
| 2018 | 0.3 | 0.46 | 4.1 | 0.57 | 1.4 | 0.19 | 0.3 |
| 2019 | 0.3 | 0.45 | 3.9 | 0.54 | 1.3 | 0.18 | 0.3 |
| 2020 | 0.3 | 0.43 | 3.4 | 0.47 | 1.1 | 0.16 | 0.3 |

Table 21. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Nets & Traps fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.25), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1945 | | | 0 | 0 | 0 | 0 | |
| 1946 | | | 0 | 0 | 0 | 0 | |
| 1947 | | | 0 | 0 | 0 | 0 | |
| 1948 | | | 0 | 0 | 0 | 0 | |
| 1949 | | | 0 | 0 | 0 | 0 | |
| 1950 | | | 0 | 0 | 0 | 0 | |
| 1951 | | | 0 | 0 | 0 | 0 | |
| 1952 | | | 0 | 0 | 0 | 0 | |
| 1953 | | | 0 | 0 | 0 | 0 | |
| 1954 | | | 0 | 0 | 0 | 0 | |
| 1955 | | | 0 | 0 | 0 | 0 | |
| 1956 | | | 0 | 0 | 0 | 0 | |
| 1957 | | | 0 | 0 | 0 | 0 | |
| 1958 | | | 0 | 0 | 0 | 0 | |
| 1959 | | | 0 | 0 | 0 | 0 | |
| 1960 | | | 0 | 0 | 0 | 0 | |
| 1961 | | | 0 | 0 | 0 | 0 | |
| 1962 | | | 0 | 0 | 0 | 0 | |
| 1963 | | | 0 | 0 | 0 | 0 | 1.6 |
| 1964 | | | 0 | 0 | 0 | 0 | |
| 1965 | | | 0 | 0 | 0 | 0 | |
| 1966 | | | 0 | 0 | 0 | 0 | |
| 1967 | | | 0 | 0 | 0 | 0 | |
| 1968 | | | 0 | 0 | 0 | 0 | |
| 1969 | | | 0 | 0 | 0 | 0 | |
| 1970 | | | 0 | 0 | 0 | 0 | |
| 1971 | | | 0 | 0 | 0 | 0 | |

Table 21 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Nets & Traps fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.25), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1972 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1973 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1974 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1975 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1976 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1977 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1978 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1979 | | | 0.000 | 0.000 | 0.000 | 0.000 | |
| 1980 | | | 0.001 | 0.000 | 0.001 | 0.000 | 1.6 |
| 1981 | | | 0.001 | 0.000 | 0.001 | 0.000 | 1.3 |
| 1982 | | | 0.002 | 0.000 | 0.002 | 0.000 | 1.2 |
| 1983 | | | 0.004 | 0.001 | 0.004 | 0.001 | 1.1 |
| 1984 | | | 0.002 | 0.000 | 0.002 | 0.000 | 1.1 |
| 1985 | | | 0.002 | 0.000 | 0.002 | 0.000 | 0.9 |
| 1986 | | | 0.002 | 0.000 | 0.002 | 0.000 | 1.2 |
| 1987 | | | 0.002 | 0.000 | 0.002 | 0.000 | 1.2 |
| 1988 | | | 0.002 | 0.000 | 0.002 | 0.000 | 0.9 |
| 1989 | | | 0.002 | 0.000 | 0.002 | 0.000 | 1.0 |
| 1990 | | | 0.199 | 0.028 | 0.115 | 0.016 | 0.6 |
| 1991 | | | 0.218 | 0.030 | 0.123 | 0.017 | 0.6 |
| 1992 | | | 0.276 | 0.039 | 0.150 | 0.021 | 0.5 |
| 1993 | 0.3 | 0.04 | 0.415 | 0.058 | 0.229 | 0.032 | 0.6 |
| 1994 | 0.3 | 0.04 | 0.055 | 0.008 | 0.031 | 0.004 | 0.6 |
| 1995 | 0.3 | 0.02 | 0.056 | 0.008 | 0.033 | 0.005 | 0.6 |
| 1996 | 0.3 | 0.02 | 0.040 | 0.006 | 0.023 | 0.003 | 0.6 |
| 1997 | 0.3 | 0.02 | 0.046 | 0.006 | 0.026 | 0.004 | 0.6 |
| 1998 | 0.3 | 0.02 | 0.056 | 0.008 | 0.031 | 0.004 | 0.5 |

Table 21 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Nets & Traps fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.25), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1999 | 0.3 | 0.04 | 0.093 | 0.013 | 0.051 | 0.007 | 0.6 |
| 2000 | 0.3 | 0.04 | 0.093 | 0.013 | 0.052 | 0.007 | 0.6 |
| 2001 | 0.3 | 0.04 | 0.078 | 0.011 | 0.045 | 0.006 | 0.6 |
| 2002 | 0.3 | 0.06 | 0.098 | 0.014 | 0.055 | 0.008 | 0.6 |
| 2003 | 0.3 | 0.04 | 0.072 | 0.010 | 0.040 | 0.006 | 0.6 |
| 2004 | 0.3 | 0.06 | 0.114 | 0.016 | 0.063 | 0.009 | 0.5 |
| 2005 | 0.3 | 0.04 | 0.103 | 0.014 | 0.056 | 0.008 | 0.5 |
| 2006 | 0.3 | 0.06 | 0.106 | 0.015 | 0.058 | 0.008 | 0.5 |
| 2007 | 0.3 | 0.04 | 0.104 | 0.014 | 0.056 | 0.008 | 0.5 |
| 2008 | 0.3 | 0.05 | 0.130 | 0.018 | 0.072 | 0.010 | 0.6 |
| 2009 | 0.3 | 0.02 | 0.135 | 0.019 | 0.077 | 0.011 | 0.6 |
| 2010 | 0.3 | 0.02 | 0.052 | 0.007 | 0.030 | 0.004 | 0.6 |
| 2011 | 0.3 | 0.03 | 0.093 | 0.013 | 0.053 | 0.007 | 0.6 |
| 2012 | 0.3 | 0.03 | 0.102 | 0.014 | 0.056 | 0.008 | 0.5 |
| 2013 | 0.3 | 0.04 | 0.102 | 0.014 | 0.055 | 0.008 | 0.5 |
| 2014 | 0.3 | 0.14 | 0.150 | 0.021 | 0.083 | 0.012 | 0.6 |
| 2015 | 0.3 | 0.16 | 0.214 | 0.030 | 0.119 | 0.017 | 0.6 |
| 2016 | 0.3 | 0.18 | 0.202 | 0.028 | 0.112 | 0.016 | 0.6 |
| 2017 | 0.3 | 0.20 | 0.170 | 0.024 | 0.093 | 0.013 | 0.5 |
| 2018 | 0.3 | 0.17 | 0.102 | 0.014 | 0.055 | 0.008 | 0.5 |
| 2019 | 0.3 | 0.15 | 0.125 | 0.018 | 0.066 | 0.009 | 0.5 |
| 2020 | 0.3 | 0.10 | 0.107 | 0.015 | 0.056 | 0.008 | 0.5 |

Table 22. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Nets & Traps fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1945 | | | 0 | 0.0 | 0.0 | 0.00 | |
| 1946 | | | 18 | 1.2 | 2.8 | 0.19 | 0.2 |
| 1947 | | | 35 | 2.4 | 5.6 | 0.39 | 0.2 |
| 1948 | | | 53 | 3.6 | 8.4 | 0.58 | 0.2 |
| 1949 | | | 71 | 4.9 | 11.3 | 0.78 | 0.2 |
| 1950 | | | 89 | 6.1 | 14.2 | 0.98 | 0.2 |
| 1951 | | | 107 | 7.4 | 17.1 | 1.18 | 0.2 |
| 1952 | | | 127 | 8.7 | 20.2 | 1.39 | 0.2 |
| 1953 | | | 146 | 10.1 | 23.2 | 1.60 | 0.2 |
| 1954 | | | 165 | 11.4 | 26.2 | 1.81 | 0.2 |
| 1955 | | | 140 | 9.7 | 22.3 | 1.54 | 0.2 |
| 1956 | | | 155 | 10.7 | 24.7 | 1.71 | 0.2 |
| 1957 | | | 170 | 11.7 | 27.1 | 1.86 | 0.2 |
| 1958 | | | 185 | 12.8 | 29.5 | 2.03 | 0.2 |
| 1959 | | | 203 | 14.0 | 32.3 | 2.23 | 0.2 |
| 1960 | | | 216 | 14.9 | 34.5 | 2.38 | 0.2 |
| 1961 | | | 219 | 15.1 | 34.9 | 2.41 | 0.2 |
| 1962 | | | 234 | 16.1 | 36.7 | 2.54 | 0.2 |
| 1963 | | | 254 | 17.5 | 40.3 | 2.78 | 0.2 |
| 1964 | | | 249 | 17.2 | 40.4 | 2.79 | 0.2 |
| 1965 | | | 233 | 16.1 | 37.6 | 2.60 | 0.2 |
| 1966 | | | 240 | 16.5 | 38.4 | 2.65 | 0.2 |
| 1967 | | | 251 | 17.3 | 40.0 | 2.76 | 0.2 |
| 1968 | | | 262 | 18.1 | 42.2 | 2.91 | 0.2 |
| 1969 | | | 261 | 18.0 | 41.8 | 2.88 | 0.2 |
| 1970 | | | 275 | 18.9 | 43.4 | 3.00 | 0.2 |
| 1971 | | | 322 | 22.2 | 50.3 | 3.47 | 0.2 |

Table 22 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Nets & Traps fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1972 | | | 382 | 26 | 60 | 4.1 | 0.2 |
| 1973 | | | 433 | 30 | 68 | 4.7 | 0.2 |
| 1974 | | | 464 | 32 | 73 | 5.0 | 0.2 |
| 1975 | | | 484 | 33 | 78 | 5.4 | 0.2 |
| 1976 | | | 453 | 31 | 73 | 5.0 | 0.2 |
| 1977 | | | 442 | 30 | 71 | 4.9 | 0.2 |
| 1978 | | | 429 | 30 | 69 | 4.7 | 0.2 |
| 1979 | | | 424 | 29 | 68 | 4.7 | 0.2 |
| 1980 | | | 430 | 30 | 67 | 4.6 | 0.2 |
| 1981 | 0.3 | 646 | 867 | 60 | 127 | 8.8 | 0.1 |
| 1982 | 0.3 | 609 | 815 | 56 | 129 | 8.9 | 0.2 |
| 1983 | 0.3 | 100 | 183 | 13 | 29 | 2.0 | 0.2 |
| 1984 | 0.3 | 661 | 977 | 67 | 153 | 10.6 | 0.2 |
| 1985 | 0.3 | 914 | 438 | 30 | 67 | 4.6 | 0.2 |
| 1986 | 0.3 | 561 | 361 | 25 | 60 | 4.1 | 0.2 |
| 1987 | 0.3 | 430 | 536 | 37 | 85 | 5.9 | 0.2 |
| 1988 | 0.3 | 817 | 563 | 39 | 80 | 5.5 | 0.1 |
| 1989 | 0.3 | 1,190 | 1,134 | 78 | 191 | 13.2 | 0.2 |
| 1990 | 0.3 | 2,222 | 2,651 | 183 | 721 | 49.8 | 0.3 |
| 1991 | 0.3 | 5,213 | 3,685 | 254 | 910 | 62.8 | 0.2 |
| 1992 | 0.3 | 4,896 | 3,144 | 217 | 822 | 56.7 | 0.3 |
| 1993 | 0.3 | 5,226 | 3,154 | 218 | 883 | 61.0 | 0.3 |
| 1994 | 0.3 | 3,996 | 2,056 | 142 | 578 | 39.9 | 0.3 |
| 1995 | 0.3 | 3,854 | 2,807 | 194 | 756 | 52.2 | 0.3 |
| 1996 | 0.3 | 4,120 | 2,265 | 156 | 590 | 40.7 | 0.3 |
| 1997 | 0.3 | 4,269 | 2,978 | 205 | 743 | 51.2 | 0.2 |
| 1998 | 0.3 | 5,879 | 4,486 | 310 | 1,182 | 81.5 | 0.3 |

Table 22 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Commercial Nets & Traps fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1999 | 0.3 | 4,667 | 2,708 | 187 | 755 | 52 | 0.3 |
| 2000 | 0.3 | 4,815 | 2,177 | 150 | 598 | 41 | 0.3 |
| 2001 | 0.3 | 3,576 | 2,190 | 151 | 588 | 41 | 0.3 |
| 2002 | 0.3 | 4,308 | 2,598 | 179 | 661 | 46 | 0.3 |
| 2003 | 0.3 | 5,406 | 3,676 | 254 | 967 | 67 | 0.3 |
| 2004 | 0.3 | 4,735 | 4,851 | 335 | 1,273 | 88 | 0.3 |
| 2005 | 0.3 | 5,118 | 3,339 | 230 | 883 | 61 | 0.3 |
| 2006 | 0.3 | 4,340 | 3,062 | 211 | 795 | 55 | 0.3 |
| 2007 | 0.3 | 4,176 | 3,737 | 258 | 994 | 69 | 0.3 |
| 2008 | 0.3 | 7,288 | 4,049 | 279 | 1,148 | 79 | 0.3 |
| 2009 | 0.3 | 4,547 | 3,617 | 250 | 1,000 | 69 | 0.3 |
| 2010 | 0.3 | 3,372 | 1,962 | 135 | 517 | 36 | 0.3 |
| 2011 | 0.3 | 3,755 | 2,508 | 173 | 631 | 44 | 0.3 |
| 2012 | 0.3 | 6,307 | 7,443 | 514 | 1,867 | 129 | 0.3 |
| 2013 | 0.3 | 8,823 | 6,338 | 437 | 1,711 | 118 | 0.3 |
| 2014 | 0.3 | 11,593 | 5,002 | 345 | 1,396 | 96 | 0.3 |
| 2015 | 0.3 | 9,381 | 5,345 | 369 | 1,394 | 96 | 0.3 |
| 2016 | 0.3 | 6,394 | 5,214 | 360 | 1,376 | 95 | 0.3 |
| 2017 | 0.3 | 7,490 | 4,296 | 296 | 1,103 | 76 | 0.3 |
| 2018 | 0.3 | 5,977 | 4,579 | 316 | 1,174 | 81 | 0.3 |
| 2019 | 0.3 | 6,005 | 4,664 | 322 | 1,190 | 82 | 0.3 |
| 2020 | 0.3 | 10,487 | 5,733 | 396 | 1,540 | 106 | 0.3 |

Table 23. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Recreational Private fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1945 | | | 0 | 0.00 | 0.00 | 0.000 | |
| 1946 | | | 4 | 0.27 | 0.51 | 0.035 | 0.1 |
| 1947 | | | 8 | 0.55 | 1.02 | 0.070 | 0.1 |
| 1948 | | | 12 | 0.82 | 1.53 | 0.105 | 0.1 |
| 1949 | | | 16 | 1.10 | 2.04 | 0.141 | 0.1 |
| 1950 | | | 20 | 1.38 | 2.56 | 0.176 | 0.1 |
| 1951 | | | 24 | 1.67 | 3.08 | 0.212 | 0.1 |
| 1952 | | | 28 | 1.96 | 3.61 | 0.249 | 0.1 |
| 1953 | | | 32 | 2.23 | 4.13 | 0.285 | 0.1 |
| 1954 | | | 36 | 2.52 | 4.65 | 0.321 | 0.1 |
| 1955 | | | 31 | 2.13 | 3.94 | 0.272 | 0.1 |
| 1956 | | | 34 | 2.35 | 4.36 | 0.301 | 0.1 |
| 1957 | | | 37 | 2.58 | 4.77 | 0.329 | 0.1 |
| 1958 | | | 41 | 2.82 | 5.20 | 0.359 | 0.1 |
| 1959 | | | 44 | 3.06 | 5.65 | 0.390 | 0.1 |
| 1960 | | | 47 | 3.24 | 6.02 | 0.415 | 0.1 |
| 1961 | | | 48 | 3.34 | 6.17 | 0.426 | 0.1 |
| 1962 | | | 52 | 3.57 | 6.52 | 0.450 | 0.1 |
| 1963 | | | 54 | 3.73 | 6.87 | 0.474 | 0.1 |
| 1964 | | | 53 | 3.62 | 6.82 | 0.471 | 0.1 |
| 1965 | | | 52 | 3.57 | 6.68 | 0.461 | 0.1 |
| 1966 | | | 54 | 3.75 | 6.98 | 0.481 | 0.1 |
| 1967 | | | 57 | 3.96 | 7.32 | 0.505 | 0.1 |
| 1968 | | | 59 | 4.07 | 7.60 | 0.525 | 0.1 |
| 1969 | | | 60 | 4.13 | 7.67 | 0.529 | 0.1 |
| 1970 | | | 63 | 4.38 | 8.04 | 0.553 | 0.1 |
| 1971 | | | 73 | 5.04 | 9.16 | 0.631 | 0.1 |

Table 23 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Recreational Private fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|--------|------------|---------|------------|-----|
| 1972 | | | 83 | 5.7 | 10.3 | 0.71 | 0.1 |
| 1973 | | | 89 | 6.2 | 11.2 | 0.78 | 0.1 |
| 1974 | | | 94 | 6.5 | 11.9 | 0.82 | 0.1 |
| 1975 | | | 97 | 6.7 | 12.4 | 0.85 | 0.1 |
| 1976 | | | 94 | 6.5 | 12.0 | 0.83 | 0.1 |
| 1977 | | | 93 | 6.4 | 11.9 | 0.82 | 0.1 |
| 1978 | | | 92 | 6.4 | 11.9 | 0.82 | 0.1 |
| 1979 | | | 92 | 6.4 | 11.9 | 0.82 | 0.1 |
| 1980 | | | 97 | 6.7 | 12.1 | 0.84 | 0.1 |
| 1981 | 0.3 | 836 | 162 | 11.2 | 19.2 | 1.33 | 0.1 |
| 1982 | 0.3 | 447 | 180 | 12.4 | 22.8 | 1.57 | 0.1 |
| 1983 | 0.3 | 1,653 | 135 | 9.3 | 16.9 | 1.17 | 0.1 |
| 1984 | 0.3 | 296 | 140 | 9.7 | 17.7 | 1.22 | 0.1 |
| 1985 | 0.3 | 344 | 72 | 5.0 | 8.9 | 0.61 | 0.1 |
| 1986 | 0.3 | 119 | 109 | 7.5 | 14.4 | 0.99 | 0.1 |
| 1987 | 0.3 | 43 | 84 | 5.8 | 10.8 | 0.74 | 0.1 |
| 1988 | 0.3 | 381 | 159 | 11.0 | 18.3 | 1.26 | 0.1 |
| 1989 | 0.3 | 682 | 262 | 18.1 | 35.2 | 2.43 | 0.1 |
| 1990 | 0.3 | 1,273 | 3,083 | 212.7 | 879.0 | 60.65 | 0.3 |
| 1991 | 0.3 | 7,836 | 15,870 | 1,095.0 | 4,022.4 | 277.54 | 0.3 |
| 1992 | 0.3 | 3,916 | 6,361 | 438.9 | 1,672.3 | 115.39 | 0.3 |
| 1993 | 0.3 | 4,092 | 5,129 | 353.9 | 1,440.1 | 99.37 | 0.3 |
| 1994 | 0.3 | 3,380 | 2,897 | 199.9 | 829.9 | 57.26 | 0.3 |
| 1995 | 0.3 | 3,467 | 3,062 | 211.3 | 836.0 | 57.68 | 0.3 |
| 1996 | 0.3 | 4,601 | 4,941 | 340.9 | 1,331.4 | 91.87 | 0.3 |
| 1997 | 0.3 | 3,543 | 3,304 | 228.0 | 848.7 | 58.56 | 0.3 |
| 1998 | 0.3 | 3,456 | 3,400 | 234.6 | 896.0 | 61.82 | 0.3 |

Table 23 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Recreational Private fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|--------|------------|-------|------------|-----|
| 1999 | 0.3 | 3,383 | 1,861 | 128 | 528 | 36 | 0.3 |
| 2000 | 0.3 | 3,501 | 2,526 | 174 | 696 | 48 | 0.3 |
| 2001 | 0.3 | 2,698 | 2,671 | 184 | 748 | 52 | 0.3 |
| 2002 | 0.3 | 2,298 | 1,998 | 138 | 516 | 36 | 0.3 |
| 2003 | 0.3 | 5,964 | 2,898 | 200 | 778 | 54 | 0.3 |
| 2004 | 0.3 | 4,781 | 3,371 | 233 | 900 | 62 | 0.3 |
| 2005 | 0.3 | 5,916 | 3,384 | 233 | 911 | 63 | 0.3 |
| 2006 | 0.3 | 2,957 | 1,673 | 115 | 447 | 31 | 0.3 |
| 2007 | 0.3 | 5,025 | 2,643 | 182 | 704 | 49 | 0.3 |
| 2008 | 0.3 | 7,291 | 3,553 | 245 | 1,016 | 70 | 0.3 |
| 2009 | 0.3 | 2,320 | 2,236 | 154 | 629 | 43 | 0.3 |
| 2010 | 0.3 | 1,621 | 653 | 45 | 177 | 12 | 0.3 |
| 2011 | 0.3 | 3,249 | 2,413 | 167 | 630 | 43 | 0.3 |
| 2012 | 0.3 | 4,346 | 5,086 | 351 | 1,305 | 90 | 0.3 |
| 2013 | 0.3 | 6,582 | 4,511 | 311 | 1,210 | 84 | 0.3 |
| 2014 | 0.3 | 5,884 | 4,340 | 299 | 1,256 | 87 | 0.3 |
| 2015 | 0.3 | 6,500 | 3,991 | 275 | 1,050 | 72 | 0.3 |
| 2016 | 0.3 | 6,993 | 5,683 | 392 | 1,548 | 107 | 0.3 |
| 2017 | 0.3 | 7,981 | 8,511 | 587 | 2,224 | 153 | 0.3 |
| 2018 | 0.3 | 7,808 | 7,426 | 512 | 1,966 | 136 | 0.3 |
| 2019 | 0.3 | 6,877 | 10,745 | 741 | 2,791 | 193 | 0.3 |
| 2020 | 0.3 | 12,176 | 7,856 | 542 | 2,121 | 146 | 0.3 |

Table 24. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Recreational Shore fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1945 | | | 0 | 0 | 0 | 0 | |
| 1946 | | | 1 | 0 | 0 | 0 | 0.5 |
| 1947 | | | 1 | 0 | 1 | 0 | 0.5 |
| 1948 | | | 2 | 0 | 1 | 0 | 0.5 |
| 1949 | | | 2 | 0 | 1 | 0 | 0.5 |
| 1950 | | | 3 | 0 | 1 | 0 | 0.5 |
| 1951 | | | 4 | 0 | 2 | 0 | 0.5 |
| 1952 | | | 4 | 0 | 2 | 0 | 0.5 |
| 1953 | | | 5 | 0 | 2 | 0 | 0.5 |
| 1954 | | | 5 | 0 | 2 | 0 | 0.5 |
| 1955 | | | 5 | 0 | 2 | 0 | 0.5 |
| 1956 | | | 5 | 0 | 2 | 0 | 0.5 |
| 1957 | | | 6 | 0 | 3 | 0 | 0.5 |
| 1958 | | | 6 | 0 | 3 | 0 | 0.5 |
| 1959 | | | 7 | 0 | 3 | 0 | 0.5 |
| 1960 | | | 7 | 1 | 3 | 0 | 0.5 |
| 1961 | | | 7 | 1 | 3 | 0 | 0.5 |
| 1962 | | | 8 | 1 | 4 | 0 | 0.5 |
| 1963 | | | 8 | 1 | 4 | 0 | 0.5 |
| 1964 | | | 9 | 1 | 4 | 0 | 0.5 |
| 1965 | | | 9 | 1 | 4 | 0 | 0.5 |
| 1966 | | | 8 | 1 | 4 | 0 | 0.5 |
| 1967 | | | 8 | 1 | 4 | 0 | 0.5 |
| 1968 | | | 9 | 1 | 4 | 0 | 0.5 |
| 1969 | | | 9 | 1 | 4 | 0 | 0.5 |
| 1970 | | | 9 | 1 | 4 | 0 | 0.5 |
| 1971 | | | 10 | 1 | 4 | 0 | 0.5 |

Table 24 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Recreational Shore fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1972 | | | 11 | 1 | 5 | 0 | 0.4 |
| 1973 | | | 13 | 1 | 6 | 0 | 0.4 |
| 1974 | | | 15 | 1 | 7 | 0 | 0.4 |
| 1975 | | | 17 | 1 | 8 | 1 | 0.4 |
| 1976 | | | 17 | 1 | 7 | 1 | 0.5 |
| 1977 | | | 16 | 1 | 7 | 0 | 0.5 |
| 1978 | | | 15 | 1 | 7 | 0 | 0.5 |
| 1979 | | | 15 | 1 | 7 | 0 | 0.5 |
| 1980 | | | 14 | 1 | 6 | 0 | 0.5 |
| 1981 | 0.3 | 29 | 12 | 1 | 6 | 0 | 0.5 |
| 1982 | 0.3 | 185 | 49 | 3 | 21 | 1 | 0.4 |
| 1983 | 0.3 | 72 | 39 | 3 | 17 | 1 | 0.4 |
| 1984 | 0.3 | 7 | 10 | 1 | 4 | 0 | 0.4 |
| 1985 | 0.3 | 17 | 10 | 1 | 4 | 0 | 0.4 |
| 1986 | 0.3 | 13 | 18 | 1 | 8 | 1 | 0.4 |
| 1987 | 0.3 | 6 | 10 | 1 | 4 | 0 | 0.5 |
| 1988 | 0.3 | 4 | 5 | 0 | 2 | 0 | 0.5 |
| 1989 | 0.3 | 19 | 14 | 1 | 6 | 0 | 0.4 |
| 1990 | 0.3 | 48 | 34 | 2 | 20 | 1 | 0.6 |
| 1991 | 0.3 | 194 | 87 | 6 | 49 | 3 | 0.6 |
| 1992 | 0.3 | 59 | 90 | 6 | 49 | 3 | 0.5 |
| 1993 | 0.3 | 38 | 57 | 4 | 32 | 2 | 0.6 |
| 1994 | 0.3 | 57 | 69 | 5 | 40 | 3 | 0.6 |
| 1995 | 0.3 | 186 | 41 | 3 | 24 | 2 | 0.6 |
| 1996 | 0.3 | 27 | 39 | 3 | 23 | 2 | 0.6 |
| 1997 | 0.3 | 58 | 29 | 2 | 16 | 1 | 0.6 |
| 1998 | 0.3 | 147 | 96 | 7 | 53 | 4 | 0.6 |

Table 24 Continued. Input (with log-scale standard errors, SE) and expected (Exp) discards for the Recreational Shore fleet in number (N, 1,000s of fish) and biomass (B, thousand pounds whole weight) for Gulf of Mexico Gray Snapper. Dead discards in numbers (discard mortality rate = 0.12), dead discards in biomass, and mean weight (MW, gutted pounds per fish) are included. Mean weight was determined by dividing the expected discards in weights by the expected discards in numbers .

| Year | Input N SE | Input N | Exp N | Exp Dead N | Exp B | Exp Dead B | MW |
|------|------------|---------|-------|------------|-------|------------|-----|
| 1999 | 0.3 | 101 | 81 | 6 | 47 | 3 | 0.6 |
| 2000 | 0.3 | 117 | 51 | 4 | 30 | 2 | 0.6 |
| 2001 | 0.3 | 79 | 78 | 5 | 45 | 3 | 0.6 |
| 2002 | 0.3 | 94 | 56 | 4 | 32 | 2 | 0.6 |
| 2003 | 0.3 | 120 | 100 | 7 | 56 | 4 | 0.6 |
| 2004 | 0.3 | 98 | 93 | 6 | 52 | 4 | 0.6 |
| 2005 | 0.3 | 101 | 94 | 6 | 53 | 4 | 0.6 |
| 2006 | 0.3 | 97 | 94 | 6 | 53 | 4 | 0.6 |
| 2007 | 0.3 | 120 | 101 | 7 | 57 | 4 | 0.6 |
| 2008 | 0.3 | 175 | 110 | 8 | 63 | 4 | 0.6 |
| 2009 | 0.3 | 119 | 120 | 8 | 71 | 5 | 0.6 |
| 2010 | 0.3 | 142 | 57 | 4 | 33 | 2 | 0.6 |
| 2011 | 0.3 | 156 | 68 | 5 | 39 | 3 | 0.6 |
| 2012 | 0.3 | 153 | 80 | 6 | 44 | 3 | 0.6 |
| 2013 | 0.3 | 288 | 171 | 12 | 95 | 7 | 0.6 |
| 2014 | 0.3 | 272 | 180 | 12 | 105 | 7 | 0.6 |
| 2015 | 0.3 | 266 | 122 | 8 | 70 | 5 | 0.6 |
| 2016 | 0.3 | 262 | 177 | 12 | 101 | 7 | 0.6 |
| 2017 | 0.3 | 250 | 156 | 11 | 88 | 6 | 0.6 |
| 2018 | 0.3 | 367 | 170 | 12 | 95 | 7 | 0.6 |
| 2019 | 0.3 | 317 | 214 | 15 | 119 | 8 | 0.6 |
| 2020 | 0.3 | 499 | 184 | 13 | 103 | 7 | 0.6 |

Table 25. Observed (Obs) versus predicted (Exp) standardized fishery-dependent catch-per-unit-effort (CPUE) indices for Gulf of Mexico Gray Snapper. Values are normalized to the mean. CVs as estimated by the standardization process were scaled to a common mean of 0.20 were converted to log-scale SEs.

| Yr | Private (Obs) | Private (Exp) | Private (SE) | Shore (Obs) | Shore (Exp) | Shore (SE) |
|------|------------------|------------------|-----------------|----------------|----------------|---------------|
| 1981 | 1.06 | 1.17 | 0.37 | 0.40 | 0.69 | 0.32 |
| 1982 | 0.58 | 1.09 | 0.57 | 0.42 | 0.86 | 0.29 |
| 1983 | 0.53 | 1.09 | 0.71 | 0.38 | 0.85 | 0.35 |
| 1984 | 0.63 | 1.06 | 0.57 | 0.47 | 0.88 | 0.33 |
| 1985 | 1.35 | 1.10 | 0.45 | 0.48 | 0.90 | 0.34 |
| 1986 | 0.83 | 1.01 | 0.29 | 0.50 | 1.00 | 0.47 |
| 1987 | 0.92 | 0.98 | 0.28 | 0.19 | 0.80 | 0.53 |
| 1988 | 0.51 | 1.13 | 0.54 | 0.31 | 0.78 | 0.57 |
| 1989 | 1.30 | 0.94 | 0.25 | 1.10 | 1.04 | 0.25 |
| 1990 | 1.00 | 1.02 | 0.27 | 0.47 | 0.74 | 0.29 |
| 1991 | 1.01 | 1.06 | 0.26 | 1.28 | 0.97 | 0.21 |
| 1992 | 1.20 | 0.98 | 0.14 | 0.79 | 1.10 | 0.18 |
| 1993 | 1.22 | 0.87 | 0.13 | 0.99 | 0.98 | 0.12 |
| 1994 | 1.18 | 0.83 | 0.13 | 0.76 | 0.79 | 0.13 |
| 1995 | 1.09 | 0.78 | 0.15 | 0.79 | 0.76 | 0.13 |
| 1996 | 0.83 | 0.84 | 0.14 | 1.08 | 0.72 | 0.14 |
| 1997 | 1.03 | 0.95 | 0.13 | 1.06 | 0.90 | 0.14 |
| 1998 | 0.99 | 0.86 | 0.11 | 0.94 | 1.08 | 0.13 |
| 1999 | 0.76 | 0.86 | 0.11 | 0.70 | 0.90 | 0.13 |
| 2000 | 0.89 | 0.78 | 0.13 | 0.86 | 0.90 | 0.15 |
| 2001 | 0.89 | 0.88 | 0.12 | 0.59 | 0.74 | 0.16 |
| 2002 | 0.89 | 0.90 | 0.11 | 0.91 | 0.96 | 0.14 |
| 2003 | 1.05 | 0.95 | 0.09 | 1.57 | 0.99 | 0.11 |

Table 25 Continued. Observed (Obs) versus predicted (Exp) standardized fishery-dependent catch-per-unit-effort (CPUE) indices for Gulf of Mexico Gray Snapper. Values are normalized to the mean. CVs as estimated by the standardization process were scaled to a common mean of 0.20 were converted to log-scale SEs.

| Yr | Private (Obs) | Private (Exp) | Private (SE) | Shore (Obs) | Shore (Exp) | Shore (SE) |
|------|------------------|------------------|-----------------|----------------|----------------|---------------|
| 2004 | 0.68 | 0.96 | 0.13 | 1.09 | 1.06 | 0.14 |
| 2005 | 0.83 | 0.99 | 0.12 | 1.67 | 1.08 | 0.12 |
| 2006 | 0.99 | 1.12 | 0.12 | 1.07 | 1.15 | 0.17 |
| 2007 | 1.11 | 1.04 | 0.11 | 1.86 | 1.35 | 0.11 |
| 2008 | 1.03 | 0.96 | 0.10 | 1.23 | 1.11 | 0.13 |
| 2009 | 0.92 | 0.93 | 0.11 | 0.79 | 0.95 | 0.14 |
| 2010 | 0.57 | 0.99 | 0.17 | 0.59 | 0.94 | 0.17 |
| 2011 | 0.68 | 1.19 | 0.17 | 0.61 | 1.10 | 0.19 |
| 2012 | 0.79 | 1.31 | 0.14 | 0.98 | 1.42 | 0.15 |
| 2013 | 1.26 | 1.11 | 0.12 | 1.23 | 1.58 | 0.12 |
| 2014 | 1.50 | 1.22 | 0.08 | 1.47 | 1.16 | 0.14 |
| 2015 | 1.33 | 1.15 | 0.09 | 1.32 | 1.37 | 0.15 |
| 2016 | 1.22 | 1.27 | 0.09 | 1.45 | 1.25 | 0.13 |
| 2017 | 1.15 | 1.31 | 0.11 | 1.43 | 1.48 | 0.13 |
| 2018 | 1.33 | 1.54 | 0.11 | 1.95 | 1.57 | 0.13 |
| 2019 | 1.26 | 1.66 | 0.10 | 1.84 | 1.95 | 0.14 |
| 2020 | 1.60 | 1.58 | 0.09 | 2.36 | 2.06 | 0.11 |

Table 26. Observed (Obs) versus predicted (Exp) standardized fishery-independent indices and associated lognormal standard error (as estimated by the standardization process) for Gulf of Mexico Gray Snapper. Values are normalized to the mean. CVs as estimated by the standardization process were converted to log-scale SEs. Trawl = SEAMAP Trawl Survey, CombVid = Combined Video Survey, RF = Reef fish visual survey.

| Year | Age0 (Obs) | Age0 (Exp) | Age0 (SE) | Age1 (Obs) | Age1 (Exp) | Age1 (SE) | Trawl (Obs) | Trawl (Exp) | Trawl (SE) | Comb Vid (Obs) | Comb Vid (Exp) | Comb Vid (SE) | RF (Obs) | RF (Exp) | RF (SE) |
|------|---------------|---------------|--------------|---------------|---------------|--------------|----------------|----------------|---------------|----------------------|----------------------|---------------------|-------------|-------------|------------|
| 1993 | | | | | | | | | | 0.720 | 0.800 | 0.315 | | | |
| 1994 | | | | | | | | | | 1.063 | 0.796 | 0.331 | | | |
| 1995 | | | | | | | | | | 0.695 | 0.760 | 0.369 | | | |
| 1996 | | | | 0.149 | 0.596 | 0.288 | | | | 0.645 | 0.715 | 0.264 | | | |
| 1997 | | | | 0.293 | 0.682 | 0.229 | | | | 1.029 | 0.703 | 0.272 | 0.371 | 0.765 | 0.137 |
| 1998 | 0.168 | 0.571 | 0.305 | 0.340 | 0.618 | 0.255 | | | | | | | 0.738 | 0.869 | 0.123 |
| 1999 | 0.927 | 0.720 | 0.233 | 0.280 | 0.623 | 0.230 | | | | | | | 0.755 | 0.801 | 0.106 |
| 2000 | 0.660 | 0.453 | 0.209 | 0.248 | 0.557 | 0.214 | | | | | | | 0.815 | 0.806 | 0.107 |
| 2001 | 1.343 | 0.915 | 0.188 | 0.226 | 0.640 | 0.249 | | | | | | | 0.958 | 0.726 | 0.118 |
| 2002 | 1.622 | 0.789 | 0.159 | 0.810 | 0.657 | 0.194 | | | | 0.863 | 0.771 | 0.280 | 1.053 | 0.828 | 0.121 |
| 2003 | 1.114 | 0.896 | 0.156 | 0.701 | 0.698 | 0.219 | | | | | | | 0.770 | 0.854 | 0.151 |
| 2004 | 1.725 | 0.877 | 0.159 | 0.716 | 0.711 | 0.211 | | | | 0.317 | 0.802 | 0.358 | 1.017 | 0.892 | 0.122 |
| 2005 | 0.446 | 0.923 | 0.200 | 0.535 | 0.737 | 0.237 | | | | 0.230 | 0.826 | 0.374 | 1.069 | 0.918 | 0.140 |
| 2006 | 1.302 | 1.205 | 0.147 | 0.221 | 0.841 | 0.239 | | | | 0.925 | 0.874 | 0.203 | 0.592 | 0.975 | 0.122 |
| 2007 | 0.970 | 0.716 | 0.165 | 1.897 | 0.777 | 0.191 | | | | 1.148 | 0.944 | 0.287 | 0.854 | 1.101 | 0.139 |
| 2008 | 0.470 | 0.659 | 0.190 | 0.955 | 0.716 | 0.201 | | | | 1.872 | 0.982 | 0.229 | 0.907 | 1.004 | 0.110 |
| 2009 | 1.320 | 0.716 | 0.150 | 0.747 | 0.686 | 0.203 | | | | 1.433 | 0.955 | 0.185 | 1.196 | 0.912 | 0.112 |
| 2010 | 0.867 | 0.922 | 0.163 | 0.274 | 0.734 | 0.234 | 1.101 | 0.791 | 0.227 | 0.744 | 0.934 | 0.185 | 0.935 | 0.898 | 0.098 |
| 2011 | 0.531 | 1.350 | 0.181 | 0.172 | 0.889 | 0.276 | 0.818 | 0.821 | 0.266 | 0.659 | 0.949 | 0.129 | 2.066 | 0.981 | 0.118 |
| 2012 | 1.395 | 1.411 | 0.173 | 0.512 | 0.989 | 0.182 | 0.822 | 0.923 | 0.228 | 0.880 | 0.975 | 0.138 | 1.167 | 1.151 | 0.095 |
| 2013 | 0.359 | 0.615 | 0.301 | 1.460 | 0.832 | 0.187 | 0.825 | 1.054 | 0.263 | 0.835 | 1.036 | 0.170 | | | |
| 2014 | 1.410 | 1.278 | 0.161 | 0.851 | 0.914 | 0.192 | 1.160 | 0.998 | 0.210 | 1.050 | 1.063 | 0.125 | 0.953 | 1.068 | 0.104 |
| 2015 | 0.615 | 0.907 | 0.190 | 2.117 | 0.863 | 0.180 | 0.955 | 0.988 | 0.249 | 1.124 | 1.053 | 0.134 | | | |
| 2016 | 1.639 | 1.360 | 0.149 | 2.220 | 0.955 | 0.188 | 1.099 | 0.971 | 0.218 | 1.096 | 1.046 | 0.119 | 0.910 | 1.105 | 0.102 |
| 2017 | 0.876 | 1.276 | 0.169 | 3.366 | 0.991 | 0.185 | 0.950 | 1.020 | 0.254 | 1.373 | 1.061 | 0.165 | | | |
| 2018 | 0.968 | 1.810 | 0.168 | 2.320 | 1.168 | 0.197 | 0.655 | 1.097 | 0.238 | 0.753 | 1.115 | 0.167 | 1.389 | 1.295 | 0.092 |
| 2019 | 1.046 | 1.732 | 0.164 | 1.948 | 1.265 | 0.182 | 1.616 | 1.259 | 0.244 | 1.204 | 1.214 | 0.167 | | | |
| 2020 | 1.228 | 1.227 | 0.167 | 1.642 | 1.196 | 0.197 | | | | 2.342 | 1.350 | 0.156 | | | |

Table 27. Summary of correlated parameters with correlation coefficients > 0.7 parameters for Gulf of Mexico Gray Snapper from the SEDAR75 base model.

| Parameter 1 | Parameter 2 | Correlation |
|--|-------------------------------------|-------------|
| Age_DblN_ascend_se_Com_LL_2(2) | Age_DblN_peak_Com_LL_2(2) | 0.87 |
| Age_DblN_ascend_se_Com_VL_1(1) | Size_DblN_end_logit_Com_VL_1(1) | 0.85 |
| Age_DblN_ascend_se_Rec_HB_CBT_6(6) | Age_DblN_peak_Rec_HB_CBT_6(6) | 0.95 |
| Age_DblN_ascend_se_Rec_PR_4(4) | Age_DblN_peak_Rec_PR_4(4) | 0.85 |
| Age_DblN_descend_se_Com_VL_1(1) | Age_DblN_top_logit_Com_VL_1(1) | -0.78 |
| Age_DblN_end_logit_Com_LL_2(2) | Age_DblN_descend_se_Com_LL_2(2) | -0.95 |
| Age_DblN_start_logit_Rec_HB_CBT_6(6) | Age_DblN_peak_Rec_HB_CBT_6(6) | -0.85 |
| Age_DblN_start_logit_Rec_HB_CBT_6(6) | Age_DblN_ascend_se_Rec_HB_CBT_6(6) | -0.78 |
| Retain_L_infl_Com_VL_1(1)_BLK1repl_1990 | Size_DblN_start_logit_Com_VL_1(1) | -0.77 |
| Retain_L_width_Rec_Shore_5(5) | Retain_L_infl_Rec_Shore_5(5) | -0.81 |
| Size_DblN_ascend_se_Com_LL_2(2) | Size_DblN_peak_Com_LL_2(2) | 0.71 |
| Size_DblN_ascend_se_Rec_HB_CBT_6(6) | Size_DblN_peak_Rec_HB_CBT_6(6) | 0.90 |
| Size_DblN_ascend_se_SEAMAP_Trawl_11(11) | Size_DblN_peak_SEAMAP_Trawl_11(11) | 0.86 |
| Size_DblN_ascend_se_Visual_Survey_13(13) | Size_DblN_peak_Visual_Survey_13(13) | 0.97 |

Table 28. Retrospective analysis and retrospective forecast spawning stock biomass (males and females combined, metric tons) and fishing mortality (F, total biomass killed age 2+ / total biomass age 2+) for the last five terminal years and combined (grey rows) for the Gulf of Mexico Gray Snapper SEDAR75. N = number of observations to compute each statistic. Values within -0.15 to 0.2 are highlighted in green and are considered acceptable levels of retrospective bias. See Carvalho et al. (2021) for additional details.

| Diagnostics | Quantity | Source | Statistic | Value | N |
|------------------------|-----------------|--------|------------|--------|---|
| Retrospective analysis | SSB (-2019) | | Mohn's Rho | 0.001 | 1 |
| Retrospective analysis | SSB (-2018) | | Mohn's Rho | 0.018 | 1 |
| Retrospective analysis | SSB (-2017) | | Mohn's Rho | 0.035 | 1 |
| Retrospective analysis | SSB (-2016) | | Mohn's Rho | -0.006 | 1 |
| Retrospective analysis | SSB (-2015) | | Mohn's Rho | -0.099 | 1 |
| Retrospective analysis | SSB (-Combined) | | Mohn's Rho | -0.010 | 5 |
| Retrospective analysis | F (-2019) | | Mohn's Rho | 0.004 | 1 |
| Retrospective analysis | F (-2018) | | Mohn's Rho | -0.027 | 1 |
| Retrospective analysis | F (-2017) | | Mohn's Rho | -0.050 | 1 |
| Retrospective analysis | F (-2016) | | Mohn's Rho | -0.016 | 1 |
| Retrospective analysis | F (-2015) | | Mohn's Rho | 0.124 | 1 |
| Retrospective analysis | F (-Combined) | | Mohn's Rho | 0.007 | 5 |

Table 29. Joint residual summary statistics for the Gulf of Mexico Gray Snapper SEDAR75. N = number of observations to compute each statistic. RMSE = root mean squared error (as a percentage), with values above 30% for joint residuals (grey rows) highlighted in red if present and acceptable values below 30% highlighted in green. See Carvalho et al. (2021) for additional details.

| Quantity | Statistic | Value | N |
|---------------------------|-----------|-------|-----|
| Index of Abundance | | | |
| Index_Private_7 | RMSE(%) | 30.4 | 40 |
| Index_Shore_8 | RMSE(%) | 44.4 | 40 |
| FWRI_Age0_9 | RMSE(%) | 51.6 | 23 |
| FWRI_Age1_10 | RMSE(%) | 81 | 25 |
| SEAMAP_Trawl_11 | RMSE(%) | 23.6 | 10 |
| Combo_Video_12 | RMSE(%) | 42.5 | 23 |
| Visual_Survey_13 | RMSE(%) | 30.4 | 19 |
| Combined | RMSE(%) | 47.5 | 180 |
| Age | | | |
| Com_VL_1 | RMSE(%) | 26.7 | 29 |
| Com_LL_2 | RMSE(%) | 8.7 | 21 |
| Rec_PR_4 | RMSE(%) | 31.1 | 20 |
| Rec_HB_CBT_6 | RMSE(%) | 21.1 | 32 |
| Combined | RMSE(%) | 23.4 | 102 |
| Length | | | |
| Com_VL_1 | RMSE(%) | 22 | 6 |
| Com_LL_2 | RMSE(%) | 6.8 | 11 |
| Com_NT_3 | RMSE(%) | 9 | 18 |
| Rec_PR_4 | RMSE(%) | 8.7 | 16 |
| Rec_Shore_5 | RMSE(%) | 8.7 | 37 |
| Rec_HB_CBT_6 | RMSE(%) | 6.1 | 8 |
| SEAMAP_Trawl_11 | RMSE(%) | 4.6 | 10 |
| Combo_Video_12 | RMSE(%) | 11.1 | 14 |
| Visual_Survey_13 | RMSE(%) | 11.2 | 19 |
| Combined | RMSE(%) | 9.9 | 139 |

Table 30. Runs tests summary statistics for the Gulf of Mexico Gray Snapper SEDAR75. N = number of observations to compute each statistic. P-values greater than 0.05% (in green) provide support for randomly distributed residuals whereas p-values less than 0.05% (in red) indicate non-randomly distributed residuals. See Carvalho et al. (2021) for additional details.

| Quantity | Statistic | Value | N |
|---------------------------|-----------|-------|----|
| Index of Abundance | | | |
| Index_Private_7 | p-value | 0.202 | 40 |
| Index_Shore_8 | p-value | 0.534 | 40 |
| FWRI_Age0_9 | p-value | 0.419 | 23 |
| FWRI_Age1_10 | p-value | 0.012 | 25 |
| SEAMAP_Trawl_11 | p-value | 0.8 | 10 |
| Combo_Video_12 | p-value | 0.145 | 23 |
| Visual_Survey_13 | p-value | 0.121 | 19 |
| Age | | | |
| Com_VL_1 | p-value | 0.612 | 29 |
| Com_LL_2 | p-value | 0.278 | 21 |
| Rec_PR_4 | p-value | 0.631 | 20 |
| Rec_HB_CBT_6 | p-value | 0.038 | 32 |
| Length | | | |
| Com_VL_1 | p-value | 0.76 | 6 |
| Com_LL_2 | p-value | 0.475 | 11 |
| Com_NT_3 | p-value | 0.709 | 18 |
| Rec_PR_4 | p-value | 0.921 | 16 |
| Rec_Shore_5 | p-value | 0.002 | 37 |
| Rec_HB_CBT_6 | p-value | 0.777 | 8 |
| SEAMAP_Trawl_11 | p-value | 0.435 | 10 |
| Combo_Video_12 | p-value | 0.003 | 14 |
| Visual_Survey_13 | p-value | 0.64 | 19 |

Table 31. Hindcast cross-validation summary statistics for the Gulf of Mexico Gray Snapper SEDAR75. N = number of observations to compute each statistic. MASE = mean absolute scaled error, with values < 1 (in green) indicative of superior prediction skill over a naïve baseline forecast (random walk) and values > 1 (in red) indicative of poor prediction skill. See Carvalho et al. (2021) for additional details.

| Quantity | Statistic | Value | N |
|---------------------------|-----------|-------|---|
| Index of Abundance | | | |
| Index_Private_7 | MASE | 0.763 | 5 |
| Index_Shore_8 | MASE | 1.655 | 5 |
| FWRI_Age0_9 | MASE | 0.599 | 5 |
| FWRI_Age1_10 | MASE | 4.293 | 5 |
| SEAMAP_Trawl_11 | MASE | 0.687 | 4 |
| Combo_Video_12 | MASE | 0.693 | 5 |
| Visual_Survey_13 | MASE | 0.57 | 1 |
| Com_VL_1 | MASE | 0.697 | 5 |
| Age | | | |
| Com_LL_2 | MASE | 1.881 | 5 |
| Rec_PR_4 | MASE | 2.841 | 5 |
| Rec_HB_CBT_6 | MASE | 1.19 | 5 |
| Rec_Shore_5 | MASE | 2.795 | 5 |
| SEAMAP_Trawl_11 | MASE | 0.797 | 4 |
| Length | | | |
| Combo_Video_12 | MASE | 1.983 | 5 |
| Visual_Survey_13 | MASE | 0.014 | 1 |

Table 32. Summary of key model building steps towards the SEDAR75 Base Model for Gulf of Mexico Gray Snapper and associated convergence diagnostics. Note that steps within each model progression are not shown due to the vast number of intermediate runs conducted.

| Model Name | Description | SS Version | NLL | Gradient | Bounded Params |
|-----------------------------|--|---------------|--------|----------|----------------|
| 0. S51 3.24 | Base model run from SEDAR 51 | 3.2 | 517 | 0.0002 | 0 |
| 1. Convert to 3.30.17 | Convert S51 base model from SS 3.24 to SS 3.30.17 | 3.3 | 519 | 0.0000 | 0 |
| 2. Corrected COM data | Correct commercial landings in S51 fleet structure | 3.3 | 513 | 0.6910 | 0 |
| 3. TY 2020 Rec | Extend out Recreational Data inputs to 2020 | 3.3 | 2,885 | 0.0004 | 1 |
| 4. TY 2020 - new COM fleets | New commercial fleet structure and data extended to 2020 | 3.3 | 5,962 | 0.0035 | 2 |
| 5. FI Length comps | Add fishery independent length comps | 3.3 | 12,108 | 0.0014 | 1 |
| 6. All Age comps added | Add age comps | 3.3 | 9,857 | 23.4000 | 1 |
| 7. max age 21 fix growth | Max age of 21 with growth fixed | 3.3 | 10,098 | 0.0113 | 0 |
| 8. max28_bias DW M | Max age 28, fix growth and bias adjustment | 3.3 | 10,096 | 0.0003 | 0 |

Table 32 Continued. Summary of key model building steps towards the SEDAR75 Base Model for Gulf of Mexico Gray Snapper and associated convergence diagnostics. Note that steps within each model progression are not shown due to the vast number of intermediate runs conducted.

| Model Name | Ln(R0) | Sigma R | Virgin SSB (mt) | Virgin Recr (1000s) | Depletion End Yr |
|-----------------------------|--------|---------|-----------------|---------------------|------------------|
| 0. S51 3.24 | 9.3 | 0.90 | 22,200 | 10,683 | 0.21 |
| 1. Convert to 3.30.17 | 9.3 | 0.90 | 22,133 | 10,651 | 0.20 |
| 2. Corrected COM data | 9.4 | 0.90 | 24,373 | 11,728 | 0.23 |
| 3. TY 2020 Rec | 9.6 | 0.47 | 30,586 | 14,739 | 0.18 |
| 4. TY 2020 - new COM fleets | 9.7 | 0.58 | 33,835 | 16,305 | 0.25 |
| 5. FI Length comps | 10.0 | 0.63 | 46,435 | 21,703 | 0.33 |
| 6. All Age comps added | 10.2 | 0.31 | 23,576 | 26,548 | 0.58 |
| 7. max age 21 fix growth | 10.0 | 0.32 | 21,246 | 22,870 | 0.48 |
| 8. max28 bias DW M | 10.1 | 0.38 | 21,719 | 23,190 | 0.48 |

Table 33. Settings used for Gulf of Mexico Gray Snapper projections.

| Parameter | Value | Comment |
|------------------------------|---|--|
| Relative F | Average from 2018 - 2020 | Average relative fishing mortality (apical F) over terminal three years of model |
| Selectivity | 2020 | Fleet specific selectivity estimated in the terminal year of the model |
| Retention | 2020 | Fleet specific retention estimated in the terminal year of the model |
| Recruitment | Beverton-Holt stock-recruitment relationship | Derived from the model estimated Beverton-Holt stock-recruitment relationship |
| Interim Landings (2021-2023) | 75.27/75.27/75.27 mt (Comm. Vertical Line) | For 2021-2023, used 3-year average of landings (2018-2020) |
| | 5.61/5.61/5.61 mt (Comm. Longline) | |
| | 0.11/0.11/0.11 mt (Comm. Nets & Traps) | |
| | 2091.08/2091.08/2091.08 thousands of fish (Private) | |
| | 1284.62/1284.62/1284.62 thousands of fish (Shore) | |
| | 340.86/340.86/340.86 thousands of fish (Charter & Headboat) | |

Table 34. Summary of Magnuson-Stevens Reauthorization Act benchmarks and reference points for the SEDAR75 Gulf of Mexico Gray Snapper assessment. Spawning Stock Biomass (SSB) is in metric tons, whereas F is a harvest rate (total biomass killed age 2+ / total biomass age 2+).

| Variable | Definition | Value |
|--------------------------------|--|--------|
| Base M | Target M for fully selected ages of Lorenzen Natural Mortality (M) | 0.15 |
| Steepness | Steepness of the Beverton-Holt stock recruit relationship (fixed) | 0.99 |
| R0 | Virgin Recruitment (1000s) | 23,191 |
| Generation Time | Fecundity-weighted mean age | 12.37 |
| SSB Unfished | Virgin spawning stock biomass (mt) | 21,719 |
| Mortality Rate Criteria | | |
| FMSY proxy | Equilibrium F that achieves 30%SPR | 0.134 |
| MFMT | $F_{MSYproxy}$ | 0.134 |
| FOY | $0.75 * \text{Directed F at } F_{30\%SPR}$ | 0.088 |
| Fcurrent | Geometric mean of the last 3 years of the assessment (F2018-2020) | 0.091 |
| Fcurrent/MFMT | Current stock status based on MFMT | 0.659 |
| Biomass Criteria | | |
| SSBMSYproxy | Equilibrium SSB at F30%SPR | 6477 |
| MSST | $0.5 * SSB_{30\%SPR}$ | 3,239 |
| SSB at Optimum Yield | Equilibrium SSB when Directed F = $0.75 * \text{Directed F at } F_{30\%SPR}$ | 7,907 |
| SSBcurrent | SSB in 2020 | 10,345 |
| SSBcurrent/SSBFMSYproxy | Current stock status based on SSB30%SPR (Equil) | 1.6 |
| SSB_2020/MSST | Current stock status based on MSST | 3.2 |
| SSBcurrent/SSB0 | SSB ratio in 2020 | 0.48 |

Table 35. Time series of fishing mortality and SSB relative to associated biological reference points. SSB is in metric tons, whereas F is a harvest rate (total biomass killed age 2+ / total biomass age 2+). Reference points include $F_{\text{FSR30}} = 0.134$, $\text{SSB}_{\text{FSR30}} = 6,477$ metric tons, and $\text{MSST}_{\text{FSR30}} = 3,238$ metric tons which was calculated as $(0.5) * \text{SSB}_{\text{FSR30}}$. SSB ratio was calculated as annual SSB divided by SSB_0 where $\text{SSB}_0 = 21,718$ metric tons.

| Year | F | F/FSR30 | SSB | SSB/ SSBFSR30 | SSB/MSST | SSB/SSB0 |
|------|-------|---------|--------|------------------|----------|----------|
| 1945 | 0.002 | 0.015 | 21,718 | 3.4 | 6.7 | 1.00 |
| 1946 | 0.003 | 0.022 | 21,669 | 3.3 | 6.7 | 1.00 |
| 1947 | 0.004 | 0.030 | 21,595 | 3.3 | 6.7 | 0.99 |
| 1948 | 0.005 | 0.037 | 21,496 | 3.3 | 6.6 | 0.99 |
| 1949 | 0.006 | 0.045 | 21,373 | 3.3 | 6.6 | 0.98 |
| 1950 | 0.007 | 0.052 | 21,227 | 3.3 | 6.6 | 0.98 |
| 1951 | 0.008 | 0.060 | 21,059 | 3.3 | 6.5 | 0.97 |
| 1952 | 0.008 | 0.060 | 20,871 | 3.2 | 6.4 | 0.96 |
| 1953 | 0.009 | 0.067 | 20,676 | 3.2 | 6.4 | 0.95 |
| 1954 | 0.010 | 0.075 | 20,469 | 3.2 | 6.3 | 0.94 |
| 1955 | 0.009 | 0.067 | 20,251 | 3.1 | 6.3 | 0.93 |
| 1956 | 0.010 | 0.075 | 20,086 | 3.1 | 6.2 | 0.92 |
| 1957 | 0.011 | 0.082 | 19,917 | 3.1 | 6.2 | 0.92 |
| 1958 | 0.012 | 0.090 | 19,734 | 3.0 | 6.1 | 0.91 |
| 1959 | 0.013 | 0.097 | 19,538 | 3.0 | 6.0 | 0.90 |
| 1960 | 0.014 | 0.105 | 19,346 | 3.0 | 6.0 | 0.89 |
| 1961 | 0.014 | 0.105 | 19,137 | 3.0 | 5.9 | 0.88 |
| 1962 | 0.015 | 0.112 | 18,889 | 2.9 | 5.8 | 0.87 |
| 1963 | 0.015 | 0.112 | 18,640 | 2.9 | 5.8 | 0.86 |
| 1964 | 0.016 | 0.120 | 18,489 | 2.9 | 5.7 | 0.85 |
| 1965 | 0.017 | 0.127 | 18,324 | 2.8 | 5.7 | 0.84 |
| 1966 | 0.017 | 0.127 | 18,009 | 2.8 | 5.6 | 0.83 |
| 1967 | 0.019 | 0.142 | 17,653 | 2.7 | 5.5 | 0.81 |
| 1968 | 0.021 | 0.157 | 17,236 | 2.7 | 5.3 | 0.79 |
| 1969 | 0.022 | 0.165 | 16,784 | 2.6 | 5.2 | 0.77 |
| 1970 | 0.023 | 0.172 | 16,249 | 2.5 | 5.0 | 0.75 |
| 1971 | 0.026 | 0.195 | 15,689 | 2.4 | 4.8 | 0.72 |

Table 35 Continued. Time series of fishing mortality and SSB relative to associated biological reference points. SSB is in metric tons, whereas F is a harvest rate (total biomass killed age 2+ / total biomass age 2+). Reference points include $F_{\text{FSR30}} = 0.134$, $\text{SSB}_{\text{FSR30}} = 6,477$ metric tons, and $\text{MSST}_{\text{FSR30}} = 3,238$ metric tons which was calculated as $(0.5) * \text{SSB}_{\text{FSR30}}$. SSB ratio was calculated as annual SSB divided by SSB_0 where $\text{SSB}_0 = 21,718$ metric tons.

| Year | F | F/FSR30 | SSB | SSB/ SSBFSR30 | SSB/MSST | SSB/SSB0 |
|------|-------|---------|--------|------------------|----------|----------|
| 1972 | 0.029 | 0.22 | 15,112 | 2.33 | 4.7 | 0.70 |
| 1973 | 0.031 | 0.23 | 14,578 | 2.25 | 4.5 | 0.67 |
| 1974 | 0.034 | 0.25 | 14,119 | 2.18 | 4.4 | 0.65 |
| 1975 | 0.035 | 0.26 | 13,692 | 2.11 | 4.2 | 0.63 |
| 1976 | 0.038 | 0.28 | 13,321 | 2.06 | 4.1 | 0.61 |
| 1977 | 0.040 | 0.30 | 12,879 | 1.99 | 4.0 | 0.59 |
| 1978 | 0.043 | 0.32 | 12,410 | 1.92 | 3.8 | 0.57 |
| 1979 | 0.046 | 0.34 | 11,870 | 1.83 | 3.7 | 0.55 |
| 1980 | 0.049 | 0.37 | 11,281 | 1.74 | 3.5 | 0.52 |
| 1981 | 0.069 | 0.52 | 10,608 | 1.64 | 3.3 | 0.49 |
| 1982 | 0.083 | 0.62 | 9,724 | 1.50 | 3.0 | 0.45 |
| 1983 | 0.053 | 0.40 | 9,102 | 1.40 | 2.8 | 0.42 |
| 1984 | 0.073 | 0.55 | 8,868 | 1.37 | 2.7 | 0.41 |
| 1985 | 0.042 | 0.31 | 8,478 | 1.31 | 2.6 | 0.39 |
| 1986 | 0.045 | 0.34 | 8,529 | 1.32 | 2.6 | 0.39 |
| 1987 | 0.059 | 0.44 | 8,723 | 1.35 | 2.7 | 0.40 |
| 1988 | 0.051 | 0.38 | 8,431 | 1.30 | 2.6 | 0.39 |
| 1989 | 0.082 | 0.61 | 8,291 | 1.28 | 2.6 | 0.38 |
| 1990 | 0.084 | 0.63 | 8,272 | 1.28 | 2.6 | 0.38 |
| 1991 | 0.133 | 0.99 | 7,697 | 1.19 | 2.4 | 0.35 |
| 1992 | 0.088 | 0.66 | 7,158 | 1.10 | 2.2 | 0.33 |
| 1993 | 0.097 | 0.73 | 7,273 | 1.12 | 2.2 | 0.34 |
| 1994 | 0.088 | 0.66 | 7,131 | 1.10 | 2.2 | 0.33 |
| 1995 | 0.101 | 0.76 | 6,897 | 1.06 | 2.1 | 0.32 |
| 1996 | 0.097 | 0.73 | 6,612 | 1.02 | 2.0 | 0.30 |
| 1997 | 0.084 | 0.63 | 6,324 | 0.98 | 2.0 | 0.29 |
| 1998 | 0.104 | 0.78 | 6,415 | 0.99 | 2.0 | 0.29 |

Table 35 Continued. Time series of fishing mortality and SSB relative to associated biological reference points. SSB is in metric tons, whereas F is a harvest rate (total biomass killed age 2+ / total biomass age 2+). Reference points include $F_{\text{FSR30\%}} = 0.134$, $\text{SSB}_{\text{FSR30}} = 6,477$ metric tons, and $\text{MSST}_{\text{FSR30}} = 3,238$ metric tons which was calculated as $(0.5) * \text{SSB}_{\text{FSR30}}$. SSB ratio was calculated as annual SSB divided by SSB_0 where $\text{SSB}_0 = 21,718$ metric tons.

| Year | F | F/FSR30 | SSB | SSB/ SSBFSR30 | SSB/MSST | SSB/SSB0 |
|------|-------|---------|--------|------------------|----------|----------|
| 1999 | 0.082 | 0.61 | 6,582 | 1.0 | 2.0 | 0.30 |
| 2000 | 0.075 | 0.56 | 6,609 | 1.0 | 2.0 | 0.30 |
| 2001 | 0.090 | 0.67 | 6,773 | 1.0 | 2.1 | 0.31 |
| 2002 | 0.076 | 0.57 | 6,561 | 1.0 | 2.0 | 0.30 |
| 2003 | 0.098 | 0.73 | 6,834 | 1.1 | 2.1 | 0.32 |
| 2004 | 0.114 | 0.85 | 6,859 | 1.1 | 2.1 | 0.32 |
| 2005 | 0.089 | 0.67 | 6,856 | 1.1 | 2.1 | 0.32 |
| 2006 | 0.073 | 0.55 | 7,075 | 1.1 | 2.2 | 0.33 |
| 2007 | 0.078 | 0.58 | 7,470 | 1.2 | 2.3 | 0.34 |
| 2008 | 0.102 | 0.76 | 8,066 | 1.2 | 2.5 | 0.37 |
| 2009 | 0.105 | 0.79 | 8,004 | 1.2 | 2.5 | 0.37 |
| 2010 | 0.055 | 0.41 | 7,794 | 1.2 | 2.4 | 0.36 |
| 2011 | 0.063 | 0.47 | 8,053 | 1.2 | 2.5 | 0.37 |
| 2012 | 0.115 | 0.86 | 8,366 | 1.3 | 2.6 | 0.38 |
| 2013 | 0.104 | 0.78 | 8,532 | 1.3 | 2.6 | 0.39 |
| 2014 | 0.118 | 0.88 | 8,973 | 1.4 | 2.8 | 0.41 |
| 2015 | 0.105 | 0.79 | 8,587 | 1.3 | 2.7 | 0.40 |
| 2016 | 0.117 | 0.88 | 8,840 | 1.4 | 2.7 | 0.41 |
| 2017 | 0.098 | 0.73 | 8,663 | 1.3 | 2.7 | 0.40 |
| 2018 | 0.093 | 0.70 | 9,055 | 1.4 | 2.8 | 0.42 |
| 2019 | 0.091 | 0.68 | 9,458 | 1.5 | 2.9 | 0.43 |
| 2020 | 0.088 | 0.66 | 10,345 | 1.6 | 3.2 | 0.48 |

Table 36. Results of the OFL projections (fishing set at F_{SPR30}) for Gulf of Mexico Gray Snapper. Recruitment (R) is in 1000s of age-0 fish, SSB is in metric tons, F is a harvest rate (total biomass killed age 2+ / total biomass age 2+), and OFL is the overfishing limit in millions of pounds gutted weight. Reference points include $F_{SPR30} = 0.134$, $SSB_{SPR30} = 6,477$ metric tons, and $MSST = 3,238$ metric tons which was calculated as $0.5 * SSB_{SPR30}$. SSB ratio was calculated as annual SSB divided by SSB_0 .

| Year | R | F | F/ F_{SPR30} | SSB | SSB/ SSB_{SPR30} | SSB/ $MSST$ | SSB/ SSB_0 | OFL |
|------|--------|------|----------------|--------|-----------------------|-------------|--------------|-------|
| 2024 | 23,143 | 0.13 | 1 | 11,963 | 1.8 | 3.7 | 0.55 | 7.970 |
| 2025 | 23,133 | 0.13 | 1 | 10,962 | 1.7 | 3.4 | 0.50 | 7.348 |
| 2026 | 23,122 | 0.13 | 1 | 10,031 | 1.5 | 3.1 | 0.46 | 6.746 |
| 2027 | 23,111 | 0.13 | 1 | 9,216 | 1.4 | 2.8 | 0.42 | 6.202 |
| 2028 | 23,100 | 0.13 | 1 | 8,533 | 1.3 | 2.6 | 0.39 | 5.735 |

Table 37. Results of the ABC projections (directed $F = 0.75 * F_{SPR30}$) for Gulf of Mexico Gray Snapper. Recruitment (R) is in 1000s of age-0 fish, SSB is in metric tons, F is a harvest rate (total biomass killed age 2+ / total biomass age 2+), and OFL is the overfishing limit in millions of pounds gutted weight. Reference points include $F_{SPR30} = 0.134$, $SSB_{SPR30} = 6,477$ metric tons, and $MSST = 3,238$ metric tons which was calculated as $0.5 * SSB_{SPR30}$. SSB ratio was calculated as annual SSB divided by SSB_0 .

| Year | R | F | F/ F_{SPR30} | SSB | SSB/ SSB_{SPR30} | SSB/ $MSST$ | SSB/ SSB_0 | ABC |
|------|--------|-----|----------------|--------|-----------------------|-------------|--------------|-------|
| 2024 | 23,143 | 0.1 | 0.75 | 11,963 | 1.5 | 3.7 | 0.55 | 5.978 |
| 2025 | 23,138 | 0.1 | 0.75 | 11,446 | 1.4 | 3.5 | 0.53 | 5.760 |
| 2026 | 23,132 | 0.1 | 0.75 | 10,906 | 1.4 | 3.4 | 0.50 | 5.514 |
| 2027 | 23,127 | 0.1 | 0.75 | 10,387 | 1.3 | 3.2 | 0.48 | 5.267 |
| 2028 | 23,121 | 0.1 | 0.75 | 9,912 | 1.3 | 3.1 | 0.46 | 5.035 |

Figures




Figure 1. Data sources used in the Gulf of Mexico Gray Snapper Stock Synthesis assessment model.




Figure 2. Mean weight-at-length (top panel), recommended and estimated growth curves (with 95% confidence intervals; middle panel), and natural mortality (bottom panel) used in the assessment model for Gulf of Mexico Gray Snapper. SEDAR51 and SEDAR75 inputs are presented for comparison.




Figure 3. Updated estimates of maturity function using functional maturity.




Figure 4. Commercial landing discrepancy between SEDAR51 Data Workshop and SEDAR75 Assessment




Figure 5. Gulf of Mexico Gray Snapper observed landings by fishery for SEDAR75 and SEDAR51. Commercial and recreational landings are in metric tons and numbers of fish, respectively.




Figure 6. Gulf of Mexico Gray Snapper observed commercial discards by fishery for SEDAR75. Commercial discards in numbers of fish.




Figure 7. Gulf of Mexico Gray Snapper observed commercial discards by fishery for SEDAR75. Commercial discards in numbers of fish. Colors match the legend in Figure 6.




Figure 8. Gulf of Mexico Gray Snapper observed recreational discards by fishery for SEDAR75. Recreational discards in numbers of fish.




Figure 9. Gulf of Mexico Gray Snapper observed recreational discards by fishery for SEDAR75. Recreational discards in numbers of fish. Colors match the legend in Figure 8.




Figure 10. Gulf of Mexico Gray Snapper observed discards by fishery for SEDAR75 and SEDAR51. Recreational discards are proportion of discards.




Figure 11. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Commercial Vertical Line fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 12. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 13. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Commercial Nets & Traps fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 14. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Recreational Private fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.





Figure 15. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Recreational Shore fishery. Input sample sizes (N input) and adjusted sample sizes (N adj) estimated by Stock Synthesis are also reported.



Proportion

Length (cm)

Figure 15 Continued. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Recreational Shore fishery. Input sample sizes (N input) and adjusted sample sizes (N adj) estimated by Stock Synthesis are also reported.




Figure 16. Observed length composition data (retained) of Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 17. Observed age composition data (retained) of Gulf of Mexico Gray Snapper in the Commercial Vertical Line fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 18. Observed age composition data (retained) of Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 19. Observed age composition data (retained) of Gulf of Mexico Gray Snapper in the Recreational Private fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 20. Observed age composition data (retained) of Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Input sample sizes (N_{input}) and adjusted sample sizes (N_{adj}) estimated by Stock Synthesis are also reported.




Figure 21. Observed relative age proportions in each year for Gulf of Mexico Gray Snapper in the Commercial Vertical Line fishery. Cohort progressions are evident.




Figure 22. Observed relative age proportions in each year for Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Cohort progressions are evident.




Figure 23. Observed relative age proportions in each year for Gulf of Mexico Gray Snapper in the Recreational Private fishery. Cohort progressions are evident.




Figure 24. Observed relative age proportions in each year for Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Cohort progressions are evident.




Figure 25. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the recreational private fishery. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 26. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the recreational shore fishery. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 27. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the FWRI Age-0 Survey. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 28. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the FWRI Age-1 Survey. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 29. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the SEAMAP Trawl. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 30. Observed length composition data of Gulf of Mexico Gray Snapper from the SEAMAP Trawl.




Figure 31. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the Combined Video Survey. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 32. Observed length composition data of Gulf of Mexico Gray Snapper from the Combined Video Survey.




Figure 33. Standardized index of relative abundance and associated 95% uncertainty interval around index values based on the model assumption of lognormal error for Gulf of Mexico Gray Snapper from the RF Visual Survey. The uncertainty displayed includes the additional SE parameter estimated as part of the data weighting process.




Figure 34. Observed length composition data of Gulf of Mexico Gray Snapper from the RF Visual Survey.




Figure 35. Catch curve analysis from the aggregated Commercial Vertical Line (top panel) and Longline (bottom panel) data. The gray dot reflects the first age fully selected for by the gear.




Figure 35 Continued. Catch curve analysis from the aggregated Commercial Vertical Line (top panel) and Longline (bottom panel) data. The gray dot reflects the first age fully selected for by the gear.



SEDAR75**SEDAR51**

Figure 36. Annual exploitation rate estimates (total biomass killed age 2+ / total biomass age 2+) for Gulf of Mexico Gray Snapper.



SEDAR75**SEDAR51**

Figure 37. Annual exploitation rate (total biomass killed age 2+ / total biomass age 2+) by fleet for Gulf of Mexico Gray Snapper.




Figure 38. Length-based selectivity for each fleet for Gulf of Mexico Gray Snapper in the terminal year of the assessment (given in parentheses). Dashed horizontal line indicates 50%, whereas the dashed vertical lines identify lengths in 25 cm FL intervals.



SEDAR75**SEDAR51**

Figure 39. Length-based selectivity for the Commercial Vertical Line fishery. Selectivity (blue line) is constant over the entire assessment time period (1945 - 2020). Retention (red line) is shown for the most recent time period. Discard mortality (orange line) is constant at 0.14.



SEDAR75**SEDAR51**

Figure 40. Length-based selectivity for the Commercial Longline fishery. Selectivity (blue line) is constant over the entire assessment time period (1945 - 2020). Retention (red line) is shown for the most recent time period. Discard mortality (orange line) is constant at 0.14.




SEDAR75**SEDAR51**

Figure 41. Length-based selectivity for the Commercial Nets and Traps fishery. Selectivity (blue line) is constant over the entire assessment time period (1945 - 2020). SEDAR 51 selectivity is for the non Monroe County selectivity.

SEDAR75



SEDAR51





Figure 42. Length-based selectivity for the Recreational Private fishery. Selectivity (blue line) is constant over the entire assessment time period (1945 - 2020). Retention (red line) is shown for the most recent time period. Discard mortality (orange line) is constant at 0.069.

SEDAR75



SEDAR51





Figure 43. Length-based selectivity for the Recreational Shore fishery. Selectivity (blue line) is constant over the entire assessment time period (1945 - 2020). Retention (red line) is shown for the most recent time period. Discard mortality (orange line) is constant at 0.069.

SEDAR75



SEDAR51




Figure 44. Length-based selectivity for the Recreational Charter & Headboat fishery. Selectivity (blue line) is constant over the entire assessment time period (1945 - 2020). Retention (red line) is shown for the most recent time period. Discard mortality (orange line) is constant at 0.069.




Figure 45. Derived age-based selectivity for each fleet for Gulf of Mexico Gray Snapper in the terminal year of the assessment (given in parentheses). Dashed horizontal line indicates 50%, whereas the dashed vertical lines identify ages in 2 year intervals. No age compositions were used in SEDAR51, resulting in the flat selectivity for SEDAR51.




Figure 46. Length-based selectivity for each survey for Gulf of Mexico Gray Snapper in the terminal year of the assessment (given in parentheses). Dashed horizontal line indicates 50%, whereas the dashed vertical lines identify lengths in 25 cm FL intervals.




Figure 47. Derived age-based selectivity for each survey for Gulf of Mexico Gray Snapper in the terminal year of the assessment (given in parentheses). Dashed horizontal line indicates 50%, whereas the dashed vertical lines identify ages in 2 year intervals. SEDAR 51 did not include age compositions.




Figure 48. Time-varying retention functions for the Commercial Vertical Line fishery for Gulf of Mexico Gray Snapper from SEDAR75 and SEDAR51.




Figure 49. Time-varying retention functions for the Commercial Longline fishery for Gulf of Mexico Gray Snapper from SEDAR75 and SEDAR51.




Figure 50. Time-varying retention functions for the Recreational Private fishery for Gulf of Mexico Gray Snapper from SEDAR75 and SEDAR51.




Figure 51. Time-varying retention functions for the Recreational Shore fishery for Gulf of Mexico Gray Snapper from SEDAR75 and SEDAR51.




Figure 52. Time-varying retention functions for the Recreational Charter & Headboat fishery for Gulf of Mexico Gray Snapper from SEDAR75 and SEDAR51.

SEDAR75




Figure 53. Predicted stock-recruitment relationship for Gulf of Mexico Gray Snapper for SEDAR75 (steepness was fixed at 0.99). Plotted are predicted annual recruitments from Stock Synthesis (circles), expected recruitment from the stock-recruit relationship (black line), and bias adjusted recruitment from the stock-recruit relationship (dashed line).

SEDAR51




Figure 53 Continued. Predicted stock-recruitment relationship for Gulf of Mexico Gray Snapper SEDAR51 (steepness 0.99). Plotted are predicted annual recruitments from Stock Synthesis (circles), expected recruitment from the stock-recruit relationship (black line), and bias adjusted recruitment from the stock-recruit relationship (dashed line).



SEDAR75**SEDAR51**

Figure 54. Top panel: Estimated Age-0 recruitment with 95% confidence intervals for Gulf of Mexico Gray Snapper (steepness was fixed at 0.99 for SEDAR75). Bottom panel: Estimated Age-0 recruitment with 95% confidence intervals for Gulf of Mexico Gray Snapper (steepness fixed at 0.99).



SEDAR75**SEDAR51**

Figure 55. Top panel: Estimated log recruitment deviations for Gulf of Mexico Gray Snapper (steepness fixed at 0.99). Bottom panel: Estimated log recruitment deviations for Gulf of Mexico Gray Snapper steepness fixed at 0.99).



SEDAR75**SEDAR51**

Figure 56. Asymptotic standard errors for recruitment deviations for Gulf of Mexico Gray Snapper. The red lines represent the estimated values of SigmaR.




Figure 57. Points are transformed variances. Red line shows current settings for bias adjustment specified for the Base Run, which coincides with the least squares estimate of alternative bias adjustment relationship for recruitment deviations (dashed orange line). For more information, see Methot and Taylor 2011.




Figure 58. Estimate of total biomass (in 1000s of metric tons) for Gulf of Mexico Gray Snapper.




Figure 59. Estimate of spawning stock biomass (in 1000s of metric tons) and associated 95% confidence intervals for Gulf of Mexico Gray Snapper.




Figure 60. Gulf of Mexico Gray Snapper observed and expected landings by fishery for SEDAR75 (left panels) and SEDAR51 (right panels). Commercial and recreational landings are in metric tons and numbers of fish, respectively. Dashed vertical lines identify ten-year intervals.




Figure 61. Input (dots with 95% confidence intervals) and expected (blue lines) discards by the Commercial Vertical Line for Gulf of Mexico Gray Snapper. Discards are in numbers of fish (1,000s) and reflect released fish (i.e., before discard mortality has been applied).




Figure 62. Input (dots with 95% confidence intervals) and expected (blue lines) discards by the Commercial Longline for Gulf of Mexico Gray Snapper. Discards are in numbers of fish (1,000s) and reflect released fish (i.e., before discard mortality has been applied).




Figure 63. Input (dots with 95% confidence intervals) and expected (blue lines) discards by the Commercial Nets & Traps for Gulf of Mexico Gray Snapper. Discards are in numbers of fish (1,000s) and reflect released fish (i.e., before discard mortality has been applied).




Figure 64. Input (dots with 95% confidence intervals) and expected (blue lines) discards by the Recreational Private for Gulf of Mexico Gray Snapper. Discards are in numbers of fish (1,000s) and reflect released fish (i.e., before discard mortality has been applied).




Figure 65. Input (dots with 95% confidence intervals) and expected (blue lines) discards by the Recreational Shore for Gulf of Mexico Gray Snapper. Discards are in numbers of fish (1,000s) and reflect released fish (i.e., before discard mortality has been applied).




Figure 66. Gulf of Mexico Gray Snapper observed and expected discards by fishery for SEDAR75. Commercial and recreational discards are in numbers of fish, respectively. Dashed vertical lines identify five-year intervals.




Figure 67. Gulf of Mexico Gray Snapper observed and expected indices for SEDAR75 (left panels) and SEDAR51 (right panels). Dashed vertical lines identify five-year intervals. The root mean squared error (RMSE) is also provided.




Figure 68. Gulf of Mexico Gray Snapper observed and expected indices for SEDAR75 (left panels) and SEDAR51 (right panels). Dashed vertical lines identify five-year intervals. The root mean squared error (RMSE) is also provided.

SEDAR75




Figure 69. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Vertical Line fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR51




Figure 69 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Vertical Line fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size. This fleet in SEDAR represents the Monroe County Handline fleet.

SEDAR75




Figure 70. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR51




Figure 70 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size. This fleet in SEDAR represents the non-Monroe County Handline fleet.

SEDAR75




Figure 70 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size. This fleet in SEDAR represents the non-Monroe County Handline fleet.

SEDAR75




Figure 71. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Nets & Traps fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR51





Figure 71 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Nets & Traps fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size. This fleet in SEDAR represents the Longline Fleet.

SEDAR75

Proportion

Length (cm)

Figure 71 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Nets & Traps fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size. This fleet in SEDAR represents the Longline Fleet

SEDAR75




Figure 72. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Private fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR51




Figure 72 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Private fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR75




Figure 72 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Private fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR75




Figure 73. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Shore fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.


SEDAR51

Figure 73 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Shore fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR75




Figure 73 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Shore fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR51




Figure 73 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Shore fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR75




Figure 74. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR51




Figure 74 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR75




Figure 74 Continued. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.

SEDAR75




Figure 75. Model fits to the length composition of discarded or retained catch aggregated across years within a given fleet for Gulf of Mexico Gray Snapper. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR72, 'Sum of N input' is the total input sample size and 'Sum of N adj.' is the total sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'Sum of N adj.' is the input sample size.

SEDAR51




Figure 75 Continued. Model fits to the length composition of discarded or retained catch aggregated across years within a given fleet for Gulf of Mexico Gray Snapper. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR72, 'Sum of N input' is the total input sample size and 'Sum of N adj.' is the total sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'Sum of N adj.' is the input sample size.

SEDAR75




Figure 76. Pearson residuals for discard and retained length composition data by year compared across fleets and surveys for Gulf of Mexico Gray Snapper for SEDAR75. Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).

SEDAR51




Figure 76 Continued. Pearson residuals for discard and retained length composition data by year compared across fleets and surveys for Gulf of Mexico Gray Snapper for SEDAR51. Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).

SEDAR75




Figure 76 Continued. Pearson residuals for discard and retained length composition data by year compared across fleets and surveys for Gulf of Mexico Gray Snapper for SEDAR75. Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 77. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the SEAMAP Trawl fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, length compositions were not used for this survey.




Figure 78. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the Combined Video Survey fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, length compositions were not used for this survey.




Figure 79. Observed and predicted length compositions (retained) for Gulf of Mexico Gray Snapper in the RF Visual Survey fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, length compositions were not used for this survey.




Figure 80. Observed and predicted age compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Vertical Line fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. Age compositions were not used in SEDAR51.




Figure 81. Observed and predicted age compositions (retained) for Gulf of Mexico Gray Snapper in the Commercial Longline fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample.




Figure 82. Observed and predicted age compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Private fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. For the SEDAR51, 'N adj.' is the input sample size.




Figure 83. Observed and predicted age compositions (retained) for Gulf of Mexico Gray Snapper in the Recreational Charter & Headboat fishery. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'N input' is the input sample size and 'N adj.' is the sample size after adjustment by the Dirichlet-Multinomial parameter. Age compositions were not used in SEDAR51.




Figure 84. Model fits to the age composition of retained catch aggregated across years within a given fleet for Gulf of Mexico Gray Snapper. Green lines represent predicted length compositions, while grey shaded regions represent observed length compositions. For SEDAR75, 'Sum of N input' is the total input sample size and 'Sum of N adj.' is the total sample size after adjustment by the Dirichlet-Multinomial parameter.




Figure 85. Pearson residuals for retained age composition data by year compared across fleets for Gulf of Mexico Gray Snapper for SEDAR75. Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).




Figure 86. The profile likelihood for the natural log of the unfished recruitment parameter of the Beverton – Holt stock-recruit function for Gulf of Mexico Gray Snapper. Each line represents the change in negative log-likelihood value for each of the data sources fit in the model across the range of fixed steepness values tested in the profile diagnostic run. The MLE for the base model was 10.052. The bottom panel shows a close up of the top panel to better detect significant differences between runs.




Figure 87. Results of the jitter analysis for various likelihood components for the Gulf of Mexico Gray Snapper Base Model. Each panel gives the results of 100 model runs where the starting parameter values for each run were randomly changed ('jittered') by 10% from the base model best fit values. The Base Run value for each panel is indicated by a red line.




Figure 88. Retrospective analysis of spawning stock biomass (top panels) and fishing mortality (F, bottom panels) estimates for Gulf of Mexico Gray Snapper conducted by re-fitting each reference model (Ref) after removing five years of observations, one year at a time sequentially. The retrospective results are shown for the entire time series and for the most recent years only. Mohn's rho statistic and the corresponding 'hindcast rho' values (in brackets) are printed at the top of each panel. One-year-ahead projections denoted by color-coded dashed lines with terminal points shown for each model. Grey shaded areas are the 95% confidence intervals from the reference model. See Carvalho et al. (2021) for additional details.




Figure 89. Joint residual plots for indices of abundance fits for Gulf of Mexico Gray Snapper. Vertical lines with points show the residuals (in colors by index), and solid black line reflects the loess smoother through all the residuals. Boxplots indicate the median and quantiles in cases where residuals from the multiple indices are available for any given year. Root-mean squared errors (RMSE) are included in the upper right-hand corner of each plot. See Carvalho et al. (2021) for additional details.




Figure 90. Joint residual plots for annual mean age estimates for Gulf of Mexico Gray Snapper. Vertical lines with points show the residuals (in colors by index), and solid black line reflects the loess smoother through all the residuals. Boxplots indicate the median and quantiles in cases where residuals from the multiple indices are available for any given year. Root-mean squared errors (RMSE) are included in the upper right-hand corner of each plot. See Carvalho et al. (2021) for additional details.




Figure 91. Joint residual plots for annual mean length estimates for Gulf of Mexico Gray Snapper. Vertical lines with points show the residuals (in colors by index), and solid black line reflects the loess smoother through all the residuals. Boxplots indicate the median and quantiles in cases where residuals from the multiple indices are available for any given year. Root-mean squared errors (RMSE) are included in the upper right-hand corner of each plot. See Carvalho et al. (2021) for additional details.




Figure 92. Hindcasting cross-validation (HCxval) results for indices of abundance fits for Gulf of Mexico Gray Snapper. Shown are observed (large points connected with dashed line), fitted (solid lines) and one-year ahead forecast values (small terminal points). HCxval was performed using one reference model (Ref) and five hindcast model runs (solid lines) relative to the expected index. The observations used for cross validation are highlighted as color-coded solid circles with associated 95% confidence intervals (light-grey shading). The model reference year refers to the endpoints of each one-year-ahead forecast and the corresponding observation (i.e., year of peel + 1). The mean absolute scaled error (MASE) score associated with each index time series is denoted in each panel. See Carvalho et al. (2021) for additional details.




Figure 93. Hindcasting cross-validation (HCxval) results for fits to annual mean age estimates for Gulf of Mexico Gray Snapper. Shown are observed (large points connected with dashed line), fitted (solid lines) and one-year ahead forecast values (small terminal points). HCxval was performed using one reference model (Ref) and five hindcast model runs (solid lines) relative to the expected mean age. The observations used for cross-validation are highlighted as color-coded solid circles with associated 95% confidence intervals (light-grey shading). The model reference year refers to the endpoints of each one-year-ahead forecast and the corresponding observation (i.e., year of peel + 1). The mean absolute scaled error (MASE) score associated with each age composition time series is denoted in each panel. See Carvalho et al. (2021) for additional details.




Figure 94. Hindcasting cross-validation (HCxval) results for fits to annual mean length estimates for Gulf of Mexico Gray Snapper. Shown are observed (large points connected with dashed line), fitted (solid lines) and one-year ahead forecast values (small terminal points). HCxval was performed using one reference model (Ref) and five hindcast model runs (solid lines) relative to the expected mean length. The observations used for cross-validation are highlighted as color-coded solid circles with associated 95% confidence intervals (light-grey shading). The model reference year refers to the endpoints of each one-year-ahead forecast and the corresponding observation (i.e., year of peel + 1). The mean absolute scaled error (MASE) score associated with each size composition time series is denoted in each panel. See Carvalho et al. (2021) for additional details.




Figure 95. Runs tests results for indices of abundance for Gulf of Mexico Gray Snapper. Green shading indicates no evidence ($p \geq 0.05$) and red shading evidence ($p < 0.05$) to reject the hypothesis of a randomly distributed time-series of residuals, respectively. The shaded (green/red) area spans three residual standard deviations to either side from zero, and the red points outside of the shading violate the 'three-sigma limit' for that series. See Carvalho et al. (2021) for additional details.




Figure 96. Runs tests results for mean length for Gulf of Mexico Gray Snapper. Green shading indicates no evidence ($p \geq 0.05$) and red shading evidence ($p < 0.05$) to reject the hypothesis of a randomly distributed time-series of residuals, respectively. The shaded (green/red) area spans three residual standard deviations to either side from zero, and the red points outside of the shading violate the 'three-sigma limit' for that series. See Carvalho et al. (2021) for additional details.




Figure 97. Runs tests results for mean age for Gulf of Mexico Gray Snapper. Green shading indicates no evidence ($p \geq 0.05$) and red shading evidence ($p < 0.05$) to reject the hypothesis of a randomly distributed time-series of residuals, respectively. The shaded (green/red) area spans three residual standard deviations to either side from zero, and the red points outside of the shading violate the 'three-sigma limit' for that series. See Carvalho et al. (2021) for additional details.




Figure 98. Bridging analysis showing changes in estimates of SSB and associated uncertainty through each major step of model building between SEDAR51 and SEDAR75.




Figure 99. Bridging analysis showing changes in estimates of fraction unfished and associated uncertainty through each major step of model building between SEDAR51 and SEDAR75.




Figure 100. Bridging analysis showing changes in estimates of annual exploitation rates (total biomass killed age 2+ / total biomass age 2+) and associated uncertainty through each major step of model building between SEDAR51 and SEDAR75.




Figure 101. Bridging analysis showing changes in estimates of annual recruitment and associated uncertainty through each major step of model building between SEDAR51 and SEDAR75.




Figure 102. SEDAR51 base model run with correct commercial landings showing changes in estimates of SSB and associated uncertainty compared with SEDAR51.




Figure 103. Bridging analysis showing changes in estimates of SSB and associated uncertainty through each major step of model building between SEDAR51 and SEDAR75.




Figure 104. Differences in SSB estimates (top panel), annual exploitation rates (total biomass killed age2+ / total biomass age 2+) (middle panel) and annual recruitment (bottom panel), and associated uncertainty between the SEDAR75 Base Run and natural mortality sensitivity runs.




Figure 105. Time series of SSB and harvest rate (total biomass killed age 2+ / total biomass age 2+) with respect to status determination criteria for the SEDAR75 Gulf of Mexico Gray Snapper assessment.




Figure 106. Kobe plot illustrating the trajectory of stock status. The orange coloring indicates regions where the stock is below the biomass target but above the biomass threshold ($MSST = 0.5 \times SSB_{SPR30\%}$). The 2020 terminal year stock status is indicated by the gray dot.




Figure 107. Historic and forecasted yields for the OFL projections and ABC projections.

10. Appendix

A summary listing of all data sets included in the assessment, along with any revisions to the contact information for who provided the analysis, has been compiled below. This will be the source of data information for the next assessment.

| Primary Categories | Data Type | Contributing Organization | Data Providers | Contact Information |
|--------------------------|--|-----------------------------|---|---|
| Life History | Raw age and length data | FWRI | Meagan Schrandt | meagan.schrandt@myfwc.com |
| | Raw age and length data | SEFSC | Beverly Barnett Robert Allman | beverly.barnett@noaa.gov robert.allman@noaa.gov |
| | Raw age and length data | GulfFIN | Gregg Bray | gregg.bray@gsmfc.org |
| | Raw age and length data | SEFSC | Beverly Barnett Robert Allman Gregg Bray Meagan Schrandt Ed Kim | beverly.barnett@noaa.gov robert.allman@noaa.gov gregg.bray@gsmfc.org meagan.schrandt@myfwc.com edward.sm.kim1@gmail.com |
| | Raw age and maturity data | University of South Alabama | Ed Kim | edward.sm.kim1@gmail.com |
| | Age-growth | SEFSC | Steve Garner | steven.garner@noaa.gov |
| | Age-error | SEFSC | Steve Garner | steven.garner@noaa.gov |
| | Reproduction | SEFSC | Heather Moncrief | heather.moncrief-cox@noaa.gov |
| Fishery Dependent | Raw recreational headboat length | SEFSC | Ken Brennan | kenneth.brennan@noaa.gov |
| | Raw recreational length data | SEFSC | Matt Nuttall | matthew.nuttall@noaa.gov |
| | Raw length data | GulfFIN | Gregg Bray | gregg.bray@gsmfc.org |
| | Raw commercial length data | SEFSC | Larry Beerkircher | lawrence.r.beerkircher@noaa.gov |
| | Recreational catch (landings+discards) estimates | SEFSC | Matt Nuttall | matthew.nuttall@noaa.gov |
| | Recreational effort estimates | SEFSC | Matt Nuttall | matthew.nuttall@noaa.gov |
| | MRIP CVs | SEFSC | Matt Nuttall | matthew.nuttall@noaa.gov |

| | | | | |
|----------------------------|---|-------------|--|---|
| | Commercial landings estimates | SEFSC | Refik Orhun | refik.orhun@noaa.gov |
| | Recreational headboat catch (landings + discards) | SEFSC | Ken Brennan | kenneth.brennan@noaa.gov |
| | Commercial discard estimates | SEFSC | Kevin McCarthy Steve Smith | kevin.j.mccarthy@noaa.gov steven.smith@noaa.gov |
| | Commercial length compositions | SEFSC | Molly Stevens | molly.stevens@noaa.gov |
| | Recreational length composition | SEFSC | Molly Stevens | molly.stevens@noaa.gov |
| | Recreational age composition | SEFSC | Molly Stevens | molly.stevens@noaa.gov |
| | Commercial age composition | SEFSC | Molly Stevens | molly.stevens@noaa.gov |
| | Recreational Private Index | SEFSC | Francesca Forrestal | francesca.forrestal@noaa.gov |
| | Recreational Shore Mode Index | SEFSC | Francesca Forrestal | francesca.forrestal@noaa.gov |
| Fishery Independent | Trawl index + size frequency | SEFSC | Adam Pollack | adam.pollack@noaa.gov |
| | Combined video index | SEFSC, FWRI | Kevin Thompson Kate Overly Adam Pollack Matt Campbell | kevin.thompson@myfwc.com katherine.overly@noaa.gov adam.pollack@noaa.gov matthew.campbell@noaa.gov |
| | Combined Video Length composition | SEFSC, FWRI | Kevin Thompson Kate Overly Adam Pollack Matt Campbell | kevin.thompson@myfwc.com katherine.overly@noaa.gov adam.pollack@noaa.gov matthew.campbell@noaa.gov |
| | FWRI Age-0 Index | FWRI | Ted Switzer Kerry Walia | ted.switzer@myfwc.com kerry.walia@myfwc.com |
| | FWRI Age-1 Index | FWRI | Ted Switzer Kerry Walia | ted.switzer@myfwc.com kerry.walia@myfwc.com |
| | RFVC Index + size frequency | FWRI | Robert Muller | robert.muller@myfwc.com |